

MATRICES

① Find the eigenvalues & eigenvectors of the matrix $A = \begin{pmatrix} 11 & -4 & -7 \\ 7 & -2 & -5 \\ 10 & -4 & -6 \end{pmatrix}$. [AIM 2018] [N/D 2016] [N/D 2011]

Sol. Given $A = \begin{pmatrix} 11 & -4 & -7 \\ 7 & -2 & -5 \\ 10 & -4 & -6 \end{pmatrix}$

Characteristic equation: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$S_1 =$ Sum of the main diagonal elements $= 11 - 2 - 6 = 3$

$S_2 =$ Sum of the minors of main diagonal elements

$= \begin{vmatrix} -2 & -5 \\ -4 & -6 \end{vmatrix} + \begin{vmatrix} 11 & -7 \\ 10 & -6 \end{vmatrix} + \begin{vmatrix} 11 & -4 \\ 7 & -2 \end{vmatrix} = (12 - 20) + (-66 + 70) + (-22 + 28)$

$= -8 + 4 + 6 = 2$

$S_3 = |A| = 11(12 - 20) + 4(-42 + 50) - 7(-28 + 20) = 11(-8) + 4(8) - 7(-8)$

$= -88 + 32 + 56 = 0$

\therefore The characteristic eqn. is $\lambda^3 - 3\lambda^2 + 2\lambda = 0$

$\lambda(\lambda^2 - 3\lambda + 2) = 0 \Rightarrow \lambda = 0, \lambda^2 - 3\lambda + 2 = 0$
 $(\lambda - 1)(\lambda - 2) = 0$

x	+
2	-3
-1	-2
$\lambda - 1$	$\lambda - 2$

$\therefore \lambda = 0, 1, 2$

Hence the eigenvalues are 0, 1, 2.

Eigenvectors: $(A - \lambda I)x = 0$

$\begin{pmatrix} 11 - \lambda & -4 & -7 \\ 7 & -2 - \lambda & -5 \\ 10 & -4 & -6 - \lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$

$\lambda = 0$ $\begin{pmatrix} 11 & -4 & -7 \\ 7 & -2 & -5 \\ 10 & -4 & -6 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$

x_1	x_2	x_3	
-4	-7	11	-4
-2	-5	7	-2

$11x_1 - 4x_2 - 7x_3 = 0$

$7x_1 - 2x_2 - 5x_3 = 0$

$10x_1 - 4x_2 - 6x_3 = 0$

$\frac{x_1}{20 - 14} = \frac{x_2}{-49 + 55} = \frac{x_3}{-22 + 28} \Rightarrow \frac{x_1}{6} = \frac{x_2}{6} = \frac{x_3}{6} \Rightarrow \frac{x_1}{1} = \frac{x_2}{1} = \frac{x_3}{1}$

$\therefore x_1 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$

$\lambda = 1$ $\begin{pmatrix} 10 & -4 & -7 \\ 7 & -3 & -5 \\ 10 & -4 & -7 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$

$$10x_1 - 4x_2 - 7x_3 = 0$$

$$7x_1 - 3x_2 - 5x_3 = 0$$

$$10x_1 - 4x_2 - 7x_3 = 0$$

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ -4 & -7 & 10 & -4 \\ -3 & -5 & 7 & -3 \end{array}$$

$$\frac{x_1}{20-21} = \frac{x_2}{-49+50} = \frac{x_3}{-30+28} \Rightarrow \frac{x_1}{-1} = \frac{x_2}{1} = \frac{x_3}{-2}$$

$$\therefore x_2 = \begin{pmatrix} -1 \\ 1 \\ -2 \end{pmatrix}$$

$$\underline{\lambda=2} \begin{pmatrix} 9 & -4 & -7 \\ 7 & -4 & -5 \\ 10 & -4 & -8 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ -4 & -7 & 9 & -4 \\ -4 & -5 & 7 & -4 \end{array}$$

$$9x_1 - 4x_2 - 7x_3 = 0$$

$$7x_1 - 4x_2 - 5x_3 = 0$$

$$10x_1 - 4x_2 - 8x_3 = 0$$

$$\frac{x_1}{20-28} = \frac{x_2}{-49+45} = \frac{x_3}{-36+28} \Rightarrow \frac{x_1}{-8} = \frac{x_2}{-4} = \frac{x_3}{-8} \Rightarrow \frac{x_1}{2} = \frac{x_2}{1} = \frac{x_3}{2}$$

$$\therefore x_3 = \begin{pmatrix} 2 \\ 1 \\ 2 \end{pmatrix}$$

② Find the eigenvalues & eigenvectors of the matrix $\begin{pmatrix} -2 & 2 & -3 \\ 2 & 1 & -6 \\ -1 & -2 & 0 \end{pmatrix}$ [M/J-2014]
 [N/D-2014]
 [M/J-2009]
 [Jan-2010]

Sol: Let $A = \begin{pmatrix} -2 & 2 & -3 \\ 2 & 1 & -6 \\ -1 & -2 & 0 \end{pmatrix}$

Characteristic equation: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$S_1 =$ Sum of the main diagonal elements $= -2 + 1 + 0 = -1$

$S_2 =$ Sum of the minors of main diagonal elements

$$= \begin{vmatrix} 1 & -6 \\ -2 & 0 \end{vmatrix} + \begin{vmatrix} -2 & -3 \\ -1 & 0 \end{vmatrix} + \begin{vmatrix} -2 & 2 \\ 2 & 1 \end{vmatrix} = (0-12) + (0-3) + (-2-4)$$

$$= -12 - 3 - 6 = -21$$

$$S_3 = |A| = -2(0-12) - 2(0-6) - 3(-4+1) = 24 + 12 + 9 = 45$$

Hence the characteristic eqn. is $\lambda^3 + \lambda^2 - 21\lambda - 45 = 0$

$$\lambda = 5 \left| \begin{array}{ccc|c} 1 & 1 & -21 & -45 \\ & 5 & 30 & 45 \\ \hline 1 & 6 & 9 & 0 \end{array} \right.$$

$$\begin{array}{r|l} \lambda & x \\ 6 & 9 \\ +3 & 3 \\ \lambda+3 & \lambda+3 \end{array}$$

$$\lambda^2 + 6\lambda + 9 = 0$$

$$(\lambda+3)(\lambda+3) = 0$$

$$\therefore \lambda = 5, -3, -3$$

Hence the eigenvalues are $-3, -3, 5$.

Eigenvectors: $(A - \lambda I)x = 0$

$$\begin{pmatrix} -2-\lambda & 2 & -3 \\ 2 & 1-\lambda & -6 \\ -1 & -2 & -\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\underline{\lambda = -3} \begin{pmatrix} 1 & 2 & -3 \\ 2 & 4 & -6 \\ -1 & -2 & 3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{aligned} x_1 + 2x_2 - 3x_3 &= 0 \\ 2x_1 + 4x_2 - 6x_3 &= 0 \\ -x_1 - 2x_2 + 3x_3 &= 0 \end{aligned}$$

Take $x_1 + 2x_2 - 3x_3 = 0$

Put $x_1 = 0 \Rightarrow 2x_2 - 3x_3 = 0 \Rightarrow 2x_2 = 3x_3 \Rightarrow \frac{x_2}{3} = \frac{x_3}{2}$

$$\therefore x_1 = \begin{pmatrix} 0 \\ 3 \\ 2 \end{pmatrix}$$

Put $x_2 = 0 \rightarrow x_1 - 3x_3 = 0 \Rightarrow x_1 = 3x_3 \Rightarrow \frac{x_1}{3} = \frac{x_3}{1}$

$$\therefore x_2 = \begin{pmatrix} 3 \\ 0 \\ 1 \end{pmatrix}$$

$$\underline{\lambda = 5} \begin{pmatrix} -7 & 2 & -3 \\ 2 & -4 & -6 \\ -1 & -2 & -5 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{aligned} -7x_1 + 2x_2 - 3x_3 &= 0 \\ 2x_1 - 4x_2 - 6x_3 &= 0 \\ -x_1 - 2x_2 - 5x_3 &= 0 \end{aligned}$$

	x_1	x_2	x_3	
	-4	-6	2	-4
	-2	-5	-1	-2

$$\frac{x_1}{20-12} = \frac{x_2}{6+10} = \frac{x_3}{-4-4} \Rightarrow \frac{x_1}{8} = \frac{x_2}{16} = \frac{x_3}{-8} \Rightarrow \frac{x_1}{1} = \frac{x_2}{2} = \frac{x_3}{-1}$$

$$\therefore x_3 = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}$$

③ Find the eigenvalues & eigenvectors of the matrix $\begin{pmatrix} 2 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 2 \end{pmatrix}$ [N/D-2015] [M/J-2013]

Sol: Let $A = \begin{pmatrix} 2 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 2 \end{pmatrix}$

Characteristic equation: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$S_1 =$ Sum of the main diagonal elements $= 2 + 2 + 2 = 6$

$S_2 =$ Sum of the minors of main diagonal elements

$$= \begin{vmatrix} 2 & 0 \\ 0 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 0 \\ 0 & 2 \end{vmatrix} = (4-0) + (4-1) + (4-0) = 4+3+4 = 11$$

$$S_3 = |A| = 2(4-0) - 0(0-0) + 1(0-2) = 8-2 = 6$$

Hence the characteristic equation is $\lambda^3 - 6\lambda^2 + 11\lambda - 6 = 0$

$$\lambda = 1 \left| \begin{array}{ccc|c} 1 & -6 & 11 & -6 \\ & 1 & -5 & 6 \\ \hline 1 & -5 & 6 & 0 \end{array} \right.$$

$$\begin{array}{r|l} x & + \\ 6 & -5 \\ -3 & -2 \\ \lambda-3 & \lambda-2 \end{array}$$

$$\lambda^2 - 5\lambda + 6 = 0$$

$$(\lambda-3)(\lambda-2) = 0$$

$$\therefore \lambda = 1, 2, 3$$

Hence the eigenvalues are 1, 2 & 3.

Eigenvectors: $(A - \lambda I)x = 0$

$$\begin{pmatrix} 2-\lambda & 0 & 1 \\ 0 & 2-\lambda & 0 \\ 1 & 0 & 2-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\underline{\lambda=1} \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{array}$$

$$x_1 + 0x_2 + x_3 = 0$$

$$0x_1 + x_2 + 0x_3 = 0$$

$$x_1 + 0x_2 + x_3 = 0$$

$$\frac{x_1}{0-1} = \frac{x_2}{0-0} = \frac{x_3}{1-0} \Rightarrow \frac{x_1}{-1} = \frac{x_2}{0} = \frac{x_3}{1}$$

$$\therefore x_1 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$$

$$\underline{\lambda=2} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array}$$

$$0x_1 + 0x_2 + x_3 = 0$$

$$0x_1 + 0x_2 + 0x_3 = 0$$

$$x_1 + 0x_2 + 0x_3 = 0$$

$$\frac{x_1}{0-0} = \frac{x_2}{1-0} = \frac{x_3}{0-0} \Rightarrow \frac{x_1}{0} = \frac{x_2}{1} = \frac{x_3}{0}$$

$$\therefore x_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$

$$\underline{\lambda=3} \begin{pmatrix} -1 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{array}{cccc} -x_1 + 0x_2 + x_3 = 0 \\ 0x_1 - x_2 + 0x_3 = 0 \\ x_1 + 0x_2 - x_3 = 0 \end{array}$$

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ 0 & 1 & -1 & 0 \\ -1 & 0 & 0 & -1 \end{array}$$

$$\frac{x_1}{0+1} = \frac{x_2}{0-0} = \frac{x_3}{1-0} \Rightarrow \frac{x_1}{1} = \frac{x_2}{0} = \frac{x_3}{1}$$

$$\therefore x_3 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$$

④ Find the eigenvalues & eigenvectors of the matrix $\begin{pmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{pmatrix}$.

Sol: Let $A = \begin{pmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{pmatrix}$

[A/M-2015]
[N/D-2015]

Characteristic equation: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$S_1 =$ Sum of the main diagonal elements $= 6 + 3 + 3 = 12$

$S_2 =$ Sum of the minors of main diagonal elements

$$= \begin{vmatrix} 3 & -1 \\ -1 & 3 \end{vmatrix} + \begin{vmatrix} 6 & 2 \\ 2 & 3 \end{vmatrix} + \begin{vmatrix} 6 & -2 \\ -2 & 3 \end{vmatrix} = (9-1) + (18-4) + (18-4)$$

$$= 8 + 14 + 14 = 36$$

$$S_3 = |A| = 6(9-1) + 2(-6+2) + 2(2-6) = 6(8) + 2(-4) + 2(-4)$$

$$= 48 - 8 - 8 = 32$$

Hence the characteristic eqn. is $\lambda^3 - 12\lambda^2 + 36\lambda - 32 = 0$

$$\lambda = 2 \quad \left| \begin{array}{ccc|c} 1 & -12 & 36 & -32 \\ & 2 & -20 & 32 \\ & & & 0 \\ \hline 1 & -10 & 16 & 0 \end{array} \right.$$

$$\begin{array}{r|l} x & + \\ 16 & -10 \\ -8 & -2 \\ \hline \lambda - 8 & \lambda - 2 \end{array}$$

$$\lambda^2 - 10\lambda + 16 = 0$$

$$(\lambda - 8)(\lambda - 2) = 0 \Rightarrow \lambda = 8, 2$$

$$\therefore \lambda = 2, 2, 8$$

Hence the eigenvalues are 2, 2, 8.

Eigenvectors: $(A - \lambda I)x = 0$

$$\begin{pmatrix} 6-\lambda & -2 & 2 \\ -2 & 3-\lambda & -1 \\ 2 & -1 & 3-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\underline{\lambda = 2} \quad \begin{pmatrix} 4 & -2 & 2 \\ -2 & 1 & -1 \\ 2 & -1 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\left. \begin{array}{l} 4x_1 - 2x_2 + 2x_3 = 0 \\ -2x_1 + x_2 - x_3 = 0 \\ 2x_1 - x_2 + x_3 = 0 \end{array} \right\} \Rightarrow 2x_1 - x_2 + x_3 = 0$$

Put $x_1 = 0 \Rightarrow -x_2 + x_3 = 0 \Rightarrow x_2 = x_3 \Rightarrow \frac{x_2}{1} = \frac{x_3}{1}$

$$\therefore x_1 = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$$

$$\lambda = 8 \quad \begin{pmatrix} -2 & -2 & 2 \\ -2 & -5 & -1 \\ 2 & -1 & -5 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0 \quad \begin{array}{l} -2x_1 - 2x_2 + 2x_3 = 0 \\ -2x_1 - 5x_2 - x_3 = 0 \\ 2x_1 - x_2 - 5x_3 = 0 \end{array} \quad \begin{array}{cccc} & x_1 & x_2 & x_3 \\ -2 & 2 & -2 & -2 \\ -5 & -1 & -2 & -5 \end{array}$$

$$\frac{x_1}{2+10} = \frac{x_2}{-4-2} = \frac{x_3}{10-4} \Rightarrow \frac{x_1}{12} = \frac{x_2}{-6} = \frac{x_3}{6} \Rightarrow \frac{x_1}{2} = \frac{x_2}{-1} = \frac{x_3}{1}$$

$$\therefore x_2 = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}$$

To find the third eigenvector orthogonal to x_1 & x_2 since the matrix A is symmetric.

Let $x_3 = \begin{pmatrix} a \\ b \\ c \end{pmatrix}$ as x_3 is orthogonal to x_1 & x_2 .

$$x_1^T x_3 = 0 \Rightarrow (0 \ 1 \ 1) \begin{pmatrix} a \\ b \\ c \end{pmatrix} = 0 \Rightarrow 0a + b + c = 0$$

$$\begin{array}{cccc} & a & b & c \\ 1 & 1 & 0 & 1 \\ -1 & 1 & 2 & -1 \end{array}$$

$$x_2^T x_3 = 0 \Rightarrow (2 \ -1 \ 1) \begin{pmatrix} a \\ b \\ c \end{pmatrix} = 0 \Rightarrow 2a - b + c = 0$$

$$\frac{a}{1+1} = \frac{b}{2-0} = \frac{c}{0-2} \Rightarrow \frac{a}{2} = \frac{b}{2} = \frac{c}{-2} \Rightarrow \frac{a}{1} = \frac{b}{1} = \frac{c}{-1}$$

$$\therefore x_3 = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$$

⑤ Find the eigenvalues & eigenvectors of the matrix $\begin{pmatrix} 2 & 2 & 1 \\ 1 & 3 & 1 \\ 1 & 2 & 2 \end{pmatrix}$. [Jan 2014]
[N/D-2010]
[M/J-2010]
[Jan-2012]

Sol: Let $A = \begin{pmatrix} 2 & 2 & 1 \\ 1 & 3 & 1 \\ 1 & 2 & 2 \end{pmatrix}$

Characteristic equation: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$$S_1 = \text{Sum of the main diagonal elements} = 2 + 3 + 2 = 7$$

$S_2 = \text{Sum of the minors of main diagonal elements}$

$$= \begin{vmatrix} 3 & 1 \\ 2 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 2 \\ 1 & 3 \end{vmatrix} = (6-2) + (4-1) + (6-2) = 4 + 3 + 4 = 11$$

$$S_3 = |A| = 2(6-2) - 2(2-1) + 1(2-3) = 8 - 2 - 1 = 5$$

Hence the characteristic eqn. is $\lambda^3 - 7\lambda^2 + 11\lambda - 5 = 0$.

$$\lambda = 1 \quad \left| \begin{array}{ccc|c} 1 & -7 & 11 & -5 \\ & 1 & -6 & 5 \\ \hline 1 & -6 & 5 & 0 \end{array} \right.$$

$$\begin{array}{c|c} x & + \\ \hline 5 & -6 \\ -3 & -2 \\ \hline \lambda - 3 & \lambda - 2 \end{array}$$

$$\lambda^2 - 6\lambda + 5 = 0$$

$$(\lambda - 3)(\lambda - 2) = 0 \Rightarrow \lambda = 2, 3$$

$$\therefore \lambda = 1, 2, 3$$

Hence the eigenvalues are 1, 2, 3.

Eigenvectors: $(A - \lambda I)x = 0$

$$\begin{pmatrix} 2-\lambda & 2 & 1 \\ 1 & 3-\lambda & 1 \\ 1 & 2 & 2-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\underline{\lambda=1} \begin{pmatrix} 1 & 2 & 1 \\ 1 & 2 & 1 \\ 1 & 2 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$x_1 + 2x_2 + x_3 = 0$$

$$x_1 + 2x_2 + x_3 = 0$$

$$x_1 + 2x_2 + x_3 = 0$$

Put $x_1 = 0 \Rightarrow 2x_2 + x_3 = 0 \Rightarrow 2x_2 = -x_3 \Rightarrow \frac{x_2}{-1} = \frac{x_3}{2}$

$$\therefore x_1 = \begin{pmatrix} 0 \\ -1 \\ 2 \end{pmatrix}$$

x_1	x_2	x_3
2	1	0
1	1	1

$$\underline{\lambda=2} \begin{pmatrix} 0 & 2 & 1 \\ 1 & 1 & 1 \\ 1 & 2 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$0x_1 + 2x_2 + x_3 = 0$$

$$x_1 + x_2 + x_3 = 0$$

$$x_1 + 2x_2 + 0x_3 = 0$$

$$\frac{x_1}{2-1} = \frac{x_2}{1-0} = \frac{x_3}{0-2} \Rightarrow \frac{x_1}{1} = \frac{x_2}{1} = \frac{x_3}{-2}$$

$$\therefore x_2 = \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix}$$

x_1	x_2	x_3
2	1	-1
0	1	1

$$\underline{\lambda=3} \begin{pmatrix} -1 & 2 & 1 \\ 1 & 0 & 1 \\ 1 & 2 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$-x_1 + 2x_2 + x_3 = 0$$

$$x_1 + 0x_2 + x_3 = 0$$

$$x_1 + 2x_2 - x_3 = 0$$

$$\frac{x_1}{2-0} = \frac{x_2}{1+1} = \frac{x_3}{0-2} \Rightarrow \frac{x_1}{2} = \frac{x_2}{2} = \frac{x_3}{-2} \Rightarrow \frac{x_1}{1} = \frac{x_2}{1} = \frac{x_3}{-1}$$

$$\therefore x_3 = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$$

Cayley-Hamilton Theorem:

Every square matrix satisfies its own characteristic equation.

Uses of Cayley-Hamilton Theorem:

- To calculate (i) the positive integral powers of A &
- (ii) the inverse of a non-singular square matrix A.

⑥ Verify Cayley-Hamilton thm. for the matrix $A = \begin{pmatrix} 2 & -1 & 2 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{pmatrix}$. Hence using it find A^{-1} & A^4 . [N/D-2014] [A/M-2017] [M/J-2013] [M/J-2010]

Sol: Given $A = \begin{pmatrix} 2 & -1 & 2 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{pmatrix}$.

Characteristic equation: $\lambda^3 - S_1 \lambda^2 + S_2 \lambda - S_3 = 0$

$S_1 =$ Sum of the main diagonal elements $= 2+2+2=6$

$S_2 =$ Sum of the minors of main diagonal elements

$$= \begin{vmatrix} 2 & -1 \\ -1 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 2 \\ 1 & 2 \end{vmatrix} + \begin{vmatrix} 2 & -1 \\ -1 & 2 \end{vmatrix} = (4-1) + (4-2) + (4-1) = 3+2+3=8$$

$$S_3 = |A| = 2(4-1) + 1(-2+1) + 2(1-2) = 2(3) + 1(-1) + 2(-1) = 6-1-2=3$$

Hence the characteristic eqn. is $\lambda^3 - 6\lambda^2 + 8\lambda - 3 = 0$.

By Cayley-Hamilton thm., every square matrix satisfies its own characteristic equation. $\therefore A^3 - 6A^2 + 8A - 3I = 0$ — (1)

Verification:

$$A^2 = \begin{pmatrix} 2 & -1 & 2 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{pmatrix} \begin{pmatrix} 2 & -1 & 2 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{pmatrix} = \begin{pmatrix} 7 & -6 & 9 \\ -5 & 6 & -6 \\ 5 & -5 & 7 \end{pmatrix}$$

$$A^3 = \begin{pmatrix} 7 & -6 & 9 \\ -5 & 6 & -6 \\ 5 & -5 & 7 \end{pmatrix} \begin{pmatrix} 2 & -1 & 2 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{pmatrix} = \begin{pmatrix} 29 & -28 & 38 \\ -22 & 23 & -28 \\ 22 & -22 & 29 \end{pmatrix}$$

$$A^3 - 6A^2 + 8A - 3I = \begin{pmatrix} 29 & -28 & 38 \\ -22 & 23 & -28 \\ 22 & -22 & 29 \end{pmatrix} - 6 \begin{pmatrix} 7 & -6 & 9 \\ -5 & 6 & -6 \\ 5 & -5 & 7 \end{pmatrix} + 8 \begin{pmatrix} 2 & -1 & 2 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{pmatrix} - 3 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 29 & -28 & 38 \\ -22 & 23 & -28 \\ 22 & -22 & 29 \end{pmatrix} - \begin{pmatrix} 42 & -36 & 54 \\ -30 & 36 & -36 \\ 30 & -30 & 42 \end{pmatrix} + \begin{pmatrix} 16 & -8 & 16 \\ -8 & 16 & -8 \\ 8 & -8 & 16 \end{pmatrix} - \begin{pmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{pmatrix}$$

$$= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

Hence Cayley-Hamilton thm. verified.

$$\textcircled{1} \Rightarrow A^3 - 6A^2 + 8A - 3I = 0 \Rightarrow A^2 - 6A + 8I - 3A^{-1} = 0 \Rightarrow 3A^{-1} = A^2 - 6A + 8I$$

$$\therefore A^{-1} = \frac{1}{3} (A^2 - 6A + 8I)$$

$$A^2 - 6A + 8I = \begin{pmatrix} 7 & -6 & 9 \\ -5 & 6 & -6 \\ 5 & -5 & 7 \end{pmatrix} - 6 \begin{pmatrix} 2 & -1 & 2 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{pmatrix} + 8 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 7 & -6 & 9 \\ -5 & 6 & -6 \\ 5 & -5 & 7 \end{pmatrix} - \begin{pmatrix} 12 & -6 & 12 \\ -6 & 12 & -6 \\ 6 & -6 & 12 \end{pmatrix} + \begin{pmatrix} 8 & 0 & 0 \\ 0 & 8 & 0 \\ 0 & 0 & 8 \end{pmatrix} = \begin{pmatrix} 3 & 0 & -3 \\ 1 & 2 & 0 \\ -1 & 1 & 3 \end{pmatrix}$$

$$\therefore A^{-1} = \frac{1}{3} \begin{pmatrix} 3 & 0 & -3 \\ 1 & 2 & 0 \\ -1 & 1 & 3 \end{pmatrix} = \begin{pmatrix} 1 & 0 & -1 \\ \frac{1}{3} & \frac{2}{3} & 0 \\ -\frac{1}{3} & \frac{1}{3} & 1 \end{pmatrix}$$

$$\textcircled{1} \Rightarrow A^3 - 6A^2 + 8A - 3I = 0 \Rightarrow A^4 - 6A^3 + 8A^2 - 3A = 0 \Rightarrow A^4 = 6A^3 - 8A^2 + 3A$$

$$A^4 = 6 \begin{pmatrix} 29 & -28 & 38 \\ -22 & 23 & -28 \\ 22 & -22 & 29 \end{pmatrix} - 8 \begin{pmatrix} 7 & -6 & 9 \\ -5 & 6 & -6 \\ 5 & -5 & 7 \end{pmatrix} + 3 \begin{pmatrix} 2 & -1 & 2 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{pmatrix}$$

$$= \begin{pmatrix} 174 & -168 & 228 \\ -132 & 138 & -168 \\ 132 & -132 & 174 \end{pmatrix} - \begin{pmatrix} 56 & -48 & 72 \\ -40 & 48 & -48 \\ 40 & -40 & 56 \end{pmatrix} + \begin{pmatrix} 6 & -3 & 6 \\ -3 & 6 & -3 \\ 3 & -3 & 6 \end{pmatrix} = \begin{pmatrix} 124 & -123 & 162 \\ -95 & 96 & -123 \\ 95 & -95 & 124 \end{pmatrix}$$

$\textcircled{7}$ Verify Cayley-Hamilton thm. for the matrix $A = \begin{pmatrix} 1 & 2 & -2 \\ 2 & 5 & -4 \\ 3 & 7 & -5 \end{pmatrix}$. Hence using it find A^{-1} . [N/D-2015]

Sol: Given $A = \begin{pmatrix} 1 & 2 & -2 \\ 2 & 5 & -4 \\ 3 & 7 & -5 \end{pmatrix}$

Characteristic eqn.: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$S_1 = \text{Sum of the main diagonal elements} = 1 + 5 - 5 = 1$

$S_2 = \text{Sum of the minors of main diagonal elements}$

$$= \begin{vmatrix} 5 & -4 \\ 7 & -5 \end{vmatrix} + \begin{vmatrix} 1 & -2 \\ 3 & -5 \end{vmatrix} + \begin{vmatrix} 1 & 2 \\ 2 & 5 \end{vmatrix} = (-25 + 28) + (-5 + 6) + (5 - 4)$$

$$= 3 + 1 + 1 = 5$$

$$S_3 = |A| = 1(-25 + 28) - 2(-10 + 12) - 2(14 - 15) = 1(3) - 2(2) - 2(-1)$$

$$= 3 - 4 + 2 = 1$$

Hence the characteristic eqn. is $\lambda^3 - \lambda^2 + 5\lambda - 1 = 0$

Verification: By C-H thm., every square matrix satisfies its own characteristic eqn. $\therefore A^3 - A^2 + 5A - I = 0$ — $\textcircled{1}$

$$A^2 = \begin{pmatrix} -1 & -2 & 0 \\ 0 & 1 & -4 \\ 2 & 6 & -9 \end{pmatrix} \quad A^3 = \begin{pmatrix} -5 & -12 & 10 \\ -10 & -23 & 16 \\ -13 & -29 & 17 \end{pmatrix}$$

$$\therefore A^3 - A^2 + 5A - I = \begin{pmatrix} -5 & -12 & 10 \\ -10 & -23 & 16 \\ -13 & -29 & 17 \end{pmatrix} - \begin{pmatrix} -1 & -2 & 0 \\ 0 & 1 & -4 \\ 2 & 6 & -9 \end{pmatrix} + 5 \begin{pmatrix} 1 & 2 & -2 \\ 2 & 5 & -4 \\ 3 & 7 & -5 \end{pmatrix} - \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} -5 & -12 & 10 \\ -10 & -23 & 16 \\ -13 & -29 & 17 \end{pmatrix} + \begin{pmatrix} 1 & 2 & 0 \\ 0 & -1 & 4 \\ -2 & -6 & 9 \end{pmatrix} + \begin{pmatrix} 5 & 10 & -10 \\ 10 & 25 & -20 \\ 15 & 35 & -25 \end{pmatrix} + \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

$$= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

Hence C-H thm. verified.

$$\textcircled{1} \Rightarrow A^3 - A^2 + 5A - I = 0 \Rightarrow A^2 - A + 5I - A^{-1} = 0 \Rightarrow A^{-1} = A^2 - A + 5I$$

$$\therefore A^{-1} = A^2 - A + 5I = \begin{pmatrix} -1 & -2 & 0 \\ 0 & 1 & -4 \\ 2 & 6 & -9 \end{pmatrix} - \begin{pmatrix} 1 & 2 & -2 \\ 2 & 5 & -4 \\ 3 & 7 & -5 \end{pmatrix} + 5 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} -1 & -2 & 0 \\ 0 & 1 & -4 \\ 2 & 6 & -9 \end{pmatrix} + \begin{pmatrix} -1 & -2 & 2 \\ -2 & -5 & 4 \\ -3 & -7 & 5 \end{pmatrix} + \begin{pmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{pmatrix}$$

$$\therefore A^{-1} = \begin{pmatrix} 3 & -4 & 2 \\ -2 & 1 & 0 \\ -1 & -1 & 1 \end{pmatrix}$$

⑧ Verify C-H thm. for the matrix $A = \begin{pmatrix} 1 & 3 & 7 \\ 4 & 2 & 3 \\ 1 & 2 & 1 \end{pmatrix}$. Hence using it find A^{-1} .

Sol: Given $A = \begin{pmatrix} 1 & 3 & 7 \\ 4 & 2 & 3 \\ 1 & 2 & 1 \end{pmatrix}$

[A/M-2015]

[N/D-2011]

Characteristic eqn.: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$S_1 =$ Sum of the main diagonal elements $= 1 + 2 + 1 = 4$

$S_2 =$ Sum of the minors of main diagonal elements

$$= \begin{vmatrix} 2 & 3 \\ 2 & 1 \end{vmatrix} + \begin{vmatrix} 1 & 7 \\ 1 & 1 \end{vmatrix} + \begin{vmatrix} 1 & 3 \\ 4 & 2 \end{vmatrix} = (2-6) + (1-7) + (2-12) = -4-6-10 = -20$$

$$S_3 = |A| = 1(2-6) - 3(4-3) + 7(8-2) = -4 - 3(1) + 7(6) = -4 - 3 + 42 = 35$$

Hence the characteristic eqn. is $\lambda^3 - 4\lambda^2 - 20\lambda - 35 = 0$

By C-H thm., every square matrix satisfies its own characteristic eqn.

$$\therefore A^3 - 4A^2 - 20A - 35I = 0 \text{ --- } \textcircled{1}$$

Verification:

$$A^2 = \begin{pmatrix} 20 & 23 & 23 \\ 15 & 22 & 37 \\ 10 & 9 & 14 \end{pmatrix}$$

$$A^3 = \begin{pmatrix} 135 & 152 & 232 \\ 140 & 163 & 208 \\ 60 & 76 & 111 \end{pmatrix}$$

$$A^3 - 4A^2 - 20A - 35I = \begin{pmatrix} 135 & 152 & 232 \\ 140 & 163 & 208 \\ 60 & 76 & 111 \end{pmatrix} - 4 \begin{pmatrix} 20 & 23 & 23 \\ 15 & 22 & 37 \\ 10 & 9 & 14 \end{pmatrix} - 20 \begin{pmatrix} 1 & 3 & 7 \\ 4 & 2 & 3 \\ 1 & 2 & 1 \end{pmatrix} - 35 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 135 & 152 & 232 \\ 140 & 163 & 208 \\ 60 & 76 & 111 \end{pmatrix} - \begin{pmatrix} 80 & 92 & 92 \\ 60 & 88 & 148 \\ 40 & 36 & 56 \end{pmatrix} - \begin{pmatrix} 20 & 60 & 140 \\ 80 & 40 & 60 \\ 20 & 40 & 20 \end{pmatrix} - \begin{pmatrix} 35 & 0 & 0 \\ 0 & 35 & 0 \\ 0 & 0 & 35 \end{pmatrix}$$

$$= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \text{ Hence Cayley-Hamilton thm. verified.}$$

$$\textcircled{1} \Rightarrow A^3 - 4A^2 - 20A - 35I = 0 \Rightarrow A^2 - 4A - 20I - 35A^{-1} = 0$$

$$\Rightarrow 35A^{-1} = A^2 - 4A - 20I \Rightarrow A^{-1} = \frac{1}{35}(A^2 - 4A - 20I)$$

$$\therefore A^2 - 4A - 20I = \begin{pmatrix} 20 & 23 & 23 \\ 15 & 22 & 37 \\ 10 & 9 & 14 \end{pmatrix} - 4 \begin{pmatrix} 1 & 3 & 7 \\ 4 & 2 & 3 \\ 1 & 2 & 1 \end{pmatrix} - 20 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 20 & 23 & 23 \\ 15 & 22 & 37 \\ 10 & 9 & 14 \end{pmatrix} - \begin{pmatrix} 4 & 12 & 28 \\ 16 & 8 & 12 \\ 4 & 8 & 4 \end{pmatrix} - \begin{pmatrix} 20 & 0 & 0 \\ 0 & 20 & 0 \\ 0 & 0 & 20 \end{pmatrix} = \begin{pmatrix} -4 & 11 & -5 \\ -1 & -6 & 25 \\ 6 & 1 & -10 \end{pmatrix}$$

$$\therefore A^{-1} = \frac{1}{35} \begin{pmatrix} -4 & 11 & -5 \\ -1 & -6 & 25 \\ 6 & 1 & -10 \end{pmatrix}$$

9) Verify Cayley-Hamilton thm. for $A = \begin{pmatrix} 1 & -1 & 1 \\ 0 & 1 & 0 \\ 2 & 0 & 3 \end{pmatrix}$. Hence using it find A^{-1} & A^4 [Jan-2011]

Sol: Given $A = \begin{pmatrix} 1 & -1 & 1 \\ 0 & 1 & 0 \\ 2 & 0 & 3 \end{pmatrix}$

Characteristic eqn.: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$S_1 =$ Sum of the main diagonal elements $= 1 + 1 + 3 = 5$

$S_2 =$ Sum of the minors of main diagonal elements
 $= \begin{vmatrix} 1 & 0 \\ 0 & 3 \end{vmatrix} + \begin{vmatrix} 1 & 1 \\ 2 & 3 \end{vmatrix} + \begin{vmatrix} 1 & -1 \\ 0 & 1 \end{vmatrix} = (3-0) + (3-2) + (1-0) = 3+1+1 = 5$

$S_3 = |A| = 1(3-0) + 1(0-2) + 1(0-2) = 3-2 = 1$

Hence the characteristic eqn. is $\lambda^3 - 5\lambda^2 + 5\lambda - 1 = 0$.

By C-H thm., every square matrix satisfies its own characteristic eqn.

$$\therefore A^3 - 5A^2 + 5A - I = 0 \text{ --- } \textcircled{1}$$

Verification:

$$A^2 = \begin{pmatrix} 3 & -2 & 4 \\ 0 & 1 & 0 \\ 8 & -2 & 11 \end{pmatrix}, \quad A^3 = \begin{pmatrix} 11 & -5 & 15 \\ 0 & 1 & 0 \\ 30 & -10 & 41 \end{pmatrix}$$

$$A^3 - 5A^2 + 5A - I = \begin{pmatrix} 11 & -5 & 15 \\ 0 & 1 & 0 \\ 30 & -10 & 41 \end{pmatrix} - 5 \begin{pmatrix} 3 & -2 & 4 \\ 0 & 1 & 0 \\ 8 & -2 & 11 \end{pmatrix} + 5 \begin{pmatrix} 1 & -1 & 1 \\ 0 & 1 & 0 \\ 2 & 0 & 3 \end{pmatrix} - \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 11 & -5 & 15 \\ 0 & 1 & 0 \\ 30 & -10 & 41 \end{pmatrix} - \begin{pmatrix} 15 & -10 & 20 \\ 0 & 5 & 0 \\ 40 & -10 & 55 \end{pmatrix} + \begin{pmatrix} 5 & -5 & 5 \\ 0 & 5 & 0 \\ 10 & 0 & 15 \end{pmatrix} - \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

Hence Cayley-Hamilton thm. verified.

$$\textcircled{1} \Rightarrow A^3 - 5A^2 + 5A - I = 0 \Rightarrow A^2 - 5A + 5I - A^{-1} = 0 \Rightarrow A^{-1} = A^2 - 5A + 5I$$

$$A^{-1} = A^2 - 5A + 5I = \begin{pmatrix} 3 & -2 & 4 \\ 0 & 1 & 0 \\ 8 & -2 & 11 \end{pmatrix} - 5 \begin{pmatrix} 1 & -1 & 1 \\ 0 & 1 & 0 \\ 2 & 0 & 3 \end{pmatrix} + 5 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 3 & -2 & 4 \\ 0 & 1 & 0 \\ 8 & -2 & 11 \end{pmatrix} - \begin{pmatrix} 5 & -5 & 5 \\ 0 & 5 & 0 \\ 10 & 0 & 15 \end{pmatrix} + \begin{pmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{pmatrix}$$

$$\therefore A^{-1} = \begin{pmatrix} 3 & 3 & -1 \\ 0 & 1 & 0 \\ -2 & -2 & 1 \end{pmatrix}$$

$$\text{From } \textcircled{1}, A^4 - 5A^3 + 5A^2 - A = 0 \Rightarrow A^4 = 5A^3 - 5A^2 + A$$

$$A^4 = 5A^3 - 5A^2 + A = 5 \begin{pmatrix} 11 & -5 & 15 \\ 0 & 1 & 0 \\ 30 & -10 & 41 \end{pmatrix} - 5 \begin{pmatrix} 3 & -2 & 4 \\ 0 & 1 & 0 \\ 8 & -2 & 11 \end{pmatrix} + \begin{pmatrix} 1 & -1 & 1 \\ 0 & 1 & 0 \\ 2 & 0 & 3 \end{pmatrix}$$

$$= \begin{pmatrix} 55 & -25 & 75 \\ 0 & 5 & 0 \\ 150 & -50 & 205 \end{pmatrix} - \begin{pmatrix} 15 & -10 & 20 \\ 0 & 5 & 0 \\ 40 & -10 & 55 \end{pmatrix} + \begin{pmatrix} 1 & -1 & 1 \\ 0 & 1 & 0 \\ 2 & 0 & 3 \end{pmatrix}$$

$$\therefore A^4 = \begin{pmatrix} 41 & -16 & 56 \\ 0 & 1 & 0 \\ 112 & -40 & 153 \end{pmatrix}$$

\textcircled{10} Using Cayley-Hamilton Thm. find the inverse of the given matrix

$$A = \begin{pmatrix} 1 & 2 & 1 \\ 2 & 2 & 1 \\ 1 & 1 & 3 \end{pmatrix} \quad [A/M 2018]$$

Sol: Given $A = \begin{pmatrix} 1 & 2 & 1 \\ 2 & 2 & 1 \\ 1 & 1 & 3 \end{pmatrix}$

Characteristic eqn: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$S_1 =$ Sum of the main diagonal elements $= 1 + 2 + 3 = 6$

$S_2 =$ Sum of the minors of main diagonal elements

$$= \begin{vmatrix} 2 & 1 \\ 1 & 3 \end{vmatrix} + \begin{vmatrix} 1 & 1 \\ 2 & 2 \end{vmatrix} + \begin{vmatrix} 1 & 2 \\ 2 & 2 \end{vmatrix} = (6-1) + (3-1) + (2-4) = 5 + 2 - 2 = 5$$

$$S_3 = |A| = 1(6-1) - 2(6-1) + 1(2-2) = 5 - 10 = -5$$

Hence the characteristic eqn. is $\lambda^3 - 6\lambda^2 + 5\lambda + 5 = 0$

Using C-H thm., we get $A^3 - 6A^2 + 5A + 5I = 0$

$$A^3 - 6A^2 + 5I + 5A^{-1} = 0 \Rightarrow 5A^{-1} = -A^3 + 6A^2 - 5I \Rightarrow A^{-1} = \frac{1}{5}(-A^3 + 6A^2 - 5I)$$

$$\therefore A^{-1} = \frac{1}{5} \left[- \begin{pmatrix} 6 & 7 & 6 \\ 7 & 9 & 7 \\ 6 & 7 & 11 \end{pmatrix} + 6 \begin{pmatrix} 1 & 2 & 1 \\ 2 & 2 & 1 \\ 1 & 1 & 3 \end{pmatrix} - 5 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right]$$

$$A^{-1} = \frac{1}{5} \left[\begin{pmatrix} -6 & -7 & -6 \\ -7 & -9 & -7 \\ -6 & -7 & -11 \end{pmatrix} + \begin{pmatrix} 6 & 12 & 6 \\ 12 & 12 & 6 \\ 6 & 6 & 18 \end{pmatrix} + \begin{pmatrix} -5 & 0 & 0 \\ 0 & -5 & 0 \\ 0 & 0 & -5 \end{pmatrix} \right]$$

$$\therefore A^{-1} = \frac{1}{5} \begin{pmatrix} -5 & 5 & 0 \\ 5 & -2 & -1 \\ 0 & -1 & 2 \end{pmatrix}$$

Q11) Use C-H thm. to find the value of the matrix given by

$$A^8 - 5A^7 + 7A^6 - 3A^5 + A^4 - 5A^3 + 8A^2 - 2A + I, \text{ if the matrix } A = \begin{pmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{pmatrix}.$$

[M/J-2009]

Sol: Given $A = \begin{pmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{pmatrix}$

Characteristic eqn.: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$S_1 =$ Sum of the main diagonal elements $= 2 + 1 + 2 = 5$

$S_2 =$ Sum of the minors of main diagonal elements

$$= \begin{vmatrix} 1 & 0 \\ 1 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ 0 & 1 \end{vmatrix} = (2-0) + (4-1) + (2-0) = 2+3+2 = 7$$

$S_3 = |A| = 2(2-0) - 1(0-0) + 1(0-1) = 4 - 1 = 3$

Hence the characteristic eqn. is $\lambda^3 - 5\lambda^2 + 7\lambda - 3 = 0$

Using C-H thm. we get, $A^3 - 5A^2 + 7A - 3I = 0$ — (1)

$A^3 - 5A^2 + 7A - 3I$	$A^8 - 5A^7 + 7A^6 - 3A^5 + A^4 - 5A^3 + 8A^2 - 2A + I$
$A^8 - 5A^7 + 7A^6 - 3A^5$	$A^8 - 5A^7 + 7A^6 - 3A^5$
$(-) (+) \quad (-) \quad (+)$	$A^4 - 5A^3 + 8A^2 - 2A$
	$A^4 - 5A^3 + 7A^2 - 3A$
	$(-) (+) \quad (-) \quad (+)$
	$A^2 + A + I$

$$\therefore A^8 - 5A^7 + 7A^6 - 3A^5 + A^4 - 5A^3 + 8A^2 - 2A + I = (A^3 - 5A^2 + 7A - 3I)(A^5 + A) + A^2 + A + I$$

$$= (0)(A^5 + A) + A^2 + A + I \quad (\because \text{by (1)})$$

$$= A^2 + A + I$$

$$= \begin{pmatrix} 5 & 1 & 1 \\ 0 & 1 & 0 \\ 4 & 1 & 5 \end{pmatrix} + \begin{pmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{pmatrix} + \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 8 & 5 & 5 \\ 0 & 3 & 0 \\ 5 & 5 & 8 \end{pmatrix}$$

⑫ Find A^n using c-H thm., taking $A = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$. Hence find A^3 .

[Jan-2012]

Sol. Given $A = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$

Characteristic eqn.: $\lambda^2 - S_1\lambda + S_2 = 0$

$S_1 =$ Sum of the main diagonal elements $= 1+3=4$

$S_2 = |A| = 3-8 = -5$

Hence the characteristic eqn. is $\lambda^2 - 4\lambda - 5 = 0$

Using c-H thm., we get, $A^2 - 4A - 5I = 0$. — ①

$A^n = (A^2 - 4A - 5I)Q(A) + aA + bI$ where $Q(A)$ is the quotient & $aA + bI$ is the remainder.

$\therefore A^n = (0)Q(A) + aA + bI$ (\because by ①)

$A^n = aA + bI \Rightarrow \lambda^n = a\lambda + b$ — ②

Eigenvalues: $\lambda^2 - 4\lambda - 5 = 0$
 $(\lambda+1)(\lambda-5) = 0$
 $\therefore \lambda = -1, 5$

	λ
$+$	-5
$-$	-4
$+$	-5
$\lambda+1$	$\lambda-5$

Subst. $\lambda = -1$ & 5 in ②,

$(-1)^n = a(-1) + b \Rightarrow (-1)^n = -a + b$ — ③

$5^n = a(5) + b \Rightarrow 5^n = 5a + b$ — ④

③ - ④ $\Rightarrow (-1)^n - 5^n = -6a \Rightarrow a = \frac{-1}{6} [(-1)^n - 5^n] = \frac{1}{6} [5^n - (-1)^n]$

Subst. a value in ③, $(-1)^n = -\frac{1}{6} [5^n - (-1)^n] + b$

$\Rightarrow b = (-1)^n + \frac{1}{6} [5^n - (-1)^n] = \frac{6(-1)^n + 5^n - (-1)^n}{6} = \frac{5(-1)^n + 5^n}{6}$

$\therefore b = \frac{1}{6} [5(-1)^n + 5^n]$

$\therefore A^n = \frac{1}{6} [5^n - (-1)^n] A + \frac{1}{6} [5(-1)^n + 5^n] I$

$A^3 = \frac{1}{6} [5^3 - (-1)^3] A + \frac{1}{6} [5(-1)^3 + 5^3] I$

$= \frac{1}{6} [125 + 1] A + \frac{1}{6} [-5 + 125] I = \frac{126}{6} A + \frac{120}{6} I$

$= 21A + 20I = 21 \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} + 20 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 21 & 84 \\ 42 & 63 \end{pmatrix} + \begin{pmatrix} 20 & 0 \\ 0 & 20 \end{pmatrix} = \begin{pmatrix} 41 & 84 \\ 42 & 83 \end{pmatrix}$

13) Reduce the matrix $\begin{pmatrix} 10 & -2 & -5 \\ -2 & 2 & 3 \\ -5 & 3 & 5 \end{pmatrix}$ to diagonal form. [A/M-2017]

Sol:

Let $A = \begin{pmatrix} 10 & -2 & -5 \\ -2 & 2 & 3 \\ -5 & 3 & 5 \end{pmatrix}$

Characteristic eqn/: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$S_1 =$ Sum of the main diagonal elements $= 10 + 2 + 5 = 17$

$S_2 =$ Sum of the minors of main diagonal elements

$= \begin{vmatrix} 2 & 3 \\ 3 & 5 \end{vmatrix} + \begin{vmatrix} 10 & -5 \\ -5 & 5 \end{vmatrix} + \begin{vmatrix} 10 & -2 \\ -2 & 2 \end{vmatrix} = (10-9) + (50-25) + (20-4) = 1+25+16 = 42$

$S_3 = |A| = 10(10-9) + 2(-10+15) - 5(-6+10) = 10+10-20 = 0$

Hence the characteristic eqn/ is $\lambda^3 - 17\lambda^2 + 42\lambda = 0$
 $\lambda(\lambda^2 - 17\lambda + 42) = 0$

$\lambda = 0, \lambda^2 - 17\lambda + 42 = 0$
 $(\lambda - 14)(\lambda - 3) = 0$

$$\begin{array}{r|l} x & + \\ 42 & -17 \\ -14 & -3 \\ \lambda - 14 & \lambda - 3 \end{array}$$

$\therefore \lambda = 0, 3, 14$

Hence the eigenvalues are 0, 3 & 14.

Eigenvectors: $(A - \lambda I)x = 0$

$\begin{pmatrix} 10-\lambda & -2 & -5 \\ -2 & 2-\lambda & 3 \\ -5 & 3 & 5-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$

$\lambda = 0 \begin{pmatrix} 10 & -2 & -5 \\ -2 & 2 & 3 \\ -5 & 3 & 5 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$

$10x_1 - 2x_2 - 5x_3 = 0$
 $-2x_1 + 2x_2 + 3x_3 = 0$
 $-5x_1 + 3x_2 + 5x_3 = 0$

$$\begin{array}{ccc|ccc} x_1 & x_2 & x_3 & & & \\ -2 & -5 & 10 & & -2 & \\ 2 & 3 & -2 & & 2 & \end{array}$$

$\frac{x_1}{-6+10} = \frac{x_2}{10-20} = \frac{x_3}{20-4} \Rightarrow \frac{x_1}{4} = \frac{x_2}{-20} = \frac{x_3}{16} \Rightarrow \frac{x_1}{1} = \frac{x_2}{-5} = \frac{x_3}{4}$

$\therefore x_1 = \begin{pmatrix} 1 \\ -5 \\ 4 \end{pmatrix}$

$\lambda = 3 \begin{pmatrix} 7 & -2 & -5 \\ -2 & -1 & 3 \\ -5 & 3 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$

$7x_1 - 2x_2 - 5x_3 = 0$
 $-2x_1 - x_2 + 3x_3 = 0$
 $-5x_1 + 3x_2 + 2x_3 = 0$

$$\begin{array}{ccc|ccc} x_1 & x_2 & x_3 & & & \\ -2 & -5 & 7 & & -2 & \\ -1 & 3 & -2 & & -1 & \end{array}$$

$\frac{x_1}{-6-5} = \frac{x_2}{10-21} = \frac{x_3}{-7-4} \Rightarrow \frac{x_1}{-11} = \frac{x_2}{-11} = \frac{x_3}{-11} \Rightarrow \frac{x_1}{1} = \frac{x_2}{1} = \frac{x_3}{1}$

$\therefore x_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$

$$\lambda = 14 \quad \begin{pmatrix} -4 & -2 & -5 \\ -2 & -12 & 3 \\ -5 & 3 & -9 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0 \quad \begin{array}{l} -4x_1 - 2x_2 - 5x_3 = 0 \\ -2x_1 - 12x_2 + 3x_3 = 0 \\ -5x_1 + 3x_2 - 9x_3 = 0 \end{array} \quad \begin{array}{ccc} x_1 & x_2 & x_3 \\ -2 & -5 & -4 \\ -12 & 3 & -2 \\ -5 & -2 & -9 \end{array}$$

$$\frac{x_1}{-6-60} = \frac{x_2}{10+12} = \frac{x_3}{48-4} \Rightarrow \frac{x_1}{-66} = \frac{x_2}{22} = \frac{x_3}{44} \Rightarrow \frac{x_1}{-6} = \frac{x_2}{2} = \frac{x_3}{4}$$

$$\Rightarrow \frac{x_1}{-3} = \frac{x_2}{1} = \frac{x_3}{2} \quad \therefore x_3 = \begin{pmatrix} -3 \\ 1 \\ 2 \end{pmatrix}$$

$$x_1^T x_2 = (1 \ -5 \ 4) \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = 1 - 5 + 4 = 0$$

$$x_1^T x_3 = (1 \ -5 \ 4) \begin{pmatrix} -3 \\ 1 \\ 2 \end{pmatrix} = -3 - 5 + 8 = 0$$

$$x_2^T x_3 = (1 \ 1 \ 1) \begin{pmatrix} -3 \\ 1 \\ 2 \end{pmatrix} = -3 + 1 + 2 = 0$$

Hence the eigenvectors are orthogonal to each other.

$$N = \begin{pmatrix} \frac{1}{\sqrt{42}} & \frac{1}{\sqrt{3}} & -\frac{3}{\sqrt{14}} \\ -\frac{5}{\sqrt{42}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{14}} \\ \frac{4}{\sqrt{42}} & \frac{1}{\sqrt{3}} & \frac{2}{\sqrt{14}} \end{pmatrix} \quad N^T = \begin{pmatrix} \frac{1}{\sqrt{42}} & -\frac{5}{\sqrt{42}} & \frac{4}{\sqrt{42}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \\ -\frac{3}{\sqrt{14}} & \frac{1}{\sqrt{14}} & \frac{2}{\sqrt{14}} \end{pmatrix}$$

$$AN = \begin{pmatrix} 10 & -2 & -5 \\ -2 & 2 & 3 \\ -5 & 3 & 5 \end{pmatrix} \begin{pmatrix} \frac{1}{\sqrt{42}} & \frac{1}{\sqrt{3}} & -\frac{3}{\sqrt{14}} \\ -\frac{5}{\sqrt{42}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{14}} \\ \frac{4}{\sqrt{42}} & \frac{1}{\sqrt{3}} & \frac{2}{\sqrt{14}} \end{pmatrix}$$

$$= \begin{pmatrix} 0 & \frac{3}{\sqrt{3}} & -\frac{42}{\sqrt{14}} \\ 0 & \frac{3}{\sqrt{3}} & \frac{14}{\sqrt{14}} \\ 0 & \frac{3}{\sqrt{3}} & \frac{28}{\sqrt{14}} \end{pmatrix}$$

$$D = N^T AN = \begin{pmatrix} \frac{1}{\sqrt{42}} & -\frac{5}{\sqrt{42}} & \frac{4}{\sqrt{42}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \\ -\frac{3}{\sqrt{14}} & \frac{1}{\sqrt{14}} & \frac{2}{\sqrt{14}} \end{pmatrix} \begin{pmatrix} 0 & \frac{3}{\sqrt{3}} & -\frac{42}{\sqrt{14}} \\ 0 & \frac{3}{\sqrt{3}} & \frac{14}{\sqrt{14}} \\ 0 & \frac{3}{\sqrt{3}} & \frac{28}{\sqrt{14}} \end{pmatrix}$$

$$= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 14 \end{pmatrix}$$

9) Diagonalize the matrix $A = \begin{pmatrix} 2 & 0 & 4 \\ 0 & 6 & 0 \\ 4 & 0 & 2 \end{pmatrix}$. [M/J-2014]

Sol: Given $A = \begin{pmatrix} 2 & 0 & 4 \\ 0 & 6 & 0 \\ 4 & 0 & 2 \end{pmatrix}$

Characteristic eqn: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$S_1 =$ Sum of the main diagonal elements $= 2 + 6 + 2 = 10$

$S_2 =$ Sum of the minors of main diagonal elements

$$= \begin{vmatrix} 6 & 0 \\ 0 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 4 \\ 4 & 2 \end{vmatrix} + \begin{vmatrix} 2 & 0 \\ 0 & 6 \end{vmatrix} = (12-0) + (4-16) + (12-0) = 12 - 12 + 12 = 12$$

$$S_3 = |A| = 2(12-0) - 0(0-0) + 4(0-24) = 24 - 96 = -72$$

Hence the characteristic eqn. is $\lambda^3 - 10\lambda^2 + 12\lambda + 72 = 0$

$$\lambda = -2 \left| \begin{array}{ccc|c} 1 & -10 & 12 & 72 \\ & -2 & 24 & -72 \\ \hline 1 & -12 & 36 & 0 \end{array} \right.$$

$$\lambda^2 - 12\lambda + 36 = 0$$

$$(\lambda - 6)(\lambda - 6) = 0$$

$$\therefore \lambda = -2, 6, 6$$

Hence the eigenvalues are $-2, 6$ & 6 .

Eigenvectors: $(A - \lambda I)x = 0$

$$\begin{pmatrix} 2-\lambda & 0 & 4 \\ 0 & 6-\lambda & 0 \\ 4 & 0 & 2-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\underline{\lambda = -2} \begin{pmatrix} 4 & 0 & 4 \\ 0 & 8 & 0 \\ 4 & 0 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$4x_1 + 0x_2 + 4x_3 = 0$$

$$0x_1 + 8x_2 + 0x_3 = 0$$

$$4x_1 + 0x_2 + 4x_3 = 0$$

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ 0 & 4 & 4 & 0 \\ 8 & 0 & 0 & 8 \end{array}$$

$$\frac{x_1}{0-32} = \frac{x_2}{0-0} = \frac{x_3}{32-0} \Rightarrow \frac{x_1}{-32} = \frac{x_2}{0} = \frac{x_3}{32} \Rightarrow \frac{x_1}{-1} = \frac{x_2}{0} = \frac{x_3}{1}$$

$$\therefore x_1 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$$

$$\underline{\lambda = 6} \begin{pmatrix} -4 & 0 & 4 \\ 0 & 0 & 0 \\ 4 & 0 & -4 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$-4x_1 + 0x_2 + 4x_3 = 0$$

$$0x_1 + 0x_2 + 0x_3 = 0$$

$$4x_1 + 0x_2 - 4x_3 = 0$$

Consider, $4x_1 + 0x_2 - 4x_3 = 0$

Put $x_1 = 0$, $0x_2 - 4x_3 = 0 \Rightarrow 4x_3 = 0x_2 \Rightarrow \frac{x_3}{0} = \frac{x_2}{4} \Rightarrow \frac{x_3}{0} = \frac{x_2}{1}$

$$\therefore x_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$

$$\text{Let } x_3 = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$$

$$x_1^T x_3 = (-1 \ 0 \ 1) \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = -x_1 + 0x_2 + x_3$$

$$\begin{matrix} x_1 & x_2 & x_3 \\ 0 & 1 & -1 & 0 \\ 1 & 0 & 0 & 1 \end{matrix}$$

$$x_2^T x_3 = (0 \ 1 \ 0) \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0x_1 + x_2 + 0x_3$$

$$\frac{x_1}{0-1} = \frac{x_2}{0-0} = \frac{x_3}{-1-0} \Rightarrow \frac{x_1}{-1} = \frac{x_2}{0} = \frac{x_3}{-1} \Rightarrow \frac{x_1}{1} = \frac{x_2}{0} = \frac{x_3}{1}$$

$$\therefore x_3 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$$

$$x_1^T x_2 = (-1 \ 0 \ 1) \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = 0 ; x_1^T x_3 = (-1 \ 0 \ 1) \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} = -1 + 0 + 1 = 0$$

$$x_2^T x_3 = (0 \ 1 \ 0) \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} = 0$$

Hence the eigenvectors are orthogonal to each other.

$$N = \begin{pmatrix} -1/\sqrt{2} & 0 & 1/\sqrt{2} \\ 0 & 1 & 0 \\ 1/\sqrt{2} & 0 & 1/\sqrt{2} \end{pmatrix}, N^T = \begin{pmatrix} -1/\sqrt{2} & 0 & 1/\sqrt{2} \\ 0 & 1 & 0 \\ 1/\sqrt{2} & 0 & 1/\sqrt{2} \end{pmatrix}$$

$$AN = \begin{pmatrix} 2 & 0 & 4 \\ 0 & 6 & 0 \\ 4 & 0 & 2 \end{pmatrix} \begin{pmatrix} -1/\sqrt{2} & 0 & 1/\sqrt{2} \\ 0 & 1 & 0 \\ 1/\sqrt{2} & 0 & 1/\sqrt{2} \end{pmatrix} = \begin{pmatrix} 2/\sqrt{2} & 0 & 6/\sqrt{2} \\ 0 & 6 & 0 \\ -2/\sqrt{2} & 0 & 6/\sqrt{2} \end{pmatrix}$$

$$D = N^T AN = \begin{pmatrix} -1/\sqrt{2} & 0 & 1/\sqrt{2} \\ 0 & 1 & 0 \\ 1/\sqrt{2} & 0 & 1/\sqrt{2} \end{pmatrix} \begin{pmatrix} 2/\sqrt{2} & 0 & 6/\sqrt{2} \\ 0 & 6 & 0 \\ -2/\sqrt{2} & 0 & 6/\sqrt{2} \end{pmatrix} = \begin{pmatrix} -2 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 6 \end{pmatrix}$$

$$\therefore D = \begin{pmatrix} -2 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 6 \end{pmatrix}$$

15) Reduce the quadratic form $3x^2 + 5y^2 + 3z^2 - 2yz + 2zx - 2xy$ to the canonical form through orthogonal transformation. [N/D-2014] [Jan-2011] [M/J-2013]

Sol: Given: Quadratic form $3x^2 + 5y^2 + 3z^2 - 2yz + 2zx - 2xy$

$$A = \begin{pmatrix} \text{coeff. } x^2 & \frac{1}{2} \text{ coeff. } xy & \frac{1}{2} \text{ coeff. } xz \\ \frac{1}{2} \text{ coeff. } xy & \text{coeff. } y^2 & \frac{1}{2} \text{ coeff. } yz \\ \frac{1}{2} \text{ coeff. } xz & \frac{1}{2} \text{ coeff. } yz & \text{coeff. } z^2 \end{pmatrix} = \begin{pmatrix} 3 & -2/2 & 2/2 \\ -2/2 & 5 & -2/2 \\ 2/2 & -2/2 & 3 \end{pmatrix}$$

$$\therefore A = \begin{pmatrix} 3 & -1 & 1 \\ -1 & 5 & -1 \\ 1 & -1 & 3 \end{pmatrix}$$

Characteristic eqn.: $\lambda^3 - S_1 \lambda^2 + S_2 \lambda - S_3 = 0$

$S_1 =$ Sum of the main diagonal elements $= 3 + 5 + 3 = 11$

$S_2 =$ Sum of the minors of main diagonal elements

$$= \begin{vmatrix} 5 & -1 \\ -1 & 3 \end{vmatrix} + \begin{vmatrix} 3 & 1 \\ 1 & 3 \end{vmatrix} + \begin{vmatrix} 3 & -1 \\ -1 & 5 \end{vmatrix} = (15-1) + (9-1) + (15-1) = 14+8+14 = 36$$

$S_3 = |A| = 3(15-1) + 1(-3+1) + 1(1-5) = 3(14) + 1(-2) + 1(-4) = 42 - 2 - 4 = 36$

Hence the characteristic eqn. is $\lambda^3 - 11\lambda^2 + 36\lambda - 36 = 0$

$$\lambda = 2 \quad \left| \begin{array}{ccc|c} 1 & -11 & 36 & -36 \\ & 2 & -18 & 36 \\ \hline 1 & -9 & 18 & 0 \end{array} \right.$$

$$\lambda^2 - 9\lambda + 18 = 0$$

$$(\lambda - 6)(\lambda - 3) = 0$$

$\therefore \lambda = 2, 3, 6$

$$\begin{array}{r|l} x & + \\ 18 & -9 \\ -6 & -3 \\ \lambda - 6 & \lambda - 3 \end{array}$$

Hence the eigenvalues are 2, 3 & 6.

Eigenvectors: $(A - \lambda I)x = 0$

$$\begin{pmatrix} 3-\lambda & -1 & 1 \\ -1 & 5-\lambda & -1 \\ 1 & -1 & 3-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ -1 & 1 & 1 & -1 \\ 3 & -1 & -1 & 3 \end{array}$$

$\lambda = 2$

$$\begin{pmatrix} 1 & -1 & 1 \\ -1 & 3 & -1 \\ 1 & -1 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{aligned} x_1 - x_2 + x_3 &= 0 \\ -x_1 + 3x_2 - x_3 &= 0 \\ x_1 - x_2 + x_3 &= 0 \end{aligned}$$

$$\frac{x_1}{1-3} = \frac{x_2}{-1+1} = \frac{x_3}{3-1} \Rightarrow \frac{x_1}{-2} = \frac{x_2}{0} = \frac{x_3}{2} \Rightarrow \frac{x_1}{-1} = \frac{x_2}{0} = \frac{x_3}{1}$$

$\therefore x_1 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ -1 & 1 & 0 & -1 \\ 2 & -1 & -1 & 2 \end{array}$$

$\lambda = 3$

$$\begin{pmatrix} 0 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{aligned} 0x_1 - x_2 + x_3 &= 0 \\ -x_1 + 2x_2 - x_3 &= 0 \\ x_1 - x_2 + 0x_3 &= 0 \end{aligned}$$

$$\frac{x_1}{1-2} = \frac{x_2}{-1+0} = \frac{x_3}{0-1} \Rightarrow \frac{x_1}{-1} = \frac{x_2}{-1} = \frac{x_3}{-1} \Rightarrow \frac{x_1}{1} = \frac{x_2}{1} = \frac{x_3}{1}$$

$\therefore x_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$

$$\lambda=6 \quad \begin{pmatrix} -3 & -1 & 1 \\ -1 & -1 & -1 \\ 1 & -1 & -3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0 \quad \begin{array}{l} -3x_1 - x_2 + x_3 = 0 \\ -x_1 - x_2 - x_3 = 0 \\ x_1 - x_2 - 3x_3 = 0 \end{array} \quad \begin{array}{cccc} x_1 & x_2 & x_3 & \\ -1 & 1 & -3 & -1 \\ -1 & -1 & -1 & -1 \end{array}$$

$$\frac{x_1}{1+1} = \frac{x_2}{-1-3} = \frac{x_3}{3-1} \Rightarrow \frac{x_1}{2} = \frac{x_2}{-4} = \frac{x_3}{2} \Rightarrow \frac{x_1}{1} = \frac{x_2}{-2} = \frac{x_3}{1}$$

$$\therefore x_3 = \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix}$$

$$x_1^T x_2 = (-1 \ 0 \ 1) \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = -1+0+1=0, \quad x_1^T x_3 = (-1 \ 0 \ 1) \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix} = -1+0+1=0$$

$$x_2^T x_3 = (1 \ 1 \ 1) \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix} = 1-2+1=0$$

Hence the eigenvectors are orthogonal to each other.

$$N = \begin{pmatrix} -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{6}} \\ 0 & \frac{1}{\sqrt{3}} & -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{6}} \end{pmatrix}, \quad N^T = \begin{pmatrix} -\frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{6}} & -\frac{2}{\sqrt{6}} & \frac{1}{\sqrt{6}} \end{pmatrix}$$

$$AN = \begin{pmatrix} 3 & -1 & 1 \\ -1 & 5 & -1 \\ 1 & -1 & 3 \end{pmatrix} \begin{pmatrix} -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{6}} \\ 0 & \frac{1}{\sqrt{3}} & -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{6}} \end{pmatrix} = \begin{pmatrix} -\frac{2}{\sqrt{2}} & \frac{3}{\sqrt{3}} & \frac{6}{\sqrt{6}} \\ 0 & \frac{3}{\sqrt{3}} & -\frac{12}{\sqrt{6}} \\ \frac{2}{\sqrt{2}} & \frac{3}{\sqrt{3}} & \frac{6}{\sqrt{6}} \end{pmatrix}$$

$$D = N^T AN = \begin{pmatrix} -\frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{6}} & -\frac{2}{\sqrt{6}} & \frac{1}{\sqrt{6}} \end{pmatrix} \begin{pmatrix} -\frac{2}{\sqrt{2}} & \frac{3}{\sqrt{3}} & \frac{6}{\sqrt{6}} \\ 0 & \frac{3}{\sqrt{3}} & -\frac{12}{\sqrt{6}} \\ \frac{2}{\sqrt{2}} & \frac{3}{\sqrt{3}} & \frac{6}{\sqrt{6}} \end{pmatrix} = \begin{pmatrix} 4/2 & 0 & 0 \\ 0 & 9/3 & 0 \\ 0 & 0 & 36/6 \end{pmatrix}$$

$$\therefore D = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 6 \end{pmatrix}$$

Canonical form:

$$(y_1 \ y_2 \ y_3) \begin{pmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 6 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = (2y_1 \ 3y_2 \ 6y_3) \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = 2y_1^2 + 3y_2^2 + 6y_3^2$$

(16) Reduce the quadratic form $2x^2 + 5y^2 + 3z^2 + 4xy$ to a canonical form through an orthogonal transformation. Find also its nature. [A/M 2018]

[M/J-2010]

Sol: Given: Quadratic form $2x^2 + 5y^2 + 3z^2 + 4xy$

[Jan-2012]

$$A = \begin{pmatrix} 2 & 2 & 0 \\ 2 & 5 & 0 \\ 0 & 0 & 3 \end{pmatrix}$$

Characteristic eqn.: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$S_1 =$ Sum of the main diagonal elements $= 2+5+3=10$

$S_2 =$ Sum of the minors of main diagonal elements

$$= \begin{vmatrix} 5 & 0 \\ 0 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 0 \\ 0 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 2 \\ 2 & 5 \end{vmatrix} = (15-0) + (6-0) + (10-4) = 15+6+6 = 27$$

$$S_3 = |A| = 2(15-0) - 2(6-0) + 0(0-0) = 30 - 12 = 18$$

Hence the characteristic eqn. is $\lambda^3 - 10\lambda^2 + 27\lambda - 18 = 0$

$$\lambda = 1 \left| \begin{array}{ccc|c} 1 & -10 & 27 & -18 \\ & 1 & -9 & 18 \\ \hline 1 & -9 & 18 & 0 \end{array} \right.$$

$$\lambda^2 - 9\lambda + 18 = 0$$

$$(\lambda - 6)(\lambda - 3) = 0$$

$$\therefore \lambda = 1, 3, 6$$

Hence the eigenvalues are 1, 3 & 6.

Eigenvectors: $(A - \lambda I)x = 0$

$$\begin{pmatrix} 2-\lambda & 2 & 0 \\ 2 & 5-\lambda & 0 \\ 0 & 0 & 3-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{aligned} x_1 + 2x_2 + 0x_3 &= 0 \\ 2x_1 + 4x_2 + 0x_3 &= 0 \\ 0x_1 + 0x_2 + 2x_3 &= 0 \end{aligned}$$

① & ③

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ 2 & 0 & 1 & 2 \\ 0 & 2 & 0 & 0 \end{array}$$

$$\underline{\lambda=1} \begin{pmatrix} 1 & 2 & 0 \\ 2 & 4 & 0 \\ 0 & 0 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\frac{x_1}{4-0} = \frac{x_2}{0-2} = \frac{x_3}{0-0} \Rightarrow \frac{x_1}{4} = \frac{x_2}{-2} = \frac{x_3}{0} \Rightarrow \frac{x_1}{2} = \frac{x_2}{-1} = \frac{x_3}{0}$$

$$\therefore x_1 = \begin{pmatrix} 2 \\ -1 \\ 0 \end{pmatrix}$$

$$\underline{\lambda=3} \begin{pmatrix} -1 & 2 & 0 \\ 2 & 2 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{aligned} -x_1 + 2x_2 + 0x_3 &= 0 \\ 2x_1 + 2x_2 + 0x_3 &= 0 \\ 0x_1 + 0x_2 + 0x_3 &= 0 \end{aligned}$$

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ 2 & 0 & -1 & 2 \\ 2 & 0 & 2 & 2 \end{array}$$

$$\frac{x_1}{0-0} = \frac{x_2}{0-0} = \frac{x_3}{-2-4} \Rightarrow \frac{x_1}{0} = \frac{x_2}{0} = \frac{x_3}{-6} \Rightarrow \frac{x_1}{0} = \frac{x_2}{0} = \frac{x_3}{-1}$$

$$\therefore x_2 = \begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix}$$

$$\underline{\lambda=6} \begin{pmatrix} -4 & 2 & 0 \\ 2 & -1 & 0 \\ 0 & 0 & -3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{aligned} -4x_1 + 2x_2 + 0x_3 &= 0 \\ 2x_1 - x_2 + 0x_3 &= 0 \\ 0x_1 + 0x_2 - 3x_3 &= 0 \end{aligned}$$

② & ③

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ -1 & 0 & 2 & -1 \\ 0 & -3 & 0 & 0 \end{array}$$

$$\frac{x_1}{3-0} = \frac{x_2}{0+6} = \frac{x_3}{0-0} \Rightarrow \frac{x_1}{3} = \frac{x_2}{6} = \frac{x_3}{0} \Rightarrow \frac{x_1}{1} = \frac{x_2}{2} = \frac{x_3}{0}$$

$$\therefore x_3 = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}$$

$$x_1^T x_2 = (2 \ -1 \ 0) \begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix} = 0, \quad x_1^T x_3 = (2 \ -1 \ 0) \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} = 2 - 2 + 0 = 0$$

$$x_2^T x_3 = (0 \ 0 \ -1) \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} = 0$$

Hence the eigenvectors are orthogonal to each other.

$$N = \begin{pmatrix} \frac{2}{\sqrt{5}} & 0 & \frac{1}{\sqrt{5}} \\ -\frac{1}{\sqrt{5}} & 0 & \frac{2}{\sqrt{5}} \\ 0 & -1 & 0 \end{pmatrix}, \quad N^T = \begin{pmatrix} \frac{2}{\sqrt{5}} & -\frac{1}{\sqrt{5}} & 0 \\ 0 & 0 & -1 \\ \frac{1}{\sqrt{5}} & \frac{2}{\sqrt{5}} & 0 \end{pmatrix}$$

$$AN = \begin{pmatrix} 2 & 2 & 0 \\ 2 & 5 & 0 \\ 0 & 0 & 3 \end{pmatrix} \begin{pmatrix} \frac{2}{\sqrt{5}} & 0 & \frac{1}{\sqrt{5}} \\ -\frac{1}{\sqrt{5}} & 0 & \frac{2}{\sqrt{5}} \\ 0 & -1 & 0 \end{pmatrix} = \begin{pmatrix} \frac{2}{\sqrt{5}} & 0 & \frac{6}{\sqrt{5}} \\ -\frac{1}{\sqrt{5}} & 0 & \frac{12}{\sqrt{5}} \\ 0 & -3 & 0 \end{pmatrix}$$

$$D = N^T AN = \begin{pmatrix} \frac{2}{\sqrt{5}} & -\frac{1}{\sqrt{5}} & 0 \\ 0 & 0 & -1 \\ \frac{1}{\sqrt{5}} & \frac{2}{\sqrt{5}} & 0 \end{pmatrix} \begin{pmatrix} \frac{2}{\sqrt{5}} & 0 & \frac{6}{\sqrt{5}} \\ -\frac{1}{\sqrt{5}} & 0 & \frac{12}{\sqrt{5}} \\ 0 & -3 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 6 \end{pmatrix}$$

∴ Canonical form:

$$(y_1 \ y_2 \ y_3) \begin{pmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 6 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = (y_1 \ 3y_2 \ 6y_3) \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = y_1^2 + 3y_2^2 + 6y_3^2$$

Canonical form contains only +ve terms. ∴ Quadratic form is said to be positive definite.

①7 Reduce the quadratic form $6x_1^2 + 3x_2^2 + 3x_3^2 - 4x_1x_2 - 2x_2x_3 + 4x_3x_1$ to canonical form. Hence find its rank, signature, index & nature.

Sol: Given: Q.F $6x_1^2 + 3x_2^2 + 3x_3^2 - 4x_1x_2 - 2x_2x_3 + 4x_3x_1$

$$A = \begin{pmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{pmatrix}$$

Characteristic eqn.: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$S_1 =$ Sum of the main diagonal elements $= 6 + 3 + 3 = 12$

$S_2 =$ Sum of the minors of main diagonal elements

$$= \begin{vmatrix} 3 & -1 \\ -1 & 3 \end{vmatrix} + \begin{vmatrix} 6 & 2 \\ 2 & 3 \end{vmatrix} + \begin{vmatrix} 6 & -2 \\ -2 & 3 \end{vmatrix} = (9 - 1) + (18 - 4) + (18 - 4) = 8 + 14 + 14 = 36$$

$$S_3 = |A| = 6(9 - 1) + 2(-6 + 2) + 2(2 - 6) = 6(8) + 2(-4) + 2(-4) = 48 - 8 - 8 = 32$$

Hence the characteristic eqn. is $\lambda^3 - 12\lambda^2 + 36\lambda - 32 = 0$

$$\lambda = 2 \left| \begin{array}{ccc|c} -12 & 36 & -32 & \\ \hline 2 & -20 & 32 & \\ \hline 1 & -10 & 16 & 0 \end{array} \right.$$

$$\lambda^2 - 10\lambda + 16 = 0$$

$$(\lambda - 8)(\lambda - 2) = 0$$

$$\therefore \lambda = 2, 2, 8$$

$$\begin{array}{c|c} \lambda & + \\ \hline 16 & -10 \\ \hline -8 & -2 \\ \hline \lambda - 8 & \lambda - 2 \end{array}$$

Hence the eigenvalues are 2, 2 & 8.

Eigenvectors: $(A - \lambda I)x = 0$

$$\begin{pmatrix} 6 - \lambda & -2 & 2 \\ -2 & 3 - \lambda & -1 \\ 2 & -1 & 3 - \lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\underline{\lambda = 2} \begin{pmatrix} 4 & -2 & 2 \\ -2 & 1 & -1 \\ 2 & -1 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{aligned} 4x_1 - 2x_2 + 2x_3 &= 0 \\ -2x_1 + x_2 - x_3 &= 0 \\ 2x_1 - x_2 + x_3 &= 0 \end{aligned}$$

Consider, $2x_1 - x_2 + x_3 = 0$

Put $x_1 = 0 \Rightarrow -x_2 + x_3 = 0 \Rightarrow x_2 = x_3$

$$\therefore x_1 = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$$

$$\underline{\lambda = 8} \begin{pmatrix} -2 & -2 & 2 \\ -2 & -5 & -1 \\ 2 & -1 & -5 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{aligned} -2x_1 - 2x_2 + 2x_3 &= 0 \\ -2x_1 - 5x_2 - x_3 &= 0 \\ 2x_1 - x_2 - 5x_3 &= 0 \end{aligned}$$

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ -2 & 2 & -2 & -2 \\ -5 & -1 & -2 & -5 \end{array}$$

$$\frac{x_1}{2+10} = \frac{x_2}{-4-2} = \frac{x_3}{10-4} \Rightarrow \frac{x_1}{12} = \frac{x_2}{-6} = \frac{x_3}{6} \Rightarrow \frac{x_1}{2} = \frac{x_2}{-1} = \frac{x_3}{1}$$

$$\therefore x_2 = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}$$

Let $x_3 = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$

$$x_1^T x_3 = (0 \ 1 \ 1) \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0x_1 + x_2 + x_3$$

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ 1 & 1 & 0 & 1 \\ -1 & 1 & 2 & -1 \end{array}$$

$$x_2^T x_3 = (2 \ -1 \ 1) \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 2x_1 - x_2 + x_3$$

$$\frac{x_1}{1+1} = \frac{x_2}{2-0} = \frac{x_3}{0-2} \Rightarrow \frac{x_1}{2} = \frac{x_2}{2} = \frac{x_3}{-2} \Rightarrow \frac{x_1}{1} = \frac{x_2}{1} = \frac{x_3}{-1}$$

$$\therefore x_3 = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$$

$$x_1^T x_2 = (0 \ 1 \ 1) \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} = 0 - 1 + 1 = 0, \quad x_1^T x_3 = (0 \ 1 \ 1) \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} = 0 + 1 - 1 = 0$$

$$x_2^T x_3 = (2 \ -1 \ 1) \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} = 2 - 1 - 1 = 0$$

Hence the eigenvectors are orthogonal to each other.

$$N = \begin{pmatrix} 0 & 2/\sqrt{6} & 1/\sqrt{3} \\ 1/\sqrt{2} & -1/\sqrt{6} & 1/\sqrt{3} \\ 1/\sqrt{2} & 1/\sqrt{6} & -1/\sqrt{3} \end{pmatrix}$$

$$N^T = \begin{pmatrix} 0 & 1/\sqrt{2} & 1/\sqrt{2} \\ 2/\sqrt{6} & -1/\sqrt{6} & 1/\sqrt{6} \\ 1/\sqrt{3} & 1/\sqrt{3} & -1/\sqrt{3} \end{pmatrix}$$

$$AN = \begin{pmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{pmatrix} \begin{pmatrix} 0 & 2/\sqrt{6} & 1/\sqrt{3} \\ 1/\sqrt{2} & -1/\sqrt{6} & 1/\sqrt{3} \\ 1/\sqrt{2} & 1/\sqrt{6} & -1/\sqrt{3} \end{pmatrix} = \begin{pmatrix} 0 & 16/\sqrt{6} & 2/\sqrt{3} \\ 2/\sqrt{2} & -8/\sqrt{6} & 2/\sqrt{3} \\ 2/\sqrt{2} & 8/\sqrt{6} & -2/\sqrt{3} \end{pmatrix}$$

$$D = N^T AN = \begin{pmatrix} 0 & 1/\sqrt{2} & 1/\sqrt{2} \\ 2/\sqrt{6} & -1/\sqrt{6} & 1/\sqrt{6} \\ 1/\sqrt{3} & 1/\sqrt{3} & -1/\sqrt{3} \end{pmatrix} \begin{pmatrix} 0 & 16/\sqrt{6} & 2/\sqrt{3} \\ 2/\sqrt{2} & -8/\sqrt{6} & 2/\sqrt{3} \\ 2/\sqrt{2} & 8/\sqrt{6} & -2/\sqrt{3} \end{pmatrix} = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 48/6 & 0 \\ 0 & 0 & 6/3 \end{pmatrix}$$

$$D = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 8 & 0 \\ 0 & 0 & 2 \end{pmatrix}$$

Canonical form:

$$(y_1, y_2, y_3) \begin{pmatrix} 2 & 0 & 0 \\ 0 & 8 & 0 \\ 0 & 0 & 2 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = 2y_1^2 + 8y_2^2 + 2y_3^2$$

Canonical form contains only +ve terms. \therefore Quadratic form is said to be positive definite.

Rank = No. of non-zero terms in C.F = 3

Index = No. of +ve terms in C.F = 3

Signature = (No. of +ve terms - No. of -ve terms) in C.F = 3 - 0 = 3

⑮ Reduce the quadratic form $x_1^2 + 5x_2^2 + x_3^2 + 2x_1x_2 + 2x_2x_3 + 6x_3x_1$ to the canonical form. Hence find its nature & rank. [M/J-2014] [A/M-2015]

Given: Q.F: $x_1^2 + 5x_2^2 + x_3^2 + 2x_1x_2 + 2x_2x_3 + 6x_3x_1$ [N/D-2014]

$$A = \begin{pmatrix} 1 & 1 & 3 \\ 1 & 5 & 1 \\ 3 & 1 & 1 \end{pmatrix}$$

Characteristic eqn/: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

S_1 = Sum of the main diagonal elements = 1 + 5 + 1 = 7

S_2 = Sum of the minors of main diagonal elements

$$= \begin{vmatrix} 5 & 1 \\ 1 & 1 \end{vmatrix} + \begin{vmatrix} 1 & 3 \\ 3 & 1 \end{vmatrix} + \begin{vmatrix} 1 & 1 \\ 1 & 5 \end{vmatrix} = (5-1) + (1-9) + (5-1) = 4 - 8 + 4 = 0$$

$$S_3 = |A| = 1(5-1) - 1(1-3) + 3(1-15) = 4 + 2 - 42 = -36$$

Hence the characteristic eqn/ is $\lambda^3 - 7\lambda^2 + 0\lambda + 36 = 0 \Rightarrow \lambda^3 - 7\lambda^2 + 36 = 0$

$$\lambda=3 \left[\begin{array}{ccc|c} 1 & -7 & 0 & 36 \\ & 3 & -12 & -36 \\ \hline 1 & -4 & -12 & 0 \end{array} \right] \begin{array}{l} x \\ -12 \\ -6 \\ \lambda-6 \end{array} \begin{array}{l} + \\ -4 \\ +2 \\ \lambda+2 \end{array}$$

$$\lambda^2 - 4\lambda - 12 = 0$$

$$(\lambda - 6)(\lambda + 2) = 0$$

$$\therefore \lambda = -2, 3, 6$$

Hence the eigenvalues are -2, 3 & 6.

Eigenvectors: $(A - \lambda I)x = 0$

$$\begin{pmatrix} 1-\lambda & 1 & 3 \\ 1 & 5-\lambda & 1 \\ 3 & 1 & 1-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\underline{\lambda = -2} \begin{pmatrix} 3 & 1 & 3 \\ 1 & 7 & 1 \\ 3 & 1 & 3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{aligned} 3x_1 + x_2 + 3x_3 &= 0 \\ x_1 + 7x_2 + x_3 &= 0 \\ 3x_1 + x_2 + 3x_3 &= 0 \end{aligned}$$

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ 1 & 3 & 3 & 1 \\ 7 & 1 & 1 & 7 \end{array}$$

$$\frac{x_1}{1-21} = \frac{x_2}{3-3} = \frac{x_3}{21-1} \Rightarrow \frac{x_1}{-20} = \frac{x_2}{0} = \frac{x_3}{20} \Rightarrow \frac{x_1}{-1} = \frac{x_2}{0} = \frac{x_3}{1}$$

$$\therefore x_1 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$$

$$\underline{\lambda = 3} \begin{pmatrix} -2 & 1 & 3 \\ 1 & 2 & 1 \\ 3 & 1 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{aligned} -2x_1 + x_2 + 3x_3 &= 0 \\ x_1 + 2x_2 + x_3 &= 0 \\ 3x_1 + x_2 - 2x_3 &= 0 \end{aligned}$$

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ 1 & 3 & -2 & 1 \\ 2 & 1 & 1 & 2 \end{array}$$

$$\frac{x_1}{1-6} = \frac{x_2}{3+2} = \frac{x_3}{-4-1} \Rightarrow \frac{x_1}{-5} = \frac{x_2}{5} = \frac{x_3}{-5} \Rightarrow \frac{x_1}{-1} = \frac{x_2}{1} = \frac{x_3}{-1}$$

$$\therefore x_2 = \begin{pmatrix} -1 \\ 1 \\ -1 \end{pmatrix}$$

$$\underline{\lambda = 6} \begin{pmatrix} -5 & 1 & 3 \\ 1 & -1 & 1 \\ 3 & 1 & -5 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\begin{aligned} -5x_1 + x_2 + 3x_3 &= 0 \\ x_1 - x_2 + x_3 &= 0 \\ 3x_1 + x_2 - 5x_3 &= 0 \end{aligned}$$

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ 1 & 3 & -5 & 1 \\ -1 & 1 & 1 & -1 \end{array}$$

$$\frac{x_1}{1+3} = \frac{x_2}{3+5} = \frac{x_3}{5-1} \Rightarrow \frac{x_1}{4} = \frac{x_2}{8} = \frac{x_3}{4} \Rightarrow \frac{x_1}{1} = \frac{x_2}{2} = \frac{x_3}{1}$$

$$\therefore x_3 = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$$

$$x_1^T x_2 = (-1 \ 0 \ 1) \begin{pmatrix} -1 \\ 1 \\ -1 \end{pmatrix} = 1 + 0 - 1 = 0, \quad x_1^T x_3 = (-1 \ 0 \ 1) \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} = -1 + 0 + 1 = 0$$

$$x_2^T x_3 = (-1 \ 1 \ -1) \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} = -1 + 2 - 1 = 0$$

Hence the eigenvectors are orthogonal to each other.

$$N = \begin{pmatrix} -1/\sqrt{2} & -1/\sqrt{3} & 1/\sqrt{6} \\ 0 & 1/\sqrt{3} & 2/\sqrt{6} \\ 1/\sqrt{2} & -1/\sqrt{3} & 1/\sqrt{6} \end{pmatrix}$$

$$N^T = \begin{pmatrix} -1/\sqrt{2} & 0 & 1/\sqrt{2} \\ -1/\sqrt{3} & 1/\sqrt{3} & -1/\sqrt{3} \\ 1/\sqrt{6} & 2/\sqrt{6} & 1/\sqrt{6} \end{pmatrix}$$

$$AN = \begin{pmatrix} 1 & 1 & 3 \\ 1 & 5 & 1 \\ 3 & 1 & 1 \end{pmatrix} \begin{pmatrix} -1/\sqrt{2} & -1/\sqrt{3} & 1/\sqrt{6} \\ 0 & 1/\sqrt{3} & 2/\sqrt{6} \\ 1/\sqrt{2} & -1/\sqrt{3} & 1/\sqrt{6} \end{pmatrix} = \begin{pmatrix} 2/\sqrt{2} & -3/\sqrt{3} & 6/\sqrt{6} \\ 0 & 3/\sqrt{3} & 12/\sqrt{6} \\ -2/\sqrt{2} & -3/\sqrt{3} & 6/\sqrt{6} \end{pmatrix}$$

$$D = N^T AN = \begin{pmatrix} -1/\sqrt{2} & 0 & 1/\sqrt{2} \\ -1/\sqrt{3} & 1/\sqrt{3} & -1/\sqrt{3} \\ 1/\sqrt{6} & 2/\sqrt{6} & 1/\sqrt{6} \end{pmatrix} \begin{pmatrix} 2/\sqrt{2} & -3/\sqrt{3} & 6/\sqrt{6} \\ 0 & 3/\sqrt{3} & 12/\sqrt{6} \\ -2/\sqrt{2} & -3/\sqrt{3} & 6/\sqrt{6} \end{pmatrix} = \begin{pmatrix} -2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 6 \end{pmatrix}$$

Canonical form:

$$(y_1 \ y_2 \ y_3) \begin{pmatrix} -2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 6 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = -2y_1^2 + 3y_2^2 + 6y_3^2$$

Canonical form contains 2 +ve terms & one -ve term. \therefore Quadratic form is said to be indefinite.

Rank = No. of non-zero terms in C.F. = 3

(19) Reduce the quadratic form $2x^2 + y^2 + z^2 + 2xy - 2xz - 4yz$ to the canonical form. Hence find its nature, rank, index & signature. [A/M-2015] [N/D-2010]

Sol: Q.F: $2x^2 + y^2 + z^2 + 2xy - 2xz - 4yz$

$$A = \begin{pmatrix} 2 & 1 & -1 \\ 1 & 1 & -2 \\ -1 & -2 & 1 \end{pmatrix}$$

Characteristic eqn.: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$S_1 =$ Sum of the main diagonal elements = $2+1+1=4$

$S_2 =$ Sum of the minors of main diagonal elements

$$= \begin{vmatrix} 1 & -2 \\ -2 & 1 \end{vmatrix} + \begin{vmatrix} 2 & -1 \\ 1 & 1 \end{vmatrix} + \begin{vmatrix} 2 & 1 \\ 1 & 1 \end{vmatrix} = (1-4) + (2-1) + (2-1) = -3+1+1 = -1$$

$S_3 = |A| = 2(1-4) - 1(1-2) - 1(-2+1) = -6+1+1 = -4$

Hence the characteristic eqn. is $\lambda^3 - 4\lambda^2 - \lambda + 4 = 0$

$$\lambda = 1 \left| \begin{array}{ccc|c} 1 & -4 & -1 & 4 \\ & 1 & -3 & -4 \\ 1 & -3 & -4 & 0 \end{array} \right.$$

$$\begin{array}{l} \lambda^2 - 3\lambda - 4 = 0 \\ (\lambda+1)(\lambda-4) = 0 \\ \therefore \lambda = -1, 1, 4 \end{array} \quad \begin{array}{l} x \quad + \\ -4 \quad -3 \\ +1 \quad -4 \\ \lambda+1 \quad \lambda-4 \end{array}$$

Hence the eigenvalues are -1, 1 & 4.

Eigenvectors: $(A - \lambda I)x = 0$

$$\begin{pmatrix} 2-\lambda & 1 & -1 \\ 1 & 1-\lambda & -2 \\ -1 & -2 & 1-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\underline{\lambda = -1} \begin{pmatrix} 3 & 1 & -1 \\ 1 & 2 & -2 \\ -1 & -2 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0 \quad \begin{matrix} 3x_1 + x_2 - x_3 = 0 \\ x_1 + 2x_2 - 2x_3 = 0 \\ -x_1 - 2x_2 + 2x_3 = 0 \end{matrix} \quad \begin{matrix} x_1 & x_2 & x_3 & | \\ 1 & -1 & 3 & | \\ 2 & -2 & 1 & | \end{matrix}$$

$$\frac{x_1}{-2+2} = \frac{x_2}{-1+6} = \frac{x_3}{6-1} \Rightarrow \frac{x_1}{0} = \frac{x_2}{5} = \frac{x_3}{5} \Rightarrow \frac{x_1}{0} = \frac{x_2}{1} = \frac{x_3}{1}$$

$$\therefore x_1 = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$$

$$\underline{\lambda = 1} \begin{pmatrix} 1 & 1 & -1 \\ 1 & 0 & -2 \\ -1 & -2 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0 \quad \begin{matrix} x_1 + x_2 - x_3 = 0 \\ x_1 + 0x_2 - 2x_3 = 0 \\ -x_1 - 2x_2 + 0x_3 = 0 \end{matrix} \quad \begin{matrix} x_1 & x_2 & x_3 & | \\ 1 & -1 & 1 & | \\ 0 & -2 & 1 & | \end{matrix}$$

$$\frac{x_1}{-2+0} = \frac{x_2}{-1+2} = \frac{x_3}{0-1} \Rightarrow \frac{x_1}{-2} = \frac{x_2}{1} = \frac{x_3}{-1}$$

$$\therefore x_2 = \begin{pmatrix} -2 \\ 1 \\ -1 \end{pmatrix}$$

$$\underline{\lambda = 4} \begin{pmatrix} -2 & 1 & -1 \\ 1 & -3 & -2 \\ -1 & -2 & -3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0 \quad \begin{matrix} -2x_1 + x_2 - x_3 = 0 \\ x_1 - 3x_2 - 2x_3 = 0 \\ -x_1 - 2x_2 - 3x_3 = 0 \end{matrix} \quad \begin{matrix} x_1 & x_2 & x_3 & | \\ 1 & -1 & -2 & | \\ -3 & -2 & 1 & | \end{matrix}$$

$$\frac{x_1}{-2-3} = \frac{x_2}{-1-4} = \frac{x_3}{6-1} \Rightarrow \frac{x_1}{-5} = \frac{x_2}{-5} = \frac{x_3}{5} \Rightarrow \frac{x_1}{-1} = \frac{x_2}{-1} = \frac{x_3}{1}$$

$$\therefore x_3 = \begin{pmatrix} -1 \\ -1 \\ 1 \end{pmatrix}$$

$$x_1^T x_2 = (0 \ 1 \ 1) \begin{pmatrix} -2 \\ 1 \\ -1 \end{pmatrix} = 0 + 1 - 1 = 0, \quad x_1^T x_3 = (0 \ 1 \ 1) \begin{pmatrix} -1 \\ -1 \\ 1 \end{pmatrix} = 0 - 1 + 1 = 0$$

$$x_2^T x_3 = (-2 \ 1 \ -1) \begin{pmatrix} -1 \\ -1 \\ 1 \end{pmatrix} = 2 - 1 - 1 = 0$$

Hence the eigenvectors are orthogonal to each other.

$$N = \begin{pmatrix} 0 & -2/\sqrt{6} & -1/\sqrt{3} \\ 1/\sqrt{2} & 1/\sqrt{6} & -1/\sqrt{3} \\ 1/\sqrt{2} & -1/\sqrt{6} & 1/\sqrt{3} \end{pmatrix} \quad N^T = \begin{pmatrix} 0 & 1/\sqrt{2} & 1/\sqrt{2} \\ -2/\sqrt{6} & 1/\sqrt{6} & -1/\sqrt{6} \\ -1/\sqrt{3} & -1/\sqrt{3} & 1/\sqrt{3} \end{pmatrix}$$

$$AN = \begin{pmatrix} 2 & 1 & -1 \\ 1 & 1 & -2 \\ -1 & -2 & 1 \end{pmatrix} \begin{pmatrix} 0 & -2/\sqrt{6} & -1/\sqrt{3} \\ 1/\sqrt{2} & 1/\sqrt{6} & -1/\sqrt{3} \\ 1/\sqrt{2} & -1/\sqrt{6} & 1/\sqrt{3} \end{pmatrix} = \begin{pmatrix} 0 & -2/\sqrt{6} & -4/\sqrt{3} \\ -1/\sqrt{2} & 1/\sqrt{6} & -4/\sqrt{3} \\ -1/\sqrt{2} & -1/\sqrt{6} & 4/\sqrt{3} \end{pmatrix}$$

$$D = N^T AN = \begin{pmatrix} 0 & 1/\sqrt{2} & 1/\sqrt{2} \\ -2/\sqrt{6} & 1/\sqrt{6} & -1/\sqrt{6} \\ -1/\sqrt{3} & -1/\sqrt{3} & 1/\sqrt{3} \end{pmatrix} \begin{pmatrix} 0 & -2/\sqrt{6} & -4/\sqrt{3} \\ -1/\sqrt{2} & 1/\sqrt{6} & -4/\sqrt{3} \\ -1/\sqrt{2} & -1/\sqrt{6} & 4/\sqrt{3} \end{pmatrix} = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 4 \end{pmatrix}$$

Canonical form: $(y_1 \ y_2 \ y_3) \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 4 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = -y_1^2 + y_2^2 + 4y_3^2$

Canonical form contains 2 +ve terms & one -ve term. \therefore Quadratic form is said to be indefinite.

Rank = No. of non-zero terms in C.F = 3

Index = No. of +ve terms in C.F = 2

Signature = (No. of +ve terms - No. of -ve terms) in C.F = 2 - 1 = 1

② Reduce the quadratic form $x_1^2 + 2x_2^2 + x_3^2 - 2x_1x_2 + 2x_2x_3$ to the canonical form through an orthogonal transformation, & hence show that it is +ve semi-definite. Also given a non-zero set of values (x_1, x_2, x_3) which makes this quadratic form zero. [M/J-2009]

Sol: Given: Q.F $x_1^2 + 2x_2^2 + x_3^2 - 2x_1x_2 + 2x_2x_3$

$$A = \begin{pmatrix} 1 & -1 & 0 \\ -1 & 2 & 1 \\ 0 & 1 & 1 \end{pmatrix}$$

Characteristic eqn: $\lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$

$S_1 = 1 + 2 + 1 = 4$

$S_2 = \begin{vmatrix} 2 & 1 \\ 1 & 1 \end{vmatrix} + \begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} + \begin{vmatrix} 1 & -1 \\ -1 & 2 \end{vmatrix} = (2-1) + (1-0) + (2-1) = 1+1+1 = 3$

$S_3 = |A| = 1(2-1) + 1(-1-0) + 0(-1-0) = 1-1 = 0$

Hence the characteristic eqn. is $\lambda^3 - 4\lambda^2 + 3\lambda = 0$

$\lambda(\lambda^2 - 4\lambda + 3) = 0$
 $\lambda = 0, (\lambda-1)(\lambda-3) = 0$

$\therefore \lambda = 0, 1, 3$

x	+
3	-4
-1	-3
$\lambda-1$	$\lambda-3$

Hence the eigenvalues are 0, 1 & 3:

Eigenvectors: $(A - \lambda I)x = 0$

$$\begin{pmatrix} 1-\lambda & -1 & 0 \\ -1 & 2-\lambda & 1 \\ 0 & 1 & 1-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$\lambda = 0$ $\begin{pmatrix} 1 & -1 & 0 \\ -1 & 2 & 1 \\ 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$ $\begin{matrix} x_1 - x_2 + 0x_3 = 0 \\ -x_1 + 2x_2 + x_3 = 0 \\ 0x_1 + x_2 + x_3 = 0 \end{matrix}$

x_1	x_2	x_3
-1	0	1
2	1	-1

$\frac{x_1}{-1-0} = \frac{x_2}{0-1} = \frac{x_3}{2-1} \Rightarrow \frac{x_1}{-1} = \frac{x_2}{-1} = \frac{x_3}{1}$

$\therefore x_1 = \begin{pmatrix} -1 \\ -1 \\ 1 \end{pmatrix}$

$$\underline{\lambda=1} \begin{pmatrix} 0 & -1 & 0 \\ -1 & 1 & 1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0 \quad \begin{matrix} 0x_1 - x_2 + 0x_3 = 0 \\ -x_1 + x_2 + x_3 = 0 \\ 0x_1 + x_2 + 0x_3 = 0 \end{matrix} \quad \begin{matrix} x_1 & x_2 & x_3 & - \\ -1 & 0 & 0 & -1 \\ 1 & 1 & -1 & 1 \end{matrix}$$

$$\frac{x_1}{-1-0} = \frac{x_2}{0-0} = \frac{x_3}{0-1} \Rightarrow \frac{x_1}{-1} = \frac{x_2}{0} = \frac{x_3}{-1} \Rightarrow \frac{x_1}{1} = \frac{x_2}{0} = \frac{x_3}{1}$$

$$\therefore x_2 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$$

$$\underline{\lambda=3} \begin{pmatrix} -2 & -1 & 0 \\ -1 & -1 & 1 \\ 0 & 1 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0 \quad \begin{matrix} -2x_1 - x_2 + 0x_3 = 0 \\ -x_1 - x_2 + x_3 = 0 \\ 0x_1 + x_2 - 2x_3 = 0 \end{matrix} \quad \begin{matrix} x_1 & x_2 & x_3 & - \\ -1 & 0 & -2 & -1 \\ -1 & 1 & -1 & -1 \end{matrix}$$

$$\frac{x_1}{-1+0} = \frac{x_2}{0+2} = \frac{x_3}{2-1} \Rightarrow \frac{x_1}{-1} = \frac{x_2}{2} = \frac{x_3}{1} \quad \therefore x_3 = \begin{pmatrix} -1 \\ 2 \\ 1 \end{pmatrix}$$

$$x_1^T x_2 = (-1 \ -1 \ 1) \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} = -1+0+1=0, \quad x_1^T x_3 = (-1 \ -1 \ 1) \begin{pmatrix} -1 \\ 2 \\ 1 \end{pmatrix} = 1-2+1=0$$

$$x_2^T x_3 = (1 \ 0 \ 1) \begin{pmatrix} -1 \\ 2 \\ 1 \end{pmatrix} = -1+0+1=0$$

Hence the eigenvectors are orthogonal to each other.

$$N = \begin{pmatrix} -1/\sqrt{3} & 1/\sqrt{2} & -1/\sqrt{6} \\ -1/\sqrt{3} & 0 & 2/\sqrt{6} \\ 1/\sqrt{3} & 1/\sqrt{2} & 1/\sqrt{6} \end{pmatrix} \quad N^T = \begin{pmatrix} -1/\sqrt{3} & -1/\sqrt{3} & 1/\sqrt{3} \\ 1/\sqrt{2} & 0 & 1/\sqrt{2} \\ -1/\sqrt{6} & 2/\sqrt{6} & 1/\sqrt{6} \end{pmatrix}$$

$$AN = \begin{pmatrix} 1 & -1 & 0 \\ -1 & 2 & 1 \\ 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} -1/\sqrt{3} & 1/\sqrt{2} & -1/\sqrt{6} \\ -1/\sqrt{3} & 0 & 2/\sqrt{6} \\ 1/\sqrt{3} & 1/\sqrt{2} & 1/\sqrt{6} \end{pmatrix} = \begin{pmatrix} 0 & 1/\sqrt{2} & -3/\sqrt{6} \\ 0 & 0 & 6/\sqrt{6} \\ 0 & 1/\sqrt{2} & 3/\sqrt{6} \end{pmatrix}$$

$$D = N^T AN = \begin{pmatrix} -1/\sqrt{3} & -1/\sqrt{3} & 1/\sqrt{3} \\ 1/\sqrt{2} & 0 & 1/\sqrt{2} \\ -1/\sqrt{6} & 2/\sqrt{6} & 1/\sqrt{6} \end{pmatrix} \begin{pmatrix} 0 & 1/\sqrt{2} & -3/\sqrt{6} \\ 0 & 0 & 6/\sqrt{6} \\ 0 & 1/\sqrt{2} & 3/\sqrt{6} \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 3 \end{pmatrix}$$

Canonical form:

$$(y_1 \ y_2 \ y_3) \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 3 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = 0y_1^2 + y_2^2 + 3y_3^2$$

Canonical form contains 2 +ve terms & one zero.

\therefore Quadratic form is said to be positive semi-definite.

The orthogonal transformation is $x = NY$

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -1/\sqrt{3} & 1/\sqrt{2} & -1/\sqrt{6} \\ -1/\sqrt{3} & 0 & 2/\sqrt{6} \\ 1/\sqrt{3} & 1/\sqrt{2} & 1/\sqrt{6} \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$$

$$x_1 = -\frac{1}{\sqrt{3}}y_1 + \frac{1}{\sqrt{2}}y_2 - \frac{1}{\sqrt{6}}y_3$$

$$x_2 = -\frac{1}{\sqrt{3}}y_1 + 0y_2 + \frac{2}{\sqrt{6}}y_3$$

$$x_3 = \frac{1}{\sqrt{3}}y_1 + \frac{1}{\sqrt{2}}y_2 + \frac{1}{\sqrt{6}}y_3$$

Take $y_1 = \sqrt{3}$, $y_2 = 0$ & $y_3 = 0$

$$x_1 = -1, x_2 = -1, x_3 = 1$$

These values x_1, x_2, x_3 make the Q.F. zero.

Verification: $x_1 = -1, x_2 = -1, x_3 = 1$

$$\begin{aligned} \text{Q.F.} &= x_1^2 + 2x_2^2 + x_3^2 - 2x_1x_2 + 2x_2x_3 \\ &= 1 + 2 + 1 - 2 - 2 = 0 \end{aligned}$$

Properties:

① Prove that the eigenvalues of a real symmetric matrix are real. [M/J-2014]

Proof: Let λ be an eigenvalue of the real symmetric matrix A . Let the corresponding eigenvector be x . Let A^T denote the transpose of A .

$$\text{We have } Ax = \lambda x$$

Pre-multiplying this eqn. by $1 \times n$ matrix \bar{x}^T , where the bar denotes the complex conjugate of x^T , we get

$$\bar{x}^T Ax = \lambda \bar{x}^T x \quad \text{--- (1)}$$

Taking complex conjugate, we get

$$x^T \bar{A} \bar{x} = \bar{\lambda} x^T \bar{x}$$

$$x^T A \bar{x} = \bar{\lambda} x^T \bar{x} \quad (\because A \text{ is real})$$

Taking transpose on both sides, we get

$$(x^T A \bar{x})^T = (\bar{\lambda} x^T \bar{x})^T$$

$$\bar{x}^T A^T x = \bar{\lambda} \bar{x}^T x$$

$$\bar{x}^T Ax = \bar{\lambda} \bar{x}^T x \quad (\because A \text{ is symmetric})$$

From (1) & (2), $\lambda \bar{x}^T x = \bar{\lambda} \bar{x}^T x \Rightarrow \lambda = \bar{\lambda}$. Hence λ is real

② If λ is an eigenvalue of a matrix A , then $\frac{1}{\lambda}$ ($\lambda \neq 0$) is the eigenvalue of A^{-1} . [N/D-2014]

Proof: Given λ is an eigenvalue of a matrix A . Let the corresponding [M/J-2012]

eigenvector be x . Then we have $Ax = \lambda x$

Pre-multiplying both sides by A^{-1} , we get

$$A^{-1}Ax = A^{-1}\lambda x$$

$$Ix = \lambda A^{-1}x$$

$$x = \lambda A^{-1}x$$

$$\div \lambda \Rightarrow \frac{1}{\lambda}x = A^{-1}x$$

From this we get, $\frac{1}{\lambda}$ is an eigenvalue of A^{-1} .

(23) If λ_i for $(i=1, 2, \dots, n)$ are the non-zero eigenvalues of A , then prove that $k\lambda_i$ are the eigenvalues of kA , where k being a non-zero scalar. [M/J-2012]

Proof: Given λ_i ($i=1, 2, \dots, n$) are the non-zero eigenvalues of A . Let the corresponding eigenvectors be x_i ($i=1, 2, \dots, n$). Then we have

$$Ax_i = \lambda_i x_i \quad (i=1, 2, \dots, n)$$

Pre-multiplying both sides by k , we get

$$kAx_i = k\lambda_i x_i$$

From this we get $k\lambda_i$ ($i=1, 2, \dots, n$) are the eigenvalues of kA .

(24) If $\lambda_1, \lambda_2, \dots, \lambda_n$ are the eigenvalues of a matrix A , then A^m has the eigenvalues $\lambda_1^m, \lambda_2^m, \dots, \lambda_n^m$. (m being a +ve integer)

Proof: Given λ_i ($i=1, 2, \dots, n$) are the eigenvalues of A . Let the corresponding eigenvectors be x_i ($i=1, 2, \dots, n$). Then we have

$$Ax_i = \lambda_i x_i \quad (i=1, 2, \dots, n)$$

$$A^2 x_i = A \lambda_i x_i = \lambda_i A x_i = \lambda_i (\lambda_i x_i) \quad (\because \text{by } \textcircled{1})$$

$$A^2 x_i = \lambda_i^2 x_i$$

$$\text{Similarly we get, } A^3 x_i = \lambda_i^3 x_i$$

$$\text{In general, } A^m x_i = \lambda_i^m x_i$$

From this we get, $\lambda_1^m, \lambda_2^m, \dots, \lambda_n^m$ are the eigenvalues of A^m .

(25) Find the sum & product of the eigenvalues of the matrix $\begin{pmatrix} -2 & 2 & -3 \\ 2 & 1 & -6 \\ -1 & -2 & 0 \end{pmatrix}$.

Sol:

$$\text{Sum of the eigenvalues} = \text{Sum of the main diagonal elements} = -2 + 1 + 0 = -1$$

Product of the eigenvalue = $|A| = -2(0-12) - 2(0-6) - 3(-4+1) = 24+12+9 = 45$

(26) The product of 2 eigenvalues of the matrix $A = \begin{pmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{pmatrix}$ is 16. Find the third eigenvalue.

Sol: Given $\lambda_1 \lambda_2 = 16$ — (1)

Product of eigenvalues = $|A| = 6(9-1) + 2(-6+2) + 2(2-6) = 48-8-8 = 32$

$$\lambda_1 \lambda_2 \lambda_3 = 32$$

$$16 \lambda_3 = 32 \quad (\because \text{by (1)})$$

$$\lambda_3 = \frac{32}{16} = 2 \quad \therefore \lambda_3 = 2$$

(27) Two of the eigenvalues of $A = \begin{pmatrix} 3 & -1 & 1 \\ -1 & 5 & -1 \\ 1 & -1 & 3 \end{pmatrix}$ are 3 & 6. Find the eigenvalues of A^{-1} .

Sol: Given $\lambda_1 = 3$ & $\lambda_2 = 6$

Sum of the eigenvalues = Sum of the main diagonal elements

$$\lambda_1 + \lambda_2 + \lambda_3 = 3 + 5 + 3 = 11$$

$$3 + 6 + \lambda_3 = 11 \Rightarrow 9 + \lambda_3 = 11 \Rightarrow \lambda_3 = 11 - 9 = 2$$

Hence the eigenvalues of A^{-1} are $\frac{1}{3}, \frac{1}{6}$ & $\frac{1}{2}$.

(28) Find the eigenvalues of A^3 given $A = \begin{pmatrix} 1 & 2 & 3 \\ 0 & 2 & -7 \\ 0 & 0 & 3 \end{pmatrix}$.

Sol: Given matrix A is a upper triangular matrix.

\therefore Eigenvalues of A are 1, 2 & 3. (Entries of main diagonal elements)

Hence the eigenvalues of A^3 are $1^3, 2^3$ & 3^3 (\bar{u}) 1, 8 & 27.

(29) The eigenvectors of a 3×3 real symmetric matrix A corresponding to the eigenvalues 2, 3, 6 are $[1, 0, -1]^T$, $[1, 1, 1]^T$ & $[-1, 2, -1]^T$ respectively, find the matrix A .

Sol: $D = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 6 \end{pmatrix}$

$$N = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{6}} \\ 0 & \frac{1}{\sqrt{3}} & \frac{2}{\sqrt{6}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{6}} \end{pmatrix}$$

$$A = NDN^T = \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{3} & -1/\sqrt{6} \\ 0 & 1/\sqrt{3} & 2/\sqrt{6} \\ -1/\sqrt{2} & 1/\sqrt{3} & -1/\sqrt{6} \end{pmatrix} \begin{pmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 6 \end{pmatrix} \begin{pmatrix} 1/\sqrt{2} & 0 & -1/\sqrt{2} \\ 1/\sqrt{3} & 1/\sqrt{3} & 1/\sqrt{3} \\ -1/\sqrt{6} & 2/\sqrt{6} & -1/\sqrt{6} \end{pmatrix}$$

$$= \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{3} & -1/\sqrt{6} \\ 0 & 1/\sqrt{3} & 2/\sqrt{6} \\ -1/\sqrt{2} & 1/\sqrt{3} & -1/\sqrt{6} \end{pmatrix} \begin{pmatrix} 2/\sqrt{2} & 0 & -2/\sqrt{2} \\ 3/\sqrt{3} & 3/\sqrt{3} & 3/\sqrt{3} \\ -6/\sqrt{6} & 12/\sqrt{6} & -6/\sqrt{6} \end{pmatrix}$$

$$\therefore A = \begin{pmatrix} 1+1+1 & 1-2 & -1+1+1 \\ 1-2 & 1+1 & 1-2 \\ -1+1+1 & 1-2 & 1+1+1 \end{pmatrix} = \begin{pmatrix} 3 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 3 \end{pmatrix}$$

Diagonalisation of non-symmetric matrix:

30 Diagonalise the matrix $\begin{pmatrix} 1 & -2 \\ -5 & 4 \end{pmatrix}$.

Sol: Let $A = \begin{pmatrix} 1 & -2 \\ -5 & 4 \end{pmatrix}$

Characteristic eqn: $\lambda^2 - S_1\lambda + S_2 = 0$

$S_1 =$ Sum of the main diagonal elements $= 1+4 = 5$

$S_2 = |A| = 4 - 10 = -6$

Hence the characteristic eqn. is $\lambda^2 - 5\lambda - 6 = 0$

$$\begin{array}{r|l} x & + \\ -6 & -5 \\ \hline -6 & +1 \\ \lambda-6 & \lambda+1 \end{array}$$

$$(\lambda - 6)(\lambda + 1) = 0$$

$$\therefore \lambda = -1, 6$$

Hence the eigenvalues are -1 & 6 .

Eigenvectors: $(A - \lambda I)x = 0$

$$\begin{pmatrix} 1-\lambda & -2 \\ -5 & 4-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 0$$

$$\frac{\lambda = -1}{\begin{pmatrix} 2 & -2 \\ -5 & 5 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 0} \quad \begin{array}{l} 2x_1 - 2x_2 = 0 \Rightarrow x_1 - x_2 = 0 \Rightarrow \frac{x_1}{1} = \frac{x_2}{1} \\ -5x_1 + 5x_2 = 0 \Rightarrow x_1 - x_2 = 0 \end{array}$$

$$\therefore x_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$\frac{\lambda = 6}{\begin{pmatrix} -5 & -2 \\ -5 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 0} \quad \begin{array}{l} -5x_1 - 2x_2 = 0 \Rightarrow -5x_1 = 2x_2 \Rightarrow \frac{x_1}{2} = \frac{x_2}{-5} \\ -5x_1 - 2x_2 = 0 \end{array}$$

$$\therefore x_2 = \begin{pmatrix} 2 \\ -5 \end{pmatrix}$$

Eigenvector matrix: $P = \begin{pmatrix} 1 & 2 \\ 1 & -5 \end{pmatrix}$

$$P^{-1} = \frac{1}{|P|} \text{Adj } P = \frac{1}{-5-2} \begin{pmatrix} -5 & -2 \\ -1 & 1 \end{pmatrix} = \frac{1}{-7} \begin{pmatrix} -5 & -2 \\ -1 & 1 \end{pmatrix}$$

$$AP = \begin{pmatrix} 1 & -2 \\ -5 & 4 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 1 & -5 \end{pmatrix} = \begin{pmatrix} -1 & 12 \\ -1 & -30 \end{pmatrix}$$

$$D = P^{-1}AP = \frac{-1}{7} \begin{pmatrix} -5 & -2 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} -1 & 12 \\ -1 & -30 \end{pmatrix} = \frac{-1}{7} \begin{pmatrix} 7 & 0 \\ 0 & -42 \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & 6 \end{pmatrix}$$

③ Reduce the matrix $\begin{pmatrix} -1 & 2 & -2 \\ 1 & 2 & 1 \\ -1 & -1 & 0 \end{pmatrix}$ to the diagonal form.

Sol:

$$\text{Let } A = \begin{pmatrix} -1 & 2 & -2 \\ 1 & 2 & 1 \\ -1 & -1 & 0 \end{pmatrix}$$

$$\text{Characteristic eqn.: } \lambda^3 - S_1\lambda^2 + S_2\lambda - S_3 = 0$$

$$S_1 = -1 + 2 + 0 = 1$$

$$S_2 = \begin{vmatrix} 2 & 1 \\ -1 & 0 \end{vmatrix} + \begin{vmatrix} -1 & -2 \\ -1 & 0 \end{vmatrix} + \begin{vmatrix} -1 & 2 \\ 1 & 2 \end{vmatrix} = (0+1) + (0-2) + (-2-2) = 1-2-4 = -5$$

$$S_3 = |A| = -1(0+1) - 2(0+1) - 2(-1+2) = -1-2-2 = -5$$

Hence the characteristic eqn. is $\lambda^3 - \lambda^2 - 5\lambda + 5 = 0$

$$\lambda = 1 \left| \begin{array}{ccc|c} 1 & -1 & -5 & 5 \\ & 1 & 0 & -5 \\ & 1 & 0 & -5 \\ \hline 1 & 0 & -5 & 0 \end{array} \right.$$

$$\lambda^2 + 0\lambda - 5 = 0 \Rightarrow \lambda^2 = 5 \Rightarrow \lambda = \pm\sqrt{5}$$

$$\therefore \lambda = -\sqrt{5}, 1, \sqrt{5}$$

Hence the eigenvalues are $-\sqrt{5}, 1$ & $\sqrt{5}$.

Eigenvectors: $(A - \lambda I)x = 0$

$$\begin{pmatrix} -1-\lambda & 2 & -2 \\ 1 & 2-\lambda & 1 \\ -1 & -1 & -\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\lambda = -\sqrt{5}$$

$$\begin{pmatrix} -1+\sqrt{5} & 2 & -2 \\ 1 & 2+\sqrt{5} & 1 \\ -1 & -1 & \sqrt{5} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$(-1+\sqrt{5})x_1 + 2x_2 - 2x_3 = 0$$

$$x_1 + (2+\sqrt{5})x_2 + x_3 = 0$$

$$-x_1 - x_2 + \sqrt{5}x_3 = 0$$

$$\frac{x_1}{2\sqrt{5}+5+1} = \frac{x_2}{-1-\sqrt{5}} = \frac{x_3}{-1+2+\sqrt{5}}$$

$$\frac{x_1}{2\sqrt{5}+6} = \frac{x_2}{-1-\sqrt{5}} = \frac{x_3}{1+\sqrt{5}}$$

$$\therefore x_1 = \begin{pmatrix} 2\sqrt{5}+6 \\ -1-\sqrt{5} \\ 1+\sqrt{5} \end{pmatrix}$$

$$\begin{matrix} \textcircled{2} & \textcircled{1} & \textcircled{3} \\ 2+\sqrt{5} & 1 & 1 & 2+\sqrt{5} \\ -1 & \sqrt{5} & -1 & -1 \end{matrix} \begin{matrix} x_1 \\ x_2 \\ x_3 \end{matrix}$$

$$\lambda = 1 \begin{pmatrix} -2 & 2 & -2 \\ 1 & 1 & 1 \\ -1 & -1 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0 \quad \begin{matrix} -2x_1 + 2x_2 - 2x_3 = 0 \\ x_1 + x_2 + x_3 = 0 \\ -x_1 - x_2 - x_3 = 0 \end{matrix} \quad \begin{matrix} x_1 & x_2 & x_3 \\ 2 & -2 & -2 \\ 1 & 1 & 1 \end{matrix}$$

$$\frac{x_1}{2+2} = \frac{x_2}{-2+2} = \frac{x_3}{-2-2} \Rightarrow \frac{x_1}{4} = \frac{x_2}{0} = \frac{x_3}{-4} \Rightarrow \frac{x_1}{1} = \frac{x_2}{0} = \frac{x_3}{-1}$$

$$\therefore X_2 = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

$$\lambda = \sqrt{5} \begin{pmatrix} -1-\sqrt{5} & 2 & -2 \\ 1 & 2-\sqrt{5} & 1 \\ -1 & -1 & -\sqrt{5} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0 \quad \begin{matrix} (-1-\sqrt{5})x_1 + 2x_2 - 2x_3 = 0 \\ x_1 + (2-\sqrt{5})x_2 + x_3 = 0 \\ -x_1 - x_2 - \sqrt{5}x_3 = 0 \end{matrix}$$

$$\frac{x_1}{-2\sqrt{5}+5+1} = \frac{x_2}{-1+\sqrt{5}} = \frac{x_3}{-1+2-\sqrt{5}}$$

$$\frac{x_1}{6-2\sqrt{5}} = \frac{x_2}{-1+\sqrt{5}} = \frac{x_3}{1-\sqrt{5}}$$

$$\therefore X_3 = \begin{pmatrix} 6-2\sqrt{5} \\ -1+\sqrt{5} \\ 1-\sqrt{5} \end{pmatrix}$$

$$\textcircled{2} \text{ \& } \textcircled{3} \quad \begin{matrix} x_1 & x_2 & x_3 \\ 2-\sqrt{5} & 1 & 1 \\ -1 & -\sqrt{5} & -1 \end{matrix} \quad \begin{matrix} 2-\sqrt{5} \\ -1 \\ 2-\sqrt{5} \end{matrix}$$

Eigenvector matrix: $P = \begin{pmatrix} 2\sqrt{5}+6 & 1 & 6-2\sqrt{5} \\ -1-\sqrt{5} & 0 & -1+\sqrt{5} \\ 1+\sqrt{5} & -1 & 1-\sqrt{5} \end{pmatrix}$

$$AP = \begin{pmatrix} -23.4164 & 1 & 3.4164 \\ 7.2361 & 0 & 2.7639 \\ -7.2361 & -1 & -2.7639 \end{pmatrix}$$

$$P^{-1} = \begin{pmatrix} 0.0691 & -0.0163 & 0.0691 \\ 0 & -1 & -1 \\ 0.1809 & 0.7663 & 0.1809 \end{pmatrix}$$

$$\therefore D = P^{-1}AP = \begin{pmatrix} -2.236 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2.236 \end{pmatrix} = \begin{pmatrix} -\sqrt{5} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \sqrt{5} \end{pmatrix}$$

32) Diagonalise the matrix $A = \begin{pmatrix} 0 & -2 & -2 \\ -1 & 1 & 2 \\ -1 & -1 & 2 \end{pmatrix}$

Sol: Given $A = \begin{pmatrix} 0 & -2 & -2 \\ -1 & 1 & 2 \\ -1 & -1 & 2 \end{pmatrix}$

Characteristic eqn: $\lambda^3 - 5\lambda^2 + 5\lambda - 5 = 0$

$$S_1 = 0 + 1 + 2 = 3$$

$$S_2 = \begin{vmatrix} 1 & 2 \\ -1 & 2 \end{vmatrix} + \begin{vmatrix} 0 & -2 \\ -1 & 2 \end{vmatrix} + \begin{vmatrix} 0 & -2 \\ -1 & 1 \end{vmatrix} = (2+2) + (0-2) + (0-2) = 4-2-2 = 0$$

$$S_3 = 0 + 2(-2+2) - 2(1+1) = -4$$

Hence the characteristic eqn. is $\lambda^3 - 3\lambda^2 + 4 = 0$

$$\lambda = 2 \left| \begin{array}{ccc|c} 1 & -3 & 0 & 4 \\ & 2 & -2 & -4 \\ \hline 1 & -1 & -2 & 0 \end{array} \right.$$

$$\lambda^2 - \lambda - 2 = 0$$

$$(\lambda+1)(\lambda-2) = 0$$

$$\therefore \lambda = -1, 2, 2$$

Hence the eigenvalues are $-1, 2$ & 2 .

Eigenvectors: $(A - \lambda I)x = 0$

$$\begin{pmatrix} -\lambda & -2 & -2 \\ -1 & 1-\lambda & 2 \\ -1 & -1 & 2-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0$$

$$\lambda = -1$$

$$\begin{pmatrix} 1 & -2 & -2 \\ -1 & 2 & 2 \\ -1 & -1 & 3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0 \quad \begin{array}{l} x_1 - 2x_2 - 2x_3 = 0 \\ -x_1 + 2x_2 + 2x_3 = 0 \\ -x_1 - x_2 + 3x_3 = 0 \end{array}$$

$$\frac{x_1}{6+2} = \frac{x_2}{-2+3} = \frac{x_3}{1+2} \Rightarrow \frac{x_1}{8} = \frac{x_2}{1} = \frac{x_3}{3}$$

(2) & (3)

$$\begin{array}{cccc} x_1 & x_2 & x_3 & \\ 2 & 2 & -1 & 2 \\ -1 & 3 & -1 & -1 \end{array}$$

$$\therefore x_1 = \begin{pmatrix} 8 \\ 1 \\ 3 \end{pmatrix}$$

$$\lambda = 2$$

$$\begin{pmatrix} -2 & -2 & -2 \\ -1 & -1 & 2 \\ -1 & -1 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = 0 \quad \begin{array}{l} -2x_1 - 2x_2 - 2x_3 = 0 \\ -x_1 - x_2 + 2x_3 = 0 \\ -x_1 - x_2 + 0x_3 = 0 \end{array}$$

$$\frac{x_1}{-4-2} = \frac{x_2}{2+2} = \frac{x_3}{2-2} \Rightarrow \frac{x_1}{-6} = \frac{x_2}{4} = \frac{x_3}{0} \Rightarrow \frac{x_1}{-3} = \frac{x_2}{2} = \frac{x_3}{0}$$

$$\therefore x_2 = \begin{pmatrix} -3 \\ 2 \\ 0 \end{pmatrix}$$

We get one eigenvector corresponding to the repeated root 2.

\therefore Diagonalisation not possible.

DIFFERENTIAL CALCULUS

Representation of function:

- (i) Verbally (by a description in words)
- (ii) Visually (by a graph)
- (iii) Numerically (by a table of values)
- (iv) Algebraically (by an explicit formula)

Definition: (Real-valued functions)


A function, whose domain & co-domain are subsets of the set of all real numbers, is known as real-valued function.

Definition:

Let $f: A \rightarrow B$, then set A is called the domain of the function & set B is called the co-domain of the function.

The set of all the images of all the elements of A under the function f is called the range of f is denoted by $f(A)$.

Thus the range of f is $f(A) = \{f(x) : x \in A\}$.



Definition: (Explicit function)

If x & y be so related that y can be expressed explicitly in terms of x , then y is called explicit function of x .

E.g: $y = x^2 - 4x + 2$

Definition: (Implicit function)

If x & y be so related that y cannot be expressed explicitly in terms of x , then y is called implicit function of x .

E.g: $x^3 + y^3 - 3xy = 0$.


Problems:

① Find the domain & range & sketch the graph of the function $f(x) = x^2$.

Sol:

Given $f(x) = x^2$

Domain (x)	$-\infty$	-2	-1	0	1	2	∞
Range (f(x))	∞	4	1	0	1	4	∞



Domain = $(-\infty, \infty)$

Range = $[0, \infty)$

② Find the domain & range of $f(x) = \sqrt{5x+10}$.

Sol: Given $f(x) = \sqrt{5x+10}$

Here $5x+10 \geq 0$

$\Rightarrow 5x \geq -10$

$\Rightarrow x \geq \frac{-10}{5}$

$\Rightarrow x \geq -2$

Domain (x)	-2	-1	0	1	2	∞
Range (f(x))	0	$\sqrt{5}$	$\sqrt{10}$	$\sqrt{15}$	$\sqrt{20}$	∞

Domain = $[-2, \infty)$

Range = $[0, \infty)$

③ Find the domain of $f(x) = \frac{x+4}{x^2-9}$.

Sol: Given $f(x) = \frac{x+4}{x^2-9}$.

$$x^2-9=0 \Rightarrow x^2=9 \Rightarrow x=\sqrt{9}=\pm 3$$

$$\text{Domain} = (-\infty, -3) \cup (-3, 3) \cup (3, \infty).$$

④ Find the domain of $f(x) = \frac{1}{\sqrt[4]{x^2-5x}}$.

Sol: Given $f(x) = \frac{1}{\sqrt[4]{x^2-5x}}$

For $x=0$, we get $x^2-5x=0-0=0$

For $x=5$, we get $x^2-5x=25-25=0$

$$\text{Domain} = (-\infty, 0) \cup (5, \infty).$$

⑤ Find the domain & sketch the graph of the function

$$f(x) = \begin{cases} x+2 & \text{if } x < 0 \\ 1-x & \text{if } x \geq 0 \end{cases}$$

Sol: Given $f(x) = \begin{cases} x+2 & \text{if } x < 0 \\ 1-x & \text{if } x \geq 0 \end{cases}$

$x < 0$


x	-1	-2	-3	-4	...
$f(x)$	1	0	-1	-2	...

$f(x) = x+2$

$x \geq 0$

x	0	1	2	3	...
$f(x)$	1	0	-1	-2	...

$f(x) = 1-x$



$$\text{Domain} = (-\infty, \infty)$$

(10) (6) Find the domain of $f(x) = \sqrt{3-x} - \sqrt{2+x}$.

Sol: Given $f(x) = \sqrt{3-x} - \sqrt{2+x}$

Here $3-x \geq 0$ & $2+x \geq 0$


$\Rightarrow 3 \geq x$ & $x \geq -2$

$\Rightarrow -2 \leq x \leq 3$

Domain = $[-2, 3]$

Definition:

Even function: $f(-x) = f(x)$ [or] symmetric about the y-axis



E.g: ① $f(x) = 1 - x^4$

$f(-x) = 1 - (-x)^4 = 1 - x^4 = f(x)$


$\therefore f(-x) = f(x)$

Hence $f(x) = 1 - x^4$ is an even function.

② $f(x) = \cos x$

$f(-x) = \cos(-x) = \cos x = f(x)$

$\therefore f(x) = \cos x$ is an even function.



Odd function: $f(-x) = -f(x)$ [or] symmetric about the x-axis

E.g: ① $f(x) = x^5 + x$

$f(-x) = (-x)^5 + (-x) = -x^5 - x = -(x^5 + x) = -f(x)$

$\therefore f(-x) = -f(x)$

Hence $f(x) = x^5 + x$ is an odd function.

② $f(x) = \sin x$

$f(-x) = \sin(-x) = -\sin x = -f(x)$

Hence $f(x) = \sin x$ is an odd function.

Example for neither even nor odd function:

① $f(x) = \frac{1}{x-1}$

$f(-x) = \frac{1}{-x-1} \neq f(x) \neq -f(x)$

Hence the given function is neither even nor odd.

② $f(x) = e^x$

$f(-x) = e^{-x} \neq f(x) \neq -f(x)$

Hence $f(x) = e^x$ is neither even nor odd function.

H.w

① Find the domain of $f(x) = \sqrt{x+2}$.

② Find the domain of $f(x) = \frac{1}{x^2 - x}$.

(15)

Limit of a function:

$\lim_{x \rightarrow a} f(x) = l$ is $f(x) \rightarrow l$ as $x \rightarrow a$ (or) $f(x)$ approaches l as

x approaches a .

Left-hand limit:

$$\lim_{x \rightarrow a^-} f(x) = l$$

Here $x \rightarrow a^-$ means $x < a$.

Right-hand limit:

$$\lim_{x \rightarrow a^+} f(x) = l$$

Here $x \rightarrow a^+$ means $x > a$.

Definition:

$\lim_{x \rightarrow a} f(x) = l$ if & only if $\lim_{x \rightarrow a^-} f(x) = l$ & $\lim_{x \rightarrow a^+} f(x) = l$.

Problems:

⑦ Guess the value of $\lim_{x \rightarrow 1} \frac{x-1}{x^2-1}$.

Sol: Here $f(x) = \frac{x-1}{x^2-1}$.

$x < 1$

x	$f(x)$
0.5	0.66667
0.6	0.625
0.7	0.58824
0.8	0.55556
0.9	0.52632
0.99	0.50251
0.999	0.50025

$x > 1$

x	$f(x)$
1.5	0.4
1.4	0.41667
1.3	0.43478
1.2	0.45455
1.1	0.47619
1.01	0.49751
1.001	0.49975

$$\therefore \lim_{x \rightarrow 1} \frac{x-1}{x^2-1} = 0.5$$

Ans

① Guess the value of $\lim_{x \rightarrow 0} \frac{\sin x}{x}$.

Q8 Guess the value of the limit (if it exists) for the function $\lim_{x \rightarrow 0} \frac{e^{5x} - 1}{x}$ by evaluating the function at the given numbers, $x = \pm 0.5, \pm 0.1, \pm 0.01, \pm 0.001, \pm 0.0001$ (correct to 6 decimal places)

Sol: Here $f(x) = \frac{e^{5x} - 1}{x}$

x	f(x)
-0.5	1.83583
-0.1	3.934693
-0.01	4.877058
-0.001	4.987521
-0.0001	4.99875

x	f(x)
0.5	22.364988
0.1	6.487213
0.01	5.12711
0.001	5.012521
0.0001	5.00125

$\therefore \lim_{x \rightarrow 0} \frac{e^{5x} - 1}{x} = 5$

Q9 Evaluate $\lim_{t \rightarrow 1} \frac{t^4 - 1}{t^3 - 1}$.

Sol: $\lim_{t \rightarrow 1} \frac{t^4 - 1}{t^3 - 1} = \lim_{t \rightarrow 1} \frac{4t^3}{3t^2} = \frac{4}{3}$

$\frac{d}{dx}(x^n) = nx^{n-1}$
 $\frac{d}{dx}(c) = 0$ where c is a constant

Q10 Given that $\lim_{x \rightarrow 2} f(x) = 4$ & $\lim_{x \rightarrow 2} g(x) = -2$. Find the limit that exists for $\lim_{x \rightarrow 2} \left[\frac{3f(x)}{g(x)} \right]$.

Sol: Given $\lim_{x \rightarrow 2} f(x) = 4$ & $\lim_{x \rightarrow 2} g(x) = -2$.

$\therefore \lim_{x \rightarrow 2} \left[\frac{3f(x)}{g(x)} \right] = \frac{3(4)}{-2} = -6$

Q11 Sketch the graph of the function $f(x) = \begin{cases} 1+x, & x < -1 \\ x^2, & -1 \leq x \leq 1 \\ 2-x, & x \geq 1 \end{cases}$ & use it

to determine the value of 'a' for which $\lim_{x \rightarrow a} f(x)$ exists?

Sol:

Sol: $f(x) = 1+x, x < -1$ $f(x) = x^2, -1 \leq x \leq 1$ $f(x) = 2-x, x \geq 1$

x	-2	-3	-4
f(x)	-1	-2	-3

x	-1	0	1
f(x)	1	0	1

x	1	2	3
f(x)	1	0	-1

At x = -1

$$\lim_{x \rightarrow -1^-} f(x) = \lim_{x \rightarrow -1^-} (1+x) = 1+(-1) = 0$$

$$\lim_{x \rightarrow -1^+} f(x) = \lim_{x \rightarrow -1^+} x^2 = (-1)^2 = 1$$

$\therefore \lim_{x \rightarrow -1} f(x)$ doesn't exist.


At x = 1

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} x^2 = 1^2 = 1$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (2-x) = 2-1 = 1$$

$\therefore \lim_{x \rightarrow 1} f(x)$ exists.

Hence $\lim_{x \rightarrow a} f(x)$ exists for all 'a' except at $a = -1$.



⑫ Sketch the graph of the function $f(x) = \begin{cases} 1+\sin x & \text{if } x < 0 \\ \cos x & \text{if } 0 \leq x \leq \pi \\ \sin x & \text{if } x > \pi \end{cases}$ ← use it to

determine the value of 'a' for which $\lim_{x \rightarrow a} f(x)$ exists.

Sol:

	$f(x) = 1+\sin x, x < 0$		$f(x) = \cos x, 0 \leq x \leq \pi$			$f(x) = \sin x, x > \pi$	
x	$-\pi/2$	$-\pi$	0	$\pi/2$	π	$3\pi/2$	2π
f(x)	0	1	1	0	-1	-1	0

At x = 0

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (1+\sin x) = 1+\sin 0 = 1+0 = 1$$

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} \cos x = \cos 0 = 1$$

$\therefore \lim_{x \rightarrow 0} f(x)$ exists.


At $x = \pi$

$$\lim_{x \rightarrow \pi^-} f(x) = \lim_{x \rightarrow \pi^-} \cos x = \cos \pi = -1$$

$$\lim_{x \rightarrow \pi^+} f(x) = \lim_{x \rightarrow \pi^+} \sin x = \sin \pi = 0$$

$\therefore \lim_{x \rightarrow \pi} f(x)$ doesn't exist.

Hence $\lim_{x \rightarrow a} f(x)$ exists for all 'a' except at $a = \pi$.



(13) Check whether $\lim_{x \rightarrow -3} \frac{3x+9}{|x+3|}$ exist.

Sol: $\lim_{x \rightarrow -3^-} \frac{3x+9}{-(x+3)} = \lim_{x \rightarrow -3^-} \frac{3(x+3)}{-(x+3)} = -3$

$$\lim_{x \rightarrow -3^+} \frac{3x+9}{x+3} = \lim_{x \rightarrow -3^+} \frac{3(x+3)}{x+3} = 3. \text{ Here } \lim_{x \rightarrow -3^-} f(x) \neq \lim_{x \rightarrow -3^+} f(x)$$

$\therefore \lim_{x \rightarrow -3} f(x)$ doesn't exist.

Definition: (Continuity)

A function f is continuous at 'a' if $\lim_{x \rightarrow a} f(x) = f(a)$.

(i) If f is continuous at a, then

(i) $f(a)$ should exist

(ii) $\lim_{x \rightarrow a} f(x)$ exists both on the left & right.

(iii) $\lim_{x \rightarrow a} f(x) = f(a)$.

Eg: Polynomials, rational functions, root functions, trigonometric functions, inverse trigonometric functions, exponential functions, logarithmic functions.

(14) Find the numbers that at which f is discontinuous, at which of these numbers if f is continuous from the right from the left or neither? When $f(x) = \begin{cases} x+2, & x < 0 \\ e^x, & 0 \leq x \leq 1 \\ 2-x, & x > 1 \end{cases}$

Sol:

At x=0

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (x+2) = 0+2 = 2$$

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} e^x = e^0 = 1$$

$$f(0) = e^0 = 1$$

$$\therefore \lim_{x \rightarrow 0^+} f(x) = f(0) \neq \lim_{x \rightarrow 0^-} f(x).$$

Hence f is continuous on the right at $x=0$ & f is discontinuous on the left at $x=0$.

$\therefore f$ is discontinuous at $x=0$.

At x=1

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} e^x = e^1 = e$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (2-x) = 2-1 = 1$$

$$f(1) = e^1 = e$$

$$\therefore \lim_{x \rightarrow 1^-} f(x) = f(1) \neq \lim_{x \rightarrow 1^+} f(x)$$

Hence f is continuous on the left at $x=1$ & f is discontinuous on the right at $x=1$.

$\therefore f$ is discontinuous at $x=1$.

Thus f is continuous in $(-\infty, 0) \cup (0, 1) \cup (1, \infty)$.

(H.w)

① Find the domain where the function f is continuous. Also find the numbers at which the function f is discontinuous, where

$$f(x) = \begin{cases} 1+x^2, & x \leq 0 \\ 2-x, & 0 < x \leq 2 \\ (x-2)^2, & x > 2 \end{cases}$$

(Av)

② For what value of the constant b , is the function f continuous on

$$(-\infty, \infty) \text{ if } f(x) = \begin{cases} bx^2 + 2x & \text{if } x < 2 \\ x^3 - bx & \text{if } x \geq 2 \end{cases}$$

Sol: At $x=2$

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} (bx^2 + 2x) = 4b + 4$$

$$f(2) = (2)^3 - b(2) = 8 - 2b$$

Since f is continuous, $\lim_{x \rightarrow 2^-} f(x) = f(2)$

$$\Rightarrow 4b + 4 = 8 - 2b \Rightarrow 4b + 2b = 8 - 4$$

$$\Rightarrow 6b = 4 \Rightarrow b = \frac{4}{6} = \frac{2}{3}$$

$$\therefore \boxed{b = \frac{2}{3}}$$

(10) (16) Find the values of a & b that make f continuous on $(-\infty, \infty)$.

$$f(x) = \begin{cases} \frac{x^3 - 8}{x - 2}, & \text{if } x < 2 \\ ax^2 - bx + 3, & \text{if } 2 \leq x < 3 \\ 2x - a + b, & \text{if } x \geq 3 \end{cases}$$

$$\boxed{\frac{d(x^n)}{dx} = nx^{n-1}}$$

Sol: At $x=2$

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} \frac{x^3 - 8}{x - 2} = \lim_{x \rightarrow 2^-} \frac{3x^2}{1} = \lim_{x \rightarrow 2^-} 3x^2 = 3(2)^2 = 12$$

$$f(2) = a(2)^2 - b(2) + 3 = 4a - 2b + 3$$

Since f is continuous, $\lim_{x \rightarrow 2^-} f(x) = f(2)$

$$\Rightarrow 12 = 4a - 2b + 3 \Rightarrow 4a - 2b = 12 - 3 = 9 \Rightarrow 4a - 2b = 9 \text{ --- (1)}$$

At $x=3$

$$\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^-} ax^2 - bx + 3 = a(3)^2 - b(3) + 3 = 9a - 3b + 3$$

$$f(3) = 2(3) - a + b = 6 - a + b$$

Since f is continuous, $\lim_{x \rightarrow 3^-} f(x) = f(3)$

$$\Rightarrow 9a - 3b + 3 = 6 - a + b \Rightarrow 9a + a - 3b - b = 6 - 3$$

$$\Rightarrow 10a - 4b = 3 \text{ --- (2)}$$

$$\textcircled{1} \times 2 \Rightarrow 8a - 4b = 18$$

$$10a - 4b = 3 \text{ --- } \textcircled{2}$$

$$\begin{array}{r} (-) \quad (+) \quad (-) \\ \hline \end{array}$$

$$-2a = 15 \Rightarrow a = \frac{15}{-2}$$

$$\therefore a = \frac{-15}{2}$$

Substituting a value in $\textcircled{1}$, $4\left(\frac{-15}{2}\right) - 2b = 9$

$$\Rightarrow -30 - 2b = 9 \Rightarrow 2b = -30 - 9 = -39 \Rightarrow b = \frac{-39}{2}$$

Hence $a = \frac{-15}{2}$ & $b = \frac{-39}{2}$.

H.w

$$\textcircled{1} \text{ If } f(x) = \begin{cases} \frac{x^2 - 4}{x - 2}, & x < 2 \\ ax^2 - bx + 3, & 2 \leq x < 3 \\ 2x - a + b, & x \geq 3 \end{cases}$$

is continuous for all real x , find the

values of a & b .

Formulae:

$$\textcircled{1} \frac{d}{dx}(x^n) = nx^{n-1}$$

$$\textcircled{2} \frac{d}{dx}(c) = 0 \text{ where } c \text{ is constant.}$$

$$\textcircled{3} \frac{d}{dx}(c f(x)) = c \frac{d}{dx}(f(x))$$

$$\textcircled{4} \text{ Equation of tangent line is } y - y_1 = m(x - x_1) \text{ where } m = \frac{dy}{dx}$$

$$\textcircled{5} \text{ Equation of normal line is } y - y_1 = \frac{-1}{m}(x - x_1) \text{ where } m = \frac{dy}{dx}$$

Problems:

$\textcircled{17}$ Find the derivative of the following:-

(i) $f(x) = x^{1000}$

$$f'(x) = 1000x^{1000-1} = 1000x^{999}$$

(ii) $y = \frac{1}{x^2}$

$y = \frac{1}{x^2} = x^{-2}$

$y' = (-2)x^{-2-1} = (-2)x^{-3} = \frac{-2}{x^3}$

(iii) $y = \sqrt[3]{x^2}$

$y = (x^2)^{1/3} = x^{2/3}$

$y' = \frac{2}{3}x^{2/3-1} = \frac{2}{3}x^{-1/3} = \frac{2}{3x^{1/3}}$

(iv) $y = x^8 + 12x^5 - 4x^4 + 10x^3 - 6x + 7$

$y' = 8x^7 + 12(5x^4) - 4(4x^3) + 10(3x^2) - 6$

$y' = 8x^7 + 60x^4 - 16x^3 + 30x^2 - 6$

(v) $y = ax^{2n} + bx^n + c$

$y' = a(2n)x^{2n-1} + bnx^{n-1} = 2anx^{2n-1} + bnx^{n-1}$

(vi) $y = \frac{x^2 + 4x + 3}{\sqrt{x}}$

$y = x^{-1/2}(x^2 + 4x + 3) = x^{-1/2}x^2 + 4xx^{-1/2} + 3x^{-1/2} = x^{3/2} + 4x^{1/2} + 3x^{-1/2}$

$y' = \frac{3}{2}x^{3/2-1} + \frac{1}{2} \times 4x^{1/2-1} + 3(-1/2)x^{-1/2-1}$

$= \frac{3}{2}x^{1/2} + 2x^{-1/2} - \frac{3}{2}x^{-3/2} = \frac{3}{2}\sqrt{x} + \frac{2}{\sqrt{x}} - \frac{3}{2}x^{-3/2}$

10 18 Does the curve $y = x^4 - 2x^2 + 2$ have any horizontal tangents? If so where?

Sol: Given $y = x^4 - 2x^2 + 2$

Horizontal tangents occur where the derivative is zero.

(a) $\frac{dy}{dx} = 0 \Rightarrow 4x^3 - 4x = 0 \Rightarrow 4x(x^2 - 1) = 0$

$\Rightarrow x = 0, x^2 - 1 = 0 \Rightarrow x^2 = 1 \Rightarrow x = \pm 1$

$\therefore x = 0, 1, -1$

x	-1	0	1
y	1	2	1

Hence the corresponding points are (-1, 1), (0, 2) & (1, 1).

(19) The equation of motion of a particle is $s = 2t^3 - 5t^2 + 3t + 7$, where s is measured in centimeters & t in seconds. Find the acceleration as a function of time. What is the acceleration after 2 seconds?

Sol:
Velocity = $\frac{ds}{dt} = 6t^2 - 10t + 3$

Acceleration = $\frac{d^2s}{dt^2} = 12t - 10$

$\left[\frac{d^2s}{dt^2}\right]_{t=2} = 12(2) - 10 = 24 - 10 = 14$

H.w Problem:

① Find the derivative of the following functions:

(i) $y = \sqrt{x}$

(ii) $y = x\sqrt{2}$

(iii) $y = x^2(1-2x)$

(iv) $y = x^{2.4} + e^{2.4}$

Formulae:

① $\frac{d}{dx}(e^x) = e^x$

② $\frac{d}{dx}(e^{2x}) = e^{2x} \cdot 2 = 2e^{2x}$

Problems:

② Find the derivative of the following functions:

(i) $y = 3e^x + \frac{4}{\sqrt[3]{x}}$

$y = 3e^x + \frac{4}{x^{1/3}} = 3e^x + 4x^{-1/3}$

$y' = 3e^x + 4(-1/3)x^{-1/3-1} = 3e^x - \frac{4}{3}x^{-4/3}$

(ii) $y = a^x$

$y = a^x = e^{\log a^x} = e^{x \log a} = e^{(\log a)x}$

$y' = e^{(\log a)x} \cdot \log a = \log a \cdot e^{x(\log a)} = \log a \cdot e^{\log a^x} = \log a \cdot a^x = a^x \log a$

H.w

① Find the derivative of the following functions:

(i) $y = e^x - x$

(ii) $y = 2^x$

(iii) $y = e^{-x} - 7$

Formulae:

① $\frac{d}{dx}(uv) = uv' + vu'$

② $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{vu' - uv'}{v^2}$

② Find $f'(x)$ & $f''(x)$ of $f(x) = x^4 e^x$.

Sol: Given $f(x) = x^4 e^x$

$f'(x) = x^4(e^x) + e^x(4x^3)$
 $= x^4 e^x + 4e^x x^3 = e^x(x^4 + 4x^3)$

$f''(x) = e^x(4x^3 + 12x^2) + (x^4 + 4x^3)e^x$
 $= e^x(4x^3 + 12x^2 + x^4 + 4x^3)$
 $= e^x(x^4 + 8x^3 + 12x^2)$

$u = x^4, v = e^x$
 $u' = 4x^3, v' = e^x$
 $d(uv) = uv' + vu'$

$u = e^x, v = x^4 + 4x^3$
 $u' = e^x, v' = 4x^3 + 12x^2$

② If $f(x) = \frac{x^2}{1+2x}$, then find $f'(x)$ & $f''(x)$.

Sol: Given $f(x) = \frac{x^2}{1+2x}$

$f'(x) = \frac{(1+2x)(2x) - x^2(2)}{(1+2x)^2}$
 $= \frac{2x + 4x^2 - 2x^2}{(1+2x)^2} = \frac{2x + 2x^2}{(1+2x)^2}$

$f''(x) = \frac{(1+2x)^2(4x+2) - (2x^2+2x)4(1+2x)}{(1+2x)^4}$
 $= \frac{(1+2x)[(1+2x)(4x+2) - 4(2x^2+2x)]}{(1+2x)^4}$
 $= \frac{4x+2+8x^2+4x-8x^2-8x}{(1+2x)^3} = \frac{2}{(1+2x)^3}$

$d\left(\frac{u}{v}\right) = \frac{vu' - uv'}{v^2}$
 $u = x^2, v = 1+2x$
 $u' = 2x, v' = 2$

$u = 2x^2 + 2x, v = (1+2x)^2$
 $u' = 4x + 2, v' = 2(1+2x) \cdot 2$
 $v' = 4(1+2x)$

(Au) ② If $f(x) = xe^x$ then find the expression for $f''(x)$.

Sol: Given $f(x) = xe^x$

$f'(x) = xe^x + e^x(1) = xe^x + e^x$

$f''(x) = xe^x + e^x(1) + e^x = xe^x + 2e^x = e^x(x+2)$

$u = x, v = e^x$
 $u' = 1, v' = e^x$
 $d(uv) = uv' + vu'$

10

(24) Find $\frac{dy}{dx}$ if $y = x^2 e^{2x} (x^2 + 1)^4$.

Sol: Given $y = x^2 e^{2x} (x^2 + 1)^4$.

$$d(uv) = uv' + v u'$$

$$u = x^2 e^{2x}$$

$$v = (x^2 + 1)^4$$

$$u' = x^2 (e^{2x} \cdot 2) + e^{2x} (2x)$$

$$v' = 4(x^2 + 1)^3 (2x)$$

$$\frac{dy}{dx} = x^2 e^{2x} (4(x^2 + 1)^3 (2x)) + (x^2 + 1)^4 (2x^2 e^{2x} + 2x e^{2x})$$

$$= (x^2 + 1)^3 [8x^3 e^{2x} + (x^2 + 1) 2x e^{2x} (x + 1)]$$

$$= (x^2 + 1)^3 2x e^{2x} [4x^2 + (x^2 + 1)(x + 1)]$$

$$= (x^2 + 1)^3 2x e^{2x} [4x^2 + x^3 + x^2 + x + 1] = (x^2 + 1)^3 2x e^{2x} (x^3 + 5x^2 + x + 1)$$

$$= 2x e^{2x} (x^2 + 1)^3 (x^3 + 5x^2 + x + 1)$$

10

(25) If $f(x) = \frac{1-x}{2+x}$ then find the equation for $f'(x)$ using the concept of derivatives.

Sol: Given $f(x) = \frac{1-x}{2+x}$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{\frac{1-(x+h)}{2+(x+h)} - \frac{1-x}{2+x}}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(1-x-h)(2+x) - (1-x)(2+x+h)}{h(2+x)(2+x+h)}$$

$$= \lim_{h \rightarrow 0} \frac{2+x-2x-x^2-2h-xh - (2+x+h-2x-x^2-xh)}{h(2+x)(2+x+h)}$$

$$= \lim_{h \rightarrow 0} \frac{2+x-2x-x^2-2h-xh - 2-x-h+2x+x^2+xh}{h(2+x)(2+x+h)}$$

$$= \lim_{h \rightarrow 0} \frac{-3h}{h(2+x)(2+x+h)} = \lim_{h \rightarrow 0} \frac{-3}{(2+x)(2+x+h)}$$

$$= \frac{-3}{(2+x)(2+x)} = \frac{-3}{(2+x)^2}$$

Q Differentiate the following functions

- (i) $f(x) = (x^3 + 2x)e^x$ (ii) $f(x) = \sqrt{x}(a+bx)$
- (iii) $f(x) = \frac{x^2+x-2}{x^3+6}$ (iv) $f(x) = \frac{e^x}{x}$

Formulae:

- ① $\frac{d}{dx}(\sin x) = \cos x$ ② $\frac{d}{dx}(\cos x) = -\sin x$
- ③ $\frac{d}{dx}(\tan x) = \sec^2 x$ ④ $\frac{d}{dx}(\cot x) = -\operatorname{cosec}^2 x$
- ⑤ $\frac{d}{dx}(\operatorname{cosec} x) = -\operatorname{cosec} x \cot x$ ⑥ $\frac{d}{dx}(\sec x) = \sec x \tan x$
- ⑦ $\frac{d}{dx}(\sin mx) = m \cos mx$ ⑧ $\frac{d}{dx}(\cos mx) = -m \sin mx$
- ⑨ $\operatorname{cosec} x = \frac{1}{\sin x}$ ⑩ $\sec x = \frac{1}{\cos x}$
- ⑪ $\cot x = \frac{1}{\tan x}$ ⑫ $\tan x = \frac{\sin x}{\cos x}$
- ⑬ $\sin^2 x + \cos^2 x = 1$ ⑭ $1 + \tan^2 x = \sec^2 x$
- ⑮ $1 + \cot^2 x = \operatorname{cosec}^2 x$

Problems:

⑳ Find the derivative of the following:

(i) $y = \operatorname{cosec} x + e^x \cot x$

$$y' = -\operatorname{cosec} x \cot x + [-e^x \operatorname{cosec}^2 x + e^x \cot x]$$

$$= -\operatorname{cosec} x \cot x + e^x (-\operatorname{cosec}^2 x + \cot x)$$

$u = e^x$	$v = \cot x$
$u' = e^x$	$v' = -\operatorname{cosec}^2 x$
$d(uv) = uv' + v u'$	

(ii) $y = \frac{\sec x}{1 + \tan x}$

$$y' = \frac{(1 + \tan x) \sec x \tan x - \sec x \sec^2 x}{(1 + \tan x)^2}$$

$$= \frac{\sec x [\tan x + \tan^2 x - \sec^2 x]}{(1 + \tan x)^2} = \frac{\sec x (\tan x - 1)}{(1 + \tan x)^2} \quad (\because 1 + \tan^2 x = \sec^2 x)$$

$u = \sec x$	$v = 1 + \tan x$
$u' = \sec x \tan x$	$v' = \sec^2 x$
$d\left(\frac{u}{v}\right) = \frac{v u' - u v'}{v^2}$	

(27) Find the 25th derivative of $\cos x$.

Sol: Given $f(x) = \cos x$.

$$f'(x) = -\sin x$$

$$f''(x) = -\cos x$$

$$f'''(x) = \sin x$$

$$f^{(4)}(x) = \cos x$$

⋮

$$f^{(24)}(x) = \cos x$$

$$f^{(25)}(x) = -\sin x$$

H.W Problem:

(1) Find the derivative of the following:

(i) $y = \frac{\cos x}{1 - \sin x}$

(ii) $y = \sin x \tan x$

(iii) $f(x) = x e^x \operatorname{cosec} x$

(2) Find $\frac{d^{99}}{dx^{99}}(\sin x)$.

Formulae:

(1) $\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}$

(2) $\frac{d}{dx}(\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}$

(3) $\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$

(4) $\frac{d}{dx}(\cot^{-1} x) = \frac{-1}{1+x^2}$

(5) $\frac{d}{dx}(\operatorname{cosec}^{-1} x) = \frac{-1}{x\sqrt{x^2-1}}$

(6) $\frac{d}{dx}(\sec^{-1} x) = \frac{1}{x\sqrt{x^2-1}}$

(7) $\frac{d}{dx}(\log x) = \frac{1}{x}$

(8) $\frac{d}{dx}(\sinh x) = \cosh x$

(9) $\frac{d}{dx}(\cosh x) = \sinh x$

(10) $\frac{d}{dx}(\tanh x) = \operatorname{sech}^2 x$

(11) $\frac{d}{dx}(\coth x) = -\operatorname{cosech}^2 x$

(12) $\frac{d}{dx}(\operatorname{cosech} x) = -\operatorname{cosech} x \coth x$

(13) $\frac{d}{dx}(\operatorname{sech} x) = -\operatorname{sech} x \tanh x$

(14) $\frac{d}{dx}(\sinh^{-1} x) = \frac{1}{\sqrt{1+x^2}}$

(15) $\frac{d}{dx}(\cosh^{-1} x) = \frac{1}{\sqrt{x^2-1}}$

(16) $\frac{d}{dx}(\tanh^{-1} x) = \frac{1}{1-x^2}$

(17) $\frac{d}{dx}(\coth^{-1} x) = \frac{1}{1-x^2}$

(18) $\frac{d}{dx}(\operatorname{cosech}^{-1} x) = \frac{-1}{x\sqrt{x^2+1}}$

(19) $\frac{d}{dx}(\operatorname{sech}^{-1}x) = \frac{-1}{x\sqrt{1-x^2}}$

(20) $\sinh x = \frac{e^x - e^{-x}}{2}$

(21) $\cosh x = \frac{e^x + e^{-x}}{2}$

(22) $\operatorname{cosech} x = \frac{1}{\sinh x}$

(23) $\operatorname{sech} x = \frac{1}{\cosh x}$

(24) $\tanh x = \frac{\sinh x}{\cosh x}$

(25) $\coth x = \frac{1}{\tanh x}$

(26) $\sinh(-x) = -\sinh x$

(27) $\cosh(-x) = \cosh x$

(28) $\cosh^2 x - \sinh^2 x = 1$

(29) $1 - \tanh^2 x = \operatorname{sech}^2 x$

(28) Find y' if $y = \sqrt{\cos \sqrt{x}}$.

Sol: Given $y = \sqrt{\cos \sqrt{x}} = (\cos \sqrt{x})^{1/2}$

$y' = \frac{1}{2} (\cos \sqrt{x})^{1/2-1} (-\sin \sqrt{x}) \left(\frac{1}{2} x^{1/2-1}\right)$

$= \frac{1}{2} (\cos \sqrt{x})^{-1/2} (-\sin \sqrt{x}) \left(\frac{1}{2} x^{-1/2}\right) = \frac{-\sin \sqrt{x}}{4 \sqrt{\cos \sqrt{x}} \sqrt{x}}$

H.w

(1) Find y' if (i) $y = \sin^5 x$ (ii) $y = \cos(x^2)$ (iii) $y = e^{\sqrt{x}}$ (iv) $y = \sin(\sin(\sin x))$

Av

(29) Find y'' if $x^4 + y^4 = 16$.

Sol: Given $x^4 + y^4 = 16$ — (1)

Differentiating (1), with respect to x , we get

$4x^3 + 4y^3 \cdot y' = 0 \Rightarrow x^3 + y^3 \cdot y' = 0$ — (2) $\Rightarrow y^3 y' = -x^3 \Rightarrow y' = \frac{-x^3}{y^3}$ — (3)

Differentiating (2) with respect to x , we get

$3x^2 + y^3 \cdot y'' + y' \cdot 3y^2 \cdot y' = 0$

$3x^2 + y^3 \cdot y'' + 3y^2 \left(\frac{-x^3}{y^3}\right)^2 = 0$

$3x^2 + y^3 \cdot y'' + 3y^2 \left(\frac{x^6}{y^6}\right) = 0 \Rightarrow 3x^2 + y^3 \cdot y'' + \frac{3x^6}{y^4} = 0$

$\Rightarrow y^3 y'' = -3x^2 - \frac{3x^6}{y^4} = -3x^2 \left(1 + \frac{x^4}{y^4}\right) = -3x^2 \left(\frac{y^4 + x^4}{y^4}\right) = -3x^2 \left(\frac{16}{y^4}\right)$
(\because by (1))

$u = y^3, v = y'$
 $u' = 3y^2 \cdot y', v' = y''$
 $d(uv) = uv' + vu'$

$$y'' = -\frac{48x^2}{y^7}$$

(A0)

30 Find y' for $\cos(xy) = 1 + \sin y$.

Sol: Given $\cos(xy) = 1 + \sin y$. — ①

Diff. ① w.r.t. x , we get

$$-\sin(xy) \cdot (xy' + y \cdot 1) = \cos y \cdot y'$$

$$-xy' \sin(xy) - y \sin(xy) = \cos y \cdot y'$$

$$-y \sin(xy) = y' \cos y + xy' \sin(xy) = y' (\cos y + x \sin(xy))$$

$$\therefore y' = \frac{-y \sin(xy)}{\cos y + x \sin(xy)}$$

$$\begin{aligned} u &= x, v = y \\ u' &= 1, v' = y' \\ d(uv) &= uv' + vu' \end{aligned}$$

(A0)

31 Find the derivative of $f(x) = \cos^{-1} \left(\frac{b + a \cos x}{a + b \cos x} \right)$.

Sol: Given $f(x) = \cos^{-1} \left(\frac{b + a \cos x}{a + b \cos x} \right)$

$$\frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}$$

$$f'(x) = \frac{-1}{\sqrt{1 - \left(\frac{b + a \cos x}{a + b \cos x} \right)^2}} \left[\frac{(a + b \cos x)(-a \sin x) - (b + a \cos x)(-b \sin x)}{(a + b \cos x)^2} \right]$$

$$\begin{aligned} u &= b + a \cos x, v = a + b \cos x \\ u' &= -a \sin x, v' = -b \sin x \\ d\left(\frac{u}{v}\right) &= \frac{vu' - uv'}{v^2} \end{aligned}$$

$$f'(x) = \frac{-(a + b \cos x)}{\sqrt{(a + b \cos x)^2 - (b + a \cos x)^2}} \left[\frac{-a^2 \sin x - ab \sin x \cos x + b^2 \sin x + ab \sin x \cos x}{(a + b \cos x)^2} \right]$$

$$= \frac{-1}{\sqrt{a^2 + b^2 \cos^2 x + 2ab \cos x - b^2 - a^2 \cos^2 x - 2ab \cos x}} \left(\frac{\sin x (b^2 - a^2)}{a + b \cos x} \right)$$

$$= \frac{(a^2 - b^2) \sin x}{(a + b \cos x) \sqrt{(a^2 - b^2) - \cos^2 x (a^2 - b^2)}} = \frac{(a^2 - b^2) \sin x}{(a + b \cos x) \sqrt{(a^2 - b^2)(1 - \cos^2 x)}}$$

$$= \frac{(a^2 - b^2) \sin x}{(a + b \cos x) \sqrt{(a^2 - b^2) \sin^2 x}} \quad (\because \sin^2 x + \cos^2 x = 1)$$

$$= \frac{(a^2 - b^2) \sin x}{(a + b \cos x) \sin x \sqrt{a^2 - b^2}} = \frac{\sqrt{a^2 - b^2}}{a + b \cos x}$$

32) Find the derivative of $f(x) = \tanh^{-1} \left[\tan \frac{x}{2} \right]$.

Sol: Given $f(x) = \tanh^{-1} \left[\tan \frac{x}{2} \right]$.

$$\frac{d}{dx} (\tanh^{-1} x) = \frac{1}{1 - x^2}$$

$$\frac{d}{dx} (\tan x) = \sec^2 x$$

$$f'(x) = \frac{1}{1 - \left(\tan \frac{x}{2}\right)^2} \left(\sec^2 \frac{x}{2}\right) \left(\frac{1}{2}\right)$$

$$= \frac{1}{1 - \tan^2 \frac{x}{2}} \left(\sec^2 \frac{x}{2}\right) \left(\frac{1}{2}\right) = \frac{1}{1 - \frac{\sin^2 x/2}{\cos^2 x/2}} \left(\sec^2 x/2\right) \left(\frac{1}{2}\right)$$

$$= \frac{\cos^2 x/2}{\cos^2 x/2 - \sin^2 x/2} \left(\frac{1}{2} \sec^2 x/2\right) = \frac{\cos^2 x/2}{\cos^2 x/2 - \sin^2 x/2} \left(\frac{1}{2 \cos^2 x/2}\right)$$

$$= \frac{1}{2(\cos^2 x/2 - \sin^2 x/2)}$$

33) Find the tangent line to the equation $x^3 + y^3 = 6xy$ at the point (3, 3) & at what point the tangent line horizontal in the first quadrant.

Sol: Given $x^3 + y^3 = 6xy$ — ①

Diff. ① w.r.t. x, we get

$$3x^2 + 3y^2 \cdot y' = 6(xy' + y \cdot 1)$$

$$\Rightarrow 3x^2 + 3y^2 y' = 6xy' + 6y \Rightarrow 3y^2 y' - 6xy' = 6y - 3x^2$$

$$\Rightarrow y'(3y^2 - 6x) = 6y - 3x^2 \Rightarrow y' = \frac{6y - 3x^2}{3y^2 - 6x}$$

$$(y')_{(3,3)} = \frac{6(3) - 3(3)^2}{3(3)^2 - 6(3)} = \frac{18 - 27}{27 - 18} = \frac{-9}{9} = -1 = m \text{ (Slope)}$$

Equation of tangent line is $y - y_1 = m(x - x_1)$

$$y - 3 = -1(x - 3) \Rightarrow y - 3 = -x + 3$$

$$\Rightarrow x + y = 3 + 3 = 6 \Rightarrow x + y = 6$$

$$u = x, \quad v = y$$

$$u' = 1, \quad v' = y'$$

$$d(uv) = uv' + vu'$$

$$m = -1$$

$$x_1 = 3$$

$$y_1 = 3$$

The tangent line is horizontal if $y' = 0$

$$(ii) y' = \frac{6y - 3x^2}{3y^2 - 6x} = \frac{2y - x^2}{y^2 - 2x} = 0$$

$$\Rightarrow 2y - x^2 = 0 \Rightarrow 2y = x^2 \Rightarrow y = \frac{x^2}{2} \quad \text{--- (2)}$$

Substituting (2) in (1),

$$x^3 + \left(\frac{x^2}{2}\right)^3 = 6x \left(\frac{x^2}{2}\right) \Rightarrow x^3 + \frac{x^6}{8} = \frac{6x^3}{2} = 3x^3$$

$$\Rightarrow \frac{x^6}{8} = 3x^3 - x^3 = 2x^3 \Rightarrow \frac{x^3}{8} = 2 \Rightarrow x^3 = 16 = 2^4$$

$$\Rightarrow \boxed{x = 2^{4/3}} \quad \text{--- (3)}$$

$$\text{Subst. (3) in (2), } y = \frac{(2^{4/3})^2}{2} = \frac{2^{8/3}}{2} = 2^{8/3} \cdot 2^{-1} = 2^{8/3-1} = 2^{5/3}$$

Hence the tangent line is horizontal at $(2^{4/3}, 2^{5/3})$.

(34) Find y' if $y = (\sin x)^{(\sin x)^{(\sin x)^{\dots}}}$

Sol. Given $y = (\sin x)^{(\sin x)^{(\sin x)^{\dots}}}$

$$\Rightarrow y = (\sin x)^y \quad \text{--- (*)}$$

$$\log y = \log(\sin x)^y = y \log(\sin x) \quad \text{--- (1)}$$

Diff. (1) w.r.t. x , we get

$$\frac{1}{y} y' = y \frac{1}{\sin x} \cos x + \log(\sin x) \cdot y'$$

$$\frac{1}{y} y' - \log(\sin x) \cdot y' = \frac{y}{\sin x} \cos x = y \cot x$$

$$y' \left(\frac{1}{y} - \log(\sin x) \right) = y \cot x \Rightarrow y' \left(\frac{1 - y \log(\sin x)}{y} \right) = y \cot x$$

$$\Rightarrow y' = \frac{y^2 \cot x}{1 - y \log(\sin x)} = \frac{y^2 \cot x}{1 - \log(\sin x)^y} = \frac{y^2 \cot x}{1 - \log y} \quad (\because \text{by } *)$$

(35) Find an equation of the normal line to the curve $y = \sqrt{x}$ at the point $(1, 1)$.

Sol. Given $y = \sqrt{x}$, $(1, 1)$.

$$y = x^{1/4}$$

$$\frac{dy}{dx} = m = \frac{1}{4} x^{1/4-1} = \frac{1}{4} x^{-3/4}$$

$$m_{(1,1)} = \frac{1}{4} (1)^{-3/4} = \frac{1}{4} \times 1 = \frac{1}{4}$$

Equation of the normal line is

$$y - y_1 = -\frac{1}{m} (x - x_1)$$

$$y - 1 = -\frac{1}{1/4} (x - 1) \Rightarrow y - 1 = -4(x - 1)$$

$$\Rightarrow y - 1 = -4x + 4 \Rightarrow 4x + y = 4 + 1 = 5 \Rightarrow 4x + y = 5$$

$$\begin{aligned} m &= \frac{1}{4} \\ x_1 &= 1 \\ y_1 &= 1 \end{aligned}$$

- (H.W)
- ① If $x^3 + y^3 = 16$ find the value of $\frac{d^2y}{dx^2}$ at $(2, 2)$
 - ② Find $\frac{dy}{dx}$ if $y = x \sin^{-1} x + \sqrt{1-x^2}$.
 - ③ If $e^x \cos x = 1 + \sin(xy)$, then find $\frac{dy}{dx}$.
 - ④ Find an equation of the tangent line to the curve $y \sin(2x) = x \cos(2y)$ at the point $(\frac{\pi}{2}, \frac{\pi}{4})$.

(AU) 36 Find the critical points of $y = 5x^3 - 6x$.

Sol: Given $y = 5x^3 - 6x$.

Critical points: $y' = 0$ (i) $\frac{dy}{dx} = 0$

$$y' = 15x^2 - 6 = 0 \Rightarrow 15x^2 = 6 \Rightarrow x^2 = \frac{6}{15} = \frac{2}{5}$$

$$\therefore x = \pm \sqrt{\frac{2}{5}}$$

Definition: (Critical number)

A critical number of a function f is a number c in the domain of f such that either $f'(c) = 0$ or $f'(c)$ does not exist.

37 Find the critical points of $f(x) = x^{3/5}(4-x)$.

Sol: Given $f(x) = x^{3/5}(4-x) = 4x^{3/5} - x x^{3/5} = 4x^{3/5} - x^{8/5}$

Critical points: $f'(x) = 0$

$$f'(x) = 4\left(\frac{3}{5}\right)x^{\frac{3}{5}-1} - \frac{8}{5}x^{\frac{8}{5}-1} = 0$$

$$\Rightarrow \frac{12}{5}x^{-2/5} - \frac{8}{5}x^{3/5} = 0$$

$$\Rightarrow \frac{12}{5}x^{-2/5} = \frac{8}{5}x^{3/5} \Rightarrow \frac{12}{5} \times \frac{5}{8} = \frac{x^{3/5}}{x^{-2/5}} = x^{3/5} x^{2/5}$$

$$\Rightarrow \boxed{\frac{3}{2} = x}$$

$f'(x)$ doesn't exist when $x=0$.

Hence the critical points are 0 & $\frac{3}{2}$.

First derivative test:

Suppose that c is a critical number of a continuous function f .

- (i) If f' changes from $+$ to $-$ at c , then f has a local maximum at c .
- (ii) If f' changes from $-$ to $+$ at c , then f has a local minimum at c .
- (iii) If f' does not change sign at c , then f has no local maximum or minimum at c .

Second derivative test:

Suppose f'' is continuous near c .

- (i) If $f'(c) = 0$ & $f''(c) > 0$, then f has a local minimum at c .
- (ii) If $f'(c) = 0$ & $f''(c) < 0$, then f has a local maximum at c .

(10) (38) If $f(x) = 2x^3 + 3x^2 - 36x$, find the intervals on which it is increasing or decreasing, the local maximum & local minimum values of $f(x)$. Also find the intervals of concavity & the inflection points.

Sol: Given $f(x) = 2x^3 + 3x^2 - 36x$
 $\Rightarrow f'(x) = 6x^2 + 6x - 36$

x	$+$
-6	1
$+3$	-2
$x+3$	$x-2$

Critical points: $f'(x) = 0$

$$f'(x) = 6x^2 + 6x - 36 = 0 \Rightarrow x^2 + x - 6 = 0$$

$$\Rightarrow (x+3)(x-2) = 0$$

$$\Rightarrow x = -3, 2$$

Critical points are -3 & 2 .



Interval	Sign of f'	Behaviour of f
$-\infty < x < -3$	+	increasing
$-3 < x < 2$	-	decreasing
$2 < x < \infty$	+	increasing

} local maximum
 } local minimum

At $x = -3$, we get local maximum & at $x = 2$, we get local minimum values.

$$f(-3) = 2(-3)^3 + 3(-3)^2 - 36(-3) = 81$$


$$f(2) = 2(2)^3 + 3(2)^2 - 36(2) = -44$$

Hence the local maximum value is 81 & the local minimum value is -44.

$$f''(x) = 12x + 6$$

$$f''(x) = 0 \Rightarrow 12x + 6 = 0 \Rightarrow 12x = -6 \Rightarrow x = \frac{-6}{12} = \frac{-1}{2}$$

$$\therefore x = \frac{-1}{2}$$



Interval	Sign of f''	Behaviour of f
$-\infty < x < -\frac{1}{2}$	-	Concave down
$-\frac{1}{2} < x < \infty$	+	Concave up

Inflection points:

$$f(-\frac{1}{2}) = 2(-\frac{1}{2})^3 + 3(-\frac{1}{2})^2 - 36(-\frac{1}{2}) = \frac{37}{2}$$

Hence the inflection point is $(-\frac{1}{2}, \frac{37}{2})$.

10 (39) For the function $f(x) = 2 + 2x^2 - x^4$, find the intervals of increase or decrease, local maximum & minimum values, the intervals of concavity & the inflection points.

Sol: Given $f(x) = 2 + 2x^2 - x^4$

$$f'(x) = 4x - 4x^3$$

Critical points: $f'(x) = 0$

$$f'(x) = 0 \Rightarrow 4x - 4x^3 = 0 \Rightarrow 4x(1 - x^2) = 0 \Rightarrow x = 0, 1 - x^2 = 0$$

$\Rightarrow x=0, x^2=1 \Rightarrow x=\sqrt{1}=\pm 1$

Hence the critical points are $-1, 0$ & 1 .



Interval	Sign of f'	Behaviour of f
$-\infty < x < -1$	+	increasing
$-1 < x < 0$	-	decreasing
$0 < x < 1$	+	increasing
$1 < x < \infty$	-	decreasing

} local maximum
 } local minimum
 } local maximum

At $x = \pm 1$, we get local maximum value.

$f(1) = 2 + 2(1)^2 - (1)^4 = 2 + 2 - 1 = 3$

\therefore Local maximum value is 3.

At $x = 0$, we get local minimum value.

$f(0) = 2 + 2(0)^2 - (0)^4 = 2$

\therefore Local minimum value is 2.

$f''(x) = 4 - 12x^2$

$f''(x) = 0 \Rightarrow 4 - 12x^2 = 0 \Rightarrow 12x^2 = 4 \Rightarrow x^2 = \frac{4}{12} = \frac{1}{3} \Rightarrow x = \pm \sqrt{\frac{1}{3}}$

$\therefore \boxed{x = \pm \sqrt{\frac{1}{3}}} \Rightarrow \boxed{x = \pm \frac{1}{\sqrt{3}}}$



Interval	Sign of f''	Behaviour of f
$-\infty < x < -\frac{1}{\sqrt{3}}$	-	Concave down
$-\frac{1}{\sqrt{3}} < x < \frac{1}{\sqrt{3}}$	+	Concave up
$\frac{1}{\sqrt{3}} < x < \infty$	-	Concave down

$\boxed{-\frac{1}{\sqrt{3}} = -0.6}$
 $\boxed{\frac{1}{\sqrt{3}} = 0.6}$

Inflection points:

$f\left(\frac{1}{\sqrt{3}}\right) = 2 + 2\left(\frac{1}{\sqrt{3}}\right)^2 - \left(\frac{1}{\sqrt{3}}\right)^4 = 2 + \frac{2}{3} - \frac{1}{9} = \frac{23}{9}$

$f\left(-\frac{1}{\sqrt{3}}\right) = 2 + 2\left(-\frac{1}{\sqrt{3}}\right)^2 - \left(-\frac{1}{\sqrt{3}}\right)^4 = 2 + \frac{2}{3} - \frac{1}{9} = \frac{23}{9}$

Hence the inflection points are $\left(-\frac{1}{\sqrt{3}}, \frac{23}{9}\right)$ & $\left(\frac{1}{\sqrt{3}}, \frac{23}{9}\right)$.

Q10
40

Find the local maximum & minimum values of $f(x) = \sqrt{x} - \sqrt[4]{x}$ using both the first & second derivative tests.

Sol: Given $f(x) = \sqrt{x} - \sqrt[4]{x} = x^{1/2} - x^{1/4}$

$$f'(x) = \frac{1}{2}x^{1/2-1} - \frac{1}{4}x^{1/4-1} = \frac{1}{2}x^{-1/2} - \frac{1}{4}x^{-3/4}$$

Critical points:

$$f'(x) = 0 \Rightarrow \frac{1}{2}x^{-1/2} - \frac{1}{4}x^{-3/4} = 0 \Rightarrow \frac{1}{2}x^{-1/2} = \frac{1}{4}x^{-3/4}$$

$$\Rightarrow \frac{4}{2} = \frac{x^{-3/4}}{x^{-1/2}} \Rightarrow 2 = x^{-3/4} \cdot x^{1/2} = x^{-3/4+1/2} = x^{-3+2 \over 4} = x^{-1/4}$$

$$\Rightarrow 2 = x^{-1/4} \Rightarrow \frac{2}{x^{-1/4}} = 1 \Rightarrow 2x^{1/4} = 1 \Rightarrow x^{1/4} = 1/2$$

$$\Rightarrow x = (1/2)^4 = 1/16$$

At $x=0$, $f'(x)$ doesn't exist.

Hence the critical points are 0 & $1/16$.

$\frac{1}{16} = 0.06$

First derivative test:

Interval	Sign of f'	Behaviour of f
$-\infty < x < 0$	(not defined)	(not defined)
$0 < x < 1/16$	-	decreasing
$1/16 < x < \infty$	+	increasing

} local minimum

At $x = 1/16$, we get local minimum value.

$$f(1/16) = \sqrt{1/16} - \sqrt[4]{1/16} = 1/4 - 1/2 = -1/4$$

Hence the local minimum value is $-1/4$.

Second derivative test:

$$f''(x) = \frac{1}{2}(-1/2)x^{-1/2-1} - 1/4(-3/4)x^{-3/4-1} = -1/4x^{-3/2} + 3/16x^{-7/4}$$

$$\therefore f''(1/16) = -16 + 24 = 8 > 0 \Rightarrow \text{local minimum at } x = 1/16$$

$$\therefore f(1/16) = -1/4$$

Hence the local minimum value is $-1/4$.

A.w

(27)

① For the function $f(x) = x^3 - 3x^2 + 1$, find the intervals of increase or decrease, local maximum & minimum values, the intervals of concavity & the inflection points.

② For the function $f(x) = 3x^4 - 4x^3 - 12x^2 + 5$, find the intervals of increase or decrease, local maximum & minimum values, the intervals of concavity & the inflection points.

FUNCTIONS OF SEVERAL VARIABLES

① Evaluate: $\lim_{\substack{x \rightarrow \infty \\ y \rightarrow 2}} \frac{xy+5}{x^2+2y^2}$

Sol: $\lim_{\substack{x \rightarrow \infty \\ y \rightarrow 2}} \frac{xy+5}{x^2+2y^2} = \lim_{x \rightarrow \infty} \left[\lim_{y \rightarrow 2} \frac{xy+5}{x^2+2y^2} \right]$

$$= \lim_{x \rightarrow \infty} \left[\frac{x(2)+5}{x^2+2(2)^2} \right] = \lim_{x \rightarrow \infty} \left[\frac{2x+5}{x^2+8} \right]$$

$$= \lim_{x \rightarrow \infty} \left[\frac{x(2+5/x)}{x^2(1+8/x^2)} \right] = \lim_{x \rightarrow \infty} \left[\frac{2+5/x}{x(1+8/x^2)} \right]$$

$$= \frac{2+5/\infty}{\infty(1+8/\infty)} = \frac{2+0}{\infty(1+0)} = \frac{2}{\infty} = 0$$

H.W. ① Evaluate: $\lim_{\substack{x \rightarrow 1 \\ y \rightarrow 2}} \frac{3x^2y}{x^2+y^2+5}$

② If $f(x,y) = \log \sqrt{x^2+y^2}$, show that $\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = 0$.

Sol: Given $f(x,y) = \log \sqrt{x^2+y^2} = \log(x^2+y^2)^{1/2} = \frac{1}{2} \log(x^2+y^2)$

$\Rightarrow f(x,y) = \frac{1}{2} \log(x^2+y^2)$

$\frac{\partial f}{\partial x} = \frac{1}{2} \cdot \frac{1}{x^2+y^2} \cdot 2x = \frac{x}{x^2+y^2}$

$\frac{\partial^2 f}{\partial x^2} = \frac{(x^2+y^2) \cdot 1 - x(2x)}{(x^2+y^2)^2} = \frac{x^2+y^2-2x^2}{(x^2+y^2)^2}$

$\frac{\partial^2 f}{\partial x^2} = \frac{y^2-x^2}{(x^2+y^2)^2}$ — ①

$\frac{\partial f}{\partial y} = \frac{1}{2} \cdot \frac{1}{x^2+y^2} \cdot 2y = \frac{y}{x^2+y^2}$

$\frac{\partial^2 f}{\partial y^2} = \frac{(x^2+y^2) \cdot 1 - y(2y)}{(x^2+y^2)^2} = \frac{x^2+y^2-2y^2}{(x^2+y^2)^2}$

$\frac{\partial^2 f}{\partial y^2} = \frac{x^2-y^2}{(x^2+y^2)^2}$ — ②

① + ② $\Rightarrow \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = \frac{y^2-x^2}{(x^2+y^2)^2} + \frac{x^2-y^2}{(x^2+y^2)^2} = \frac{y^2-x^2+x^2-y^2}{(x^2+y^2)^2} = 0$

w.r.t. x

$u = x$	$v = x^2+y^2$
$u' = 1$	$v' = 2x$
$d\left(\frac{u}{v}\right) = \frac{v u' - u v'}{v^2}$	

w.r.t. y

$u = y$	$v = x^2+y^2$
$u' = 1$	$v' = 2y$

③ If $x = r \cos \theta$, $y = r \sin \theta$, find (i) $\frac{\partial x}{\partial r}$ (ii) $\frac{\partial y}{\partial \theta}$ (iii) $\frac{\partial r}{\partial x}$ (iv) $\frac{\partial \theta}{\partial y}$

Sol: Given $x = r \cos \theta$, $y = r \sin \theta$

(i) $\frac{\partial x}{\partial r} = \cos \theta$

(ii) $\frac{\partial y}{\partial \theta} = r \cos \theta$

We know that $r = \sqrt{x^2 + y^2}$ & $\theta = \tan^{-1}\left(\frac{y}{x}\right)$

$\Rightarrow r = (x^2 + y^2)^{1/2}$ & $\theta = \tan^{-1}\left(\frac{y}{x}\right)$

(iii) $\frac{\partial r}{\partial x} = \frac{1}{2}(x^2 + y^2)^{1/2 - 1} \cdot 2x = x(x^2 + y^2)^{-1/2} = \frac{x}{(x^2 + y^2)^{1/2}} = \frac{x}{\sqrt{x^2 + y^2}}$

(iv) $\frac{\partial \theta}{\partial y} = \frac{1}{1 + \left(\frac{y}{x}\right)^2} \cdot \frac{1}{x} = \frac{1}{\frac{x^2 + y^2}{x^2}} \cdot \frac{1}{x} = \frac{x^2}{x^2 + y^2} \cdot \frac{1}{x} = \frac{x}{x^2 + y^2}$

$\frac{d}{dx}(\tan^{-1}x) = \frac{1}{1+x^2}$

④ Find $\frac{du}{dt}$ in terms of t , if $u = x^3 + y^3$ where $x = at^2$, $y = 2at$.

Sol: Given $u = x^3 + y^3$, $x = at^2$, $y = 2at$

$\therefore u = (at^2)^3 + (2at)^3 = a^3t^6 + 8a^3t^3$

$\frac{du}{dt} = a^3 \cdot 6t^5 + 8a^3 \cdot 3t^2 = 6a^3t^5 + 24a^3t^2 = 6a^3(t^5 + 4t^2) = 6a^3t^2(t^3 + 4)$

Euler's theorem on homogeneous function:

If u is a homogeneous function of degree n in x & y , then

$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = nu$

① If $u = (x-y)(y-z)(z-x)$, then show that $\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} = 0$.

⑤ If $u = \tan^{-1}\left(\frac{x^3 + y^3}{x - y}\right)$, then prove that $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = \sin 2u$.

Sol: Given $u = \tan^{-1}\left(\frac{x^3 + y^3}{x - y}\right)$

$\Rightarrow \tan u = \frac{x^3 + y^3}{x - y} = f(x, y)$

$$f(tx, ty) = \frac{(tx)^3 + (ty)^3}{tx - ty} = \frac{t^3x^3 + t^3y^3}{t(x-y)} = \frac{t^3(x^3 + y^3)}{t(x-y)} = t^2 \left(\frac{x^3 + y^3}{x-y} \right) = t^2 f(x, y)$$

∴ f is a homogeneous function of degree 2 in x & y.

∴ By Euler's theorem, we get $x \frac{\partial f}{\partial x} + y \frac{\partial f}{\partial y} = 2f$ — ①

Here $f = \tan u$

$$\frac{\partial f}{\partial x} = \sec^2 u \frac{\partial u}{\partial x}$$

$$\frac{\partial f}{\partial y} = \sec^2 u \frac{\partial u}{\partial y}$$

$$\therefore \text{①} \Rightarrow x \sec^2 u \frac{\partial u}{\partial x} + y \sec^2 u \frac{\partial u}{\partial y} = 2 \tan u$$

$$\sec^2 u \left(x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} \right) = 2 \tan u$$

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = 2 \tan u \times \frac{1}{\sec^2 u} = 2 \frac{\sin u}{\cos u} \times \cos^2 u = 2 \sin u \cos u = \sin 2u$$

$2 \sin A \cos A = \sin 2A$

$$\therefore x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = \sin 2u$$

H.w
① If $u = \sin^{-1} \left(\frac{x^3 - y^3}{x + y} \right)$, then prove that $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = 2 \tan u$.

② If $u = \cos^{-1} \left(\frac{x+y}{\sqrt{x+y}} \right)$, then prove that $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = -\frac{1}{2} \cot u$

10
⑥ Verify the Euler's theorem for the function $u = x^2 + y^2 + 2xy$.

Sol: Given $u = x^2 + y^2 + 2xy = u(x, y)$

Euler's theorem: $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = nu$

LHS $\frac{\partial u}{\partial x} = 2x + 2y$, $\frac{\partial u}{\partial y} = 2y + 2x$

$$\therefore x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = x(2x + 2y) + y(2y + 2x) = 2x^2 + 2xy + 2y^2 + 2xy = 2x^2 + 2y^2 + 4xy$$

$$\therefore x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = 2x^2 + 2y^2 + 4xy \text{ — ①}$$

RHS $u(tx, ty) = (tx)^2 + (ty)^2 + 2(tx)(ty)$
 $= t^2x^2 + t^2y^2 + 2t^2xy = t^2(x^2 + y^2 + 2xy) = t^2 u(x, y)$

∴ u is a homogeneous function of degree 2 in x & y.

$$\therefore n = 2$$

$$nu = 2(x^2 + y^2 + 2xy) = 2x^2 + 2y^2 + 4xy \quad \text{--- (2)}$$

$$\text{From (1) \& (2), } x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = nu$$

Definition:

A function $f(x, y)$ is said to be a homogeneous function of degree n in x & y , if $f(tx, ty) = t^n f(x, y)$ for any positive t .

$$\textcircled{7} \text{ If } u = \sin^{-1} \left(\frac{x+2y+3z}{\sqrt{x^8+y^8+z^8}} \right), \text{ then show that } x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z} + 3 \tan u = 0.$$

$$\text{Sol: Given } u = \sin^{-1} \left(\frac{x+2y+3z}{\sqrt{x^8+y^8+z^8}} \right) \Rightarrow \sin u = \frac{x+2y+3z}{\sqrt{x^8+y^8+z^8}} = f(x, y, z)$$

$$f(tx, ty, tz) = \frac{tx+2ty+3tz}{\sqrt{(tx)^8+(ty)^8+(tz)^8}} = \frac{t(x+2y+3z)}{t^4 \sqrt{x^8+y^8+z^8}} = t^{-3} \left(\frac{x+2y+3z}{\sqrt{x^8+y^8+z^8}} \right) = t^{-3} f(x, y, z)$$

$\therefore f$ is a homogeneous function of degree (-3) in x, y & z .

\therefore By Euler's theorem, we get

$$x \frac{\partial f}{\partial x} + y \frac{\partial f}{\partial y} + z \frac{\partial f}{\partial z} = n f = -3 f \quad (\because n = -3) \quad \text{--- (1)}$$

$$\text{Here } f = \sin u$$

$$\frac{\partial f}{\partial x} = \cos u \frac{\partial u}{\partial x}$$

$$\frac{\partial f}{\partial y} = \cos u \frac{\partial u}{\partial y}$$

$$\frac{\partial f}{\partial z} = \cos u \frac{\partial u}{\partial z} \quad \text{--- (2)}$$

Subst. (2) in (1),

$$x \cos u \frac{\partial u}{\partial x} + y \cos u \frac{\partial u}{\partial y} + z \cos u \frac{\partial u}{\partial z} = -3 \sin u$$

$$\Rightarrow x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z} = \frac{-3 \sin u}{\cos u} = -3 \tan u$$

$$\Rightarrow x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z} + 3 \tan u = 0$$

8) If $u = f\left(\frac{x}{y}, \frac{y}{z}, \frac{z}{x}\right)$, then prove that $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z} = 0$.

Sol: Given $u(x, y, z) = f\left(\frac{x}{y}, \frac{y}{z}, \frac{z}{x}\right)$

$$u(tx, ty, tz) = f\left(\frac{tx}{ty}, \frac{ty}{tz}, \frac{tz}{tx}\right) = f\left(\frac{x}{y}, \frac{y}{z}, \frac{z}{x}\right) = t^0 u(x, y, z)$$

$\therefore u$ is a homogeneous function of degree 0 in x, y & z .

\therefore By Euler's theorem, we get

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z} = nu = 0 \cdot u = 0 \quad (\because n=0)$$

9) If $u = \sin^{-1} \frac{x+y}{\sqrt{x}+\sqrt{y}}$, then prove that (i) $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = \frac{1}{2} \tan u$

(ii) $x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = -\frac{\sin u \cos 2u}{4 \cos^3 u}$

Sol: Given $u = \sin^{-1} \frac{x+y}{\sqrt{x}+\sqrt{y}} \Rightarrow \sin u = \frac{x+y}{\sqrt{x}+\sqrt{y}} = f(x, y)$

$$f(tx, ty) = \frac{tx+ty}{\sqrt{tx}+\sqrt{ty}} = \frac{t(x+y)}{\sqrt{t}(\sqrt{x}+\sqrt{y})} = \sqrt{t} \left(\frac{x+y}{\sqrt{x}+\sqrt{y}} \right) = t^{1/2} f(x, y)$$

$\therefore f$ is a homogeneous function of degree $1/2$ in x & y .

\therefore By Euler's theorem, we get

(i) $x \frac{\partial f}{\partial x} + y \frac{\partial f}{\partial y} = nf = \frac{1}{2} f \quad \text{--- (1)}$

Here $f = \sin u$

$$\frac{\partial f}{\partial x} = \cos u \frac{\partial u}{\partial x}, \quad \frac{\partial f}{\partial y} = \cos u \frac{\partial u}{\partial y} \quad \text{--- (2)}$$

Subst. (2) in (1), $x \cos u \frac{\partial u}{\partial x} + y \cos u \frac{\partial u}{\partial y} = \frac{1}{2} \sin u$

$$\Rightarrow x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = \frac{1}{2} \frac{\sin u}{\cos u} = \frac{1}{2} \tan u \quad \text{--- (3)}$$

(ii) Diff. (3) partially w.r.t. x , we get

$$x \frac{\partial^2 u}{\partial x^2} + \frac{\partial u}{\partial x} \cdot 1 + y \frac{\partial^2 u}{\partial x \partial y} = \frac{1}{2} \sec^2 u \frac{\partial u}{\partial x}$$

$$x \frac{\partial^2 u}{\partial x^2} + y \frac{\partial^2 u}{\partial x \partial y} = \frac{1}{2} \sec^2 u \frac{\partial u}{\partial x} - \frac{\partial u}{\partial x} = \frac{\partial u}{\partial x} \left(\frac{1}{2} \sec^2 u - 1 \right) \quad \text{--- (4)}$$

Diff. (3) partially w.r.t. y , we get

$$x \frac{\partial^2 u}{\partial x \partial y} + y \frac{\partial^2 u}{\partial y^2} + \frac{\partial u}{\partial y} \cdot 1 = \frac{1}{2} \sec^2 u \frac{\partial u}{\partial y}$$

$$x \frac{\partial^2 u}{\partial x \partial y} + y \frac{\partial^2 u}{\partial y^2} = \frac{1}{2} \sec^2 u \frac{\partial u}{\partial y} - \frac{\partial u}{\partial y} = \frac{\partial u}{\partial y} \left(\frac{1}{2} \sec^2 u - 1 \right) \quad \text{--- (5)}$$

$$(4) \times x + (5) \times y \Rightarrow$$

$$x^2 \frac{\partial^2 u}{\partial x^2} + xy \frac{\partial^2 u}{\partial x \partial y} + xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = x \frac{\partial u}{\partial x} \left(\frac{1}{2} \sec^2 u - 1 \right) + y \frac{\partial u}{\partial y} \left(\frac{1}{2} \sec^2 u - 1 \right)$$

$$x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = \left(\frac{1}{2} \sec^2 u - 1 \right) \left(x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} \right)$$

$$= \left(\frac{1}{2} \sec^2 u - 1 \right) \frac{1}{2} \tan u \quad (\because \text{by (3)})$$

$$= \left(\frac{1}{2 \cos^2 u} - 1 \right) \frac{1}{2} \tan u = \left(\frac{1 - 2 \cos^2 u}{2 \cos^2 u} \right) \frac{1}{2} \frac{\sin u}{\cos u}$$

$$= - \left(\frac{2 \cos^2 u - 1}{2 \cos^2 u} \right) \frac{1}{2} \frac{\sin u}{\cos u}$$

$$= - \frac{\sin u \cos 2u}{4 \cos^3 u}$$

$\cos^2 u = \frac{1 + \cos 2u}{2}$ $2 \cos^2 u = 1 + \cos 2u$ $2 \cos^2 u - 1 = \cos 2u$
--

$$(10) \text{ If } u = (x-y) f\left(\frac{y}{x}\right), \text{ then find } x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2}.$$

Sol: Given $u = (x-y) f\left(\frac{y}{x}\right)$

$$u(tx, ty) = (tx - ty) f\left(\frac{ty}{tx}\right) = t(x-y) f\left(\frac{y}{x}\right) = t u(x, y)$$

$\therefore u$ is a homogeneous function of degree 1 in x & y .

\therefore By Euler's Theorem, we get

$$x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = n(n-1)u = 1(1-1)u = 1(0)u = 0 \quad (\because n=1)$$

(11) (1) If $u = \frac{x^2 + y^2}{\sqrt{x+y}}$, then prove that $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = \frac{3}{2} u$.

(2) If $u = \cos^{-1} \frac{x}{y} + \tan^{-1} \frac{y}{x}$, then prove that $x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = 0$.

(3) If $u = \sin^{-1} \frac{\sqrt{x+y}}{\sqrt{x-y}}$, then (i) $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = 0$

(ii) $x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = 0$.

Jacobians:

(11) If $x = r \cos \theta$ & $y = r \sin \theta$, then find $\frac{\partial(x,y)}{\partial(r,\theta)}$.

Sol: Given $x = r \cos \theta$, $y = r \sin \theta$

$$\frac{\partial x}{\partial r} = \cos \theta$$

$$\frac{\partial y}{\partial r} = \sin \theta$$

$$\frac{\partial x}{\partial \theta} = -r \sin \theta$$

$$\frac{\partial y}{\partial \theta} = r \cos \theta$$

$$\frac{\partial(x,y)}{\partial(r,\theta)} = \begin{vmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{vmatrix} = \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix} = \cos \theta (r \cos \theta) + r \sin \theta (\sin \theta)$$

$$= r(\cos^2 \theta + \sin^2 \theta) = r \quad (\because \cos^2 \theta + \sin^2 \theta = 1)$$

(12) If $x = uv$ & $y = \frac{u}{v}$ then find $\frac{\partial(x,y)}{\partial(u,v)}$.

Sol: Given $x = uv$, $y = \frac{u}{v} = uv^{-1}$

$$\frac{\partial x}{\partial u} = v$$

$$\frac{\partial y}{\partial u} = \frac{1}{v}$$

$$\frac{\partial x}{\partial v} = u$$

$$\frac{\partial y}{\partial v} = u(-1)v^{-1-1} = -uv^{-2} = -\frac{u}{v^2}$$

$$\frac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} v & u \\ \frac{1}{v} & -\frac{u}{v^2} \end{vmatrix} = v\left(-\frac{u}{v^2}\right) - u\left(\frac{1}{v}\right)$$

$$= -\frac{u}{v} - \frac{u}{v} = -\frac{2u}{v}$$

(13) If $x = u^2 - v^2$, $y = 2uv$ find the Jacobian of x, y with respect to u & v .
[Hint: $\frac{\partial(x,y)}{\partial(u,v)}$]

(13) State the properties of Jacobians.

Sol: ① If u & v are the functions of x & y , then

$$\frac{\partial(u,v)}{\partial(x,y)} \times \frac{\partial(x,y)}{\partial(u,v)} = 1.$$

② If u, v are functions of x, y & x, y are functions of r, s then

$$\frac{\partial(u,v)}{\partial(x,y)} \frac{\partial(x,y)}{\partial(r,s)} = \frac{\partial(u,v)}{\partial(r,s)}$$

③ If u, v, w are functionally dependent functions of three independent variables x, y, z then

$$\frac{\partial(u,v,w)}{\partial(x,y,z)} = 0.$$

14) If $u = 2xy$, $v = x^2 - y^2$ & $x = r \cos \theta$, $y = r \sin \theta$. Evaluate $\frac{\partial(u, v)}{\partial(r, \theta)}$.

Sol: Given $u = 2xy$, $v = x^2 - y^2$, $x = r \cos \theta$, $y = r \sin \theta$

$$\frac{\partial u}{\partial x} = 2y \quad \left| \quad \frac{\partial v}{\partial x} = 2x \quad \left| \quad \frac{\partial x}{\partial r} = \cos \theta \quad \left| \quad \frac{\partial y}{\partial r} = \sin \theta \right. \right.$$

$$\frac{\partial u}{\partial y} = 2x \quad \left| \quad \frac{\partial v}{\partial y} = -2y \quad \left| \quad \frac{\partial x}{\partial \theta} = -r \sin \theta \quad \left| \quad \frac{\partial y}{\partial \theta} = r \cos \theta \right. \right.$$

$$\frac{\partial(u, v)}{\partial(r, \theta)} = \frac{\partial(u, v)}{\partial(x, y)} \cdot \frac{\partial(x, y)}{\partial(r, \theta)} = \begin{vmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{vmatrix} \cdot \begin{vmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{vmatrix}$$

$$= \begin{vmatrix} 2y & 2x \\ 2x & -2y \end{vmatrix} \cdot \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix}$$

$$= (-4y^2 - 4x^2) (r \cos^2 \theta + r \sin^2 \theta)$$

$$= -4(x^2 + y^2) r (\cos^2 \theta + \sin^2 \theta) = -4(x^2 + y^2) r \quad (\because \cos^2 \theta + \sin^2 \theta = 1)$$

$$= -4r^3 \quad (\because x^2 + y^2 = r^2)$$

15) Show that the functions $u = x + y - z$, $v = x - y + z$, $w = x^2 + y^2 + z^2 - 2yz$ are dependent. Find the relation between them.

Sol: Given $u = x + y - z$, $v = x - y + z$, $w = x^2 + y^2 + z^2 - 2yz$.

$$\frac{\partial u}{\partial x} = 1 \quad \frac{\partial v}{\partial x} = 1 \quad \frac{\partial w}{\partial x} = 2x$$

$$\frac{\partial u}{\partial y} = 1 \quad \frac{\partial v}{\partial y} = -1 \quad \frac{\partial w}{\partial y} = 2y - 2z$$

$$\frac{\partial u}{\partial z} = -1 \quad \frac{\partial v}{\partial z} = 1 \quad \frac{\partial w}{\partial z} = 2z - 2y$$

$$\frac{\partial(u, v, w)}{\partial(x, y, z)} = \begin{vmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} & \frac{\partial u}{\partial z} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} & \frac{\partial v}{\partial z} \\ \frac{\partial w}{\partial x} & \frac{\partial w}{\partial y} & \frac{\partial w}{\partial z} \end{vmatrix} = \begin{vmatrix} 1 & 1 & -1 \\ 1 & -1 & 1 \\ 2x & 2y - 2z & 2z - 2y \end{vmatrix}$$

$$= 1(-1(2z - 2y) - 1(2y - 2z)) - 1(2z - 2y - 2x) - 1(2y - 2z + 2x)$$

$$= -2z + 2y - 2y + 2z - 2z + 2y + 2x - 2y + 2z - 2x$$

$$= 0$$

$\therefore u, v$ & w are functionally dependent.

$$u+v = x+y-z + x-y+z = 2x \Rightarrow u+v = 2x \text{ --- (1)}$$

$$u-v = x+y-z - (x-y+z) = x+y-z-x+y-z = 2y-2z$$

$$\Rightarrow u-v = 2y-2z \text{ --- (2)}$$

$$\begin{aligned} \therefore (u+v)^2 + (u-v)^2 &= (2x)^2 + (2y-2z)^2 = 4x^2 + 4(y-z)^2 \\ &= 4x^2 + 4(y^2 + z^2 - 2yz) = 4(x^2 + y^2 + z^2 - 2yz) = 4w \quad (\because \text{Given}) \end{aligned}$$

$$\Rightarrow u^2 + v^2 + 2uv + u^2 + v^2 - 2uv = 4w$$

$$\Rightarrow 2u^2 + 2v^2 = 4w \Rightarrow u^2 + v^2 = 2w$$

(H.w) (1) Find the Jacobian of y_1, y_2, y_3 with respect to x_1, x_2, x_3 , if

$$y_1 = \frac{x_2 x_3}{x_1}, \quad y_2 = \frac{x_3 x_1}{x_2}, \quad y_3 = \frac{x_1 x_2}{x_3}$$

(2) Find the Jacobian $\frac{\partial(x, y, z)}{\partial(r, \theta, \phi)}$ of the transformation $x = r \sin \theta \cos \phi$,

$$y = r \sin \theta \sin \phi, \quad z = r \cos \theta$$

(3) Prove $u = x+y+z, v = xy+yz+zx, w = x^2+y^2+z^2$ are functionally dependent. Find the relationship between them.

(A.U) (1b) For the given function $z = \tan^{-1}\left(\frac{x}{y}\right) - (xy)$, verify whether the statement $\frac{d\left(\frac{u}{v}\right)}{dx} = \frac{v u' - u v'}{v^2}$ is correct or not.

$$\frac{d\left(\frac{u}{v}\right)}{dx} = \frac{v u' - u v'}{v^2}$$

$$\frac{d(\tan^{-1} x)}{dx} = \frac{1}{1+x^2}$$

Sol: Given $z = \tan^{-1}\left(\frac{x}{y}\right) - (xy)$

$$\text{LHS } \frac{\partial z}{\partial y} = \frac{1}{1+\left(\frac{x}{y}\right)^2} \times \left(\frac{-1}{y^2}\right) - x = \frac{y^2}{y^2+x^2} \cdot \frac{-x}{y^2} - x = \frac{-x}{x^2+y^2} - x$$

$$\frac{\partial^2 z}{\partial x \partial y} = \frac{(x^2+y^2)(-1) - (-x)2x}{(x^2+y^2)^2} - 1 = \frac{-x^2-y^2+2x^2}{(x^2+y^2)^2} - 1 = \frac{x^2-y^2}{(x^2+y^2)^2} - 1 \text{ --- (1)}$$

$$\text{RHS } \frac{\partial z}{\partial x} = \frac{1}{1+\left(\frac{x}{y}\right)^2} \cdot \frac{1}{y} - y = \frac{y^2}{y^2+x^2} \cdot \frac{1}{y} - y = \frac{y}{x^2+y^2} - y$$

$$\frac{\partial^2 z}{\partial y \partial x} = \frac{(x^2+y^2) \cdot 1 - y(2y)}{(x^2+y^2)^2} - 1 = \frac{x^2+y^2-2y^2}{(x^2+y^2)^2} - 1 = \frac{x^2-y^2}{(x^2+y^2)^2} - 1 \text{ --- (2)}$$

From (1) & (2), $\frac{\partial^2 z}{\partial x \partial y} = \frac{\partial^2 z}{\partial y \partial x}$

(17) If $u = (x^2 + y^2 + z^2)^{-1/2}$ then find the value of $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2}$.

Sol: Given $u = (x^2 + y^2 + z^2)^{-1/2}$

$$\frac{\partial u}{\partial x} = -\frac{1}{2}(x^2 + y^2 + z^2)^{-3/2} \cdot 2x = -x(x^2 + y^2 + z^2)^{-3/2}$$

$$\frac{\partial^2 u}{\partial x^2} = - \left[x(-3/2)(x^2 + y^2 + z^2)^{-5/2} (2x) + (x^2 + y^2 + z^2)^{-3/2} \cdot 1 \right]$$

$$= 3x^2(x^2 + y^2 + z^2)^{-5/2} - (x^2 + y^2 + z^2)^{-3/2} \quad \text{--- (1)}$$

Similarly,

$$\frac{\partial^2 u}{\partial y^2} = 3y^2(x^2 + y^2 + z^2)^{-5/2} - (x^2 + y^2 + z^2)^{-3/2} \quad \text{--- (2)}$$

$$\frac{\partial^2 u}{\partial z^2} = 3z^2(x^2 + y^2 + z^2)^{-5/2} - (x^2 + y^2 + z^2)^{-3/2} \quad \text{--- (3)}$$

$$\begin{aligned} \text{(1) + (2) + (3)} &\Rightarrow \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = 3(x^2 + y^2 + z^2)^{-5/2} (x^2 + y^2 + z^2) - 3(x^2 + y^2 + z^2)^{-3/2} \\ &= 3(x^2 + y^2 + z^2)^{-5/2 + 1} - 3(x^2 + y^2 + z^2)^{-3/2} \\ &= 3(x^2 + y^2 + z^2)^{-3/2} - 3(x^2 + y^2 + z^2)^{-3/2} = 0 \end{aligned}$$

$$\therefore \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = 0$$

(18) If $u = f\left(\frac{y-x}{xy}, \frac{z-x}{xz}\right)$, find $x^2 \frac{\partial u}{\partial x} + y^2 \frac{\partial u}{\partial y} + z^2 \frac{\partial u}{\partial z}$.

Sol: Given $u = f\left(\frac{y-x}{xy}, \frac{z-x}{xz}\right)$


Let $a = \frac{y-x}{xy}$, $b = \frac{z-x}{xz}$

$u = f(a, b)$

$$\begin{aligned} \frac{\partial u}{\partial x} &= \frac{\partial u}{\partial a} \cdot \frac{\partial a}{\partial x} + \frac{\partial u}{\partial b} \cdot \frac{\partial b}{\partial x} \\ &= \frac{\partial u}{\partial a} \cdot \frac{-1}{x^2} + \frac{\partial u}{\partial b} \cdot \frac{-1}{x^2} \end{aligned}$$

$$\frac{\partial u}{\partial x} = -\frac{1}{x^2} \frac{\partial u}{\partial a} - \frac{1}{x^2} \frac{\partial u}{\partial b}$$

$$\frac{\partial u}{\partial y} = \frac{\partial u}{\partial a} \cdot \frac{\partial a}{\partial y} = \frac{\partial u}{\partial a} \cdot \frac{1}{x} \left[\frac{y(1) - (y-x) \cdot 1}{y^2} \right]$$



$$\begin{aligned} \frac{\partial a}{\partial x} &= \frac{1}{y} \left[\frac{x(-1) - (y-x) \cdot 1}{x^2} \right] \\ &= \frac{1}{y} \left[\frac{-x - y + x}{x^2} \right] = \frac{1}{y} \left(\frac{-y}{x^2} \right) \\ \frac{\partial a}{\partial x} &= \frac{-1}{x^2} \\ \frac{\partial b}{\partial x} &= \frac{1}{z} \left[\frac{x(-1) - (z-x) \cdot 1}{x^2} \right] \\ &= \frac{1}{z} \left[\frac{-x - z + x}{x^2} \right] = \frac{-1}{x^2} \end{aligned}$$

$$\frac{\partial u}{\partial y} = \frac{\partial u}{\partial a} \cdot \frac{1}{x} \left[\frac{y - y + x}{y^2} \right] = \frac{1}{y^2} \frac{\partial u}{\partial a}$$

$$\begin{aligned} \frac{\partial u}{\partial z} &= \frac{\partial u}{\partial b} \cdot \frac{\partial b}{\partial z} = \frac{\partial u}{\partial b} \cdot \frac{1}{x} \left[\frac{z(1) - (z-x) \cdot 1}{z^2} \right] \\ &= \frac{\partial u}{\partial b} \cdot \frac{1}{x} \left[\frac{z - z + x}{z^2} \right] = \frac{1}{z^2} \frac{\partial u}{\partial b} \end{aligned}$$


$$\therefore x^2 \frac{\partial u}{\partial x} + y^2 \frac{\partial u}{\partial y} + z^2 \frac{\partial u}{\partial z} = -\frac{\partial u}{\partial a} - \frac{\partial u}{\partial b} + \frac{\partial u}{\partial a} + \frac{\partial u}{\partial b} = 0$$

19) If $u = f(2x - 3y, 3y - 4z, 4z - 2x)$, then find $\frac{1}{2} \frac{\partial u}{\partial x} + \frac{1}{3} \frac{\partial u}{\partial y} + \frac{1}{4} \frac{\partial u}{\partial z}$.

Sol: Given $u = f(2x - 3y, 3y - 4z, 4z - 2x)$

Let $a = 2x - 3y, b = 3y - 4z, c = 4z - 2x$

$u = f(a, b, c)$



$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial a} \cdot \frac{\partial a}{\partial x} + \frac{\partial u}{\partial c} \cdot \frac{\partial c}{\partial x} = 2 \frac{\partial u}{\partial a} - 2 \frac{\partial u}{\partial c}$$

$$\frac{\partial u}{\partial y} = \frac{\partial u}{\partial a} \cdot \frac{\partial a}{\partial y} + \frac{\partial u}{\partial b} \cdot \frac{\partial b}{\partial y} = -3 \frac{\partial u}{\partial a} + 3 \frac{\partial u}{\partial b}$$

$$\frac{\partial u}{\partial z} = \frac{\partial u}{\partial b} \cdot \frac{\partial b}{\partial z} + \frac{\partial u}{\partial c} \cdot \frac{\partial c}{\partial z} = -4 \frac{\partial u}{\partial b} + 4 \frac{\partial u}{\partial c}$$

$$\begin{aligned} \frac{\partial a}{\partial x} &= 2, \quad \frac{\partial c}{\partial x} = -2 \\ \frac{\partial a}{\partial y} &= -3, \quad \frac{\partial b}{\partial y} = 3 \\ \frac{\partial b}{\partial z} &= -4, \quad \frac{\partial c}{\partial z} = 4 \end{aligned}$$

$$\therefore \frac{1}{2} \frac{\partial u}{\partial x} + \frac{1}{3} \frac{\partial u}{\partial y} + \frac{1}{4} \frac{\partial u}{\partial z} = \frac{\partial u}{\partial a} - \frac{\partial u}{\partial c} - \frac{\partial u}{\partial a} + \frac{\partial u}{\partial b} - \frac{\partial u}{\partial b} + \frac{\partial u}{\partial c} = 0$$

20) Find $\frac{dy}{dx}$, if $x^y + y^x = c$, where c is a constant.

$$\frac{d}{dx}(a^x) = a^x \log a$$

Sol: Given $x^y + y^x = c$

Let $f(x, y) = x^y + y^x - c = 0$

$$\frac{\partial f}{\partial x} = yx^{y-1} + y^x \log y \quad \frac{\partial f}{\partial y} = x^y \log x + xy^{x-1}$$


$$\therefore \frac{dy}{dx} = - \frac{\frac{\partial f}{\partial x}}{\frac{\partial f}{\partial y}} = - \left(\frac{yx^{y-1} + y^x \log y}{x^y \log x + xy^{x-1}} \right)$$

1) If $u = f(y - z, z - x, x - y)$, show that $\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} = 0$.

2) Find $\frac{dy}{dx}$ when $y \sin x = x \cos y$.

(21) If $g(x,y) = \psi(u,v)$ where $u = x^2 - y^2$ & $v = 2xy$, then prove that

$$\frac{\partial^2 g}{\partial x^2} + \frac{\partial^2 g}{\partial y^2} = 4(x^2 + y^2) \left[\frac{\partial^2 \psi}{\partial u^2} + \frac{\partial^2 \psi}{\partial v^2} \right].$$



Sol: Given $g(x,y) = \psi(u,v)$, $u = x^2 - y^2$, $v = 2xy$.

$$\frac{\partial u}{\partial x} = 2x \quad \frac{\partial v}{\partial x} = 2y$$

$$\frac{\partial u}{\partial y} = -2y \quad \frac{\partial v}{\partial y} = 2x$$

$$\frac{\partial g}{\partial x} = \frac{\partial \psi}{\partial u} \cdot \frac{\partial u}{\partial x} + \frac{\partial \psi}{\partial v} \cdot \frac{\partial v}{\partial x}$$

$$\frac{\partial g}{\partial y} = \frac{\partial \psi}{\partial u} \cdot \frac{\partial u}{\partial y} + \frac{\partial \psi}{\partial v} \cdot \frac{\partial v}{\partial y}$$

$$\frac{\partial g}{\partial x} = 2x \frac{\partial \psi}{\partial u} + 2y \frac{\partial \psi}{\partial v}$$

$$\frac{\partial g}{\partial y} = -2y \frac{\partial \psi}{\partial u} + 2x \frac{\partial \psi}{\partial v}$$

$$\frac{\partial}{\partial x} = 2x \frac{\partial}{\partial u} + 2y \frac{\partial}{\partial v}$$

$$\frac{\partial}{\partial y} = -2y \frac{\partial}{\partial u} + 2x \frac{\partial}{\partial v}$$

$$\frac{\partial^2 g}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial g}{\partial x} \right) = \left(2x \frac{\partial}{\partial u} + 2y \frac{\partial}{\partial v} \right) \left(2x \frac{\partial \psi}{\partial u} + 2y \frac{\partial \psi}{\partial v} \right)$$

$$\frac{\partial^2 g}{\partial x^2} = 4x^2 \frac{\partial^2 \psi}{\partial u^2} + 4xy \frac{\partial^2 \psi}{\partial u \partial v} + 4xy \frac{\partial^2 \psi}{\partial v \partial u} + 4y^2 \frac{\partial^2 \psi}{\partial v^2} \quad \text{--- (1)}$$

$$\frac{\partial^2 g}{\partial y^2} = \frac{\partial}{\partial y} \left(\frac{\partial g}{\partial y} \right) = \left(-2y \frac{\partial}{\partial u} + 2x \frac{\partial}{\partial v} \right) \left(-2y \frac{\partial \psi}{\partial u} + 2x \frac{\partial \psi}{\partial v} \right)$$

$$\frac{\partial^2 g}{\partial y^2} = 4y^2 \frac{\partial^2 \psi}{\partial u^2} - 4xy \frac{\partial^2 \psi}{\partial u \partial v} - 4xy \frac{\partial^2 \psi}{\partial v \partial u} + 4x^2 \frac{\partial^2 \psi}{\partial v^2} \quad \text{--- (2)}$$

$$\text{(1) + (2)} \Rightarrow \frac{\partial^2 g}{\partial x^2} + \frac{\partial^2 g}{\partial y^2} = 4x^2 \frac{\partial^2 \psi}{\partial u^2} + 4y^2 \frac{\partial^2 \psi}{\partial v^2} + 4y^2 \frac{\partial^2 \psi}{\partial u^2} + 4x^2 \frac{\partial^2 \psi}{\partial v^2}$$

$$= \frac{\partial^2 \psi}{\partial u^2} (4x^2 + 4y^2) + \frac{\partial^2 \psi}{\partial v^2} (4x^2 + 4y^2)$$

$$= (4x^2 + 4y^2) \left(\frac{\partial^2 \psi}{\partial u^2} + \frac{\partial^2 \psi}{\partial v^2} \right)$$

$$\frac{\partial^2 g}{\partial x^2} + \frac{\partial^2 g}{\partial y^2} = 4(x^2 + y^2) \left(\frac{\partial^2 \psi}{\partial u^2} + \frac{\partial^2 \psi}{\partial v^2} \right)$$

(22) If $u = \log(x^3 + y^3 + z^3 - 3xyz)$, show that $\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right)^2 u = \frac{-9}{(x+y+z)^2}$.

Sol: Given $u = \log(x^3 + y^3 + z^3 - 3xyz)$

$$\frac{\partial u}{\partial x} = \frac{1}{x^3 + y^3 + z^3 - 3xyz} (3x^2 - 3yz) = \frac{3(x^2 - yz)}{x^3 + y^3 + z^3 - 3xyz}$$

Similarly,

$$\frac{\partial u}{\partial y} = \frac{3(y^2 - xz)}{x^3 + y^3 + z^3 - 3xyz} \quad \& \quad \frac{\partial u}{\partial z} = \frac{3(z^2 - xy)}{x^3 + y^3 + z^3 - 3xyz}$$

$$\begin{aligned} \therefore \frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} &= \frac{3(x^2 - yz + y^2 - xz + z^2 - xy)}{x^3 + y^3 + z^3 - 3xyz} \\ &= \frac{3(x^2 + y^2 + z^2 - xy - yz - xz)}{(x+y+z)(x^2 + y^2 + z^2 - xy - yz - xz)} = \frac{3}{x+y+z} \end{aligned}$$

$$\Rightarrow \left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right) u = \frac{3}{x+y+z} = 3(x+y+z)^{-1}$$

$$\frac{\partial}{\partial x} \left(\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right) u \right) = 3(-1)(x+y+z)^{-1-1} \cdot 1 = \frac{-3}{(x+y+z)^2} \quad \text{--- (1)}$$

Similarly,

$$\frac{\partial}{\partial y} \left(\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right) u \right) = \frac{-3}{(x+y+z)^2} \quad \text{--- (2)}$$

$$\frac{\partial}{\partial z} \left(\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right) u \right) = \frac{-3}{(x+y+z)^2} \quad \text{--- (3)}$$

$$\text{(1) + (2) + (3)} \Rightarrow \left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right) \left(\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right) u \right) = \frac{-9}{(x+y+z)^2}$$

$$\Rightarrow \left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right)^2 u = \frac{-9}{(x+y+z)^2}$$

(11.10) (1) If $z = f(x, y)$ where $x = r \cos \theta$ & $y = r \sin \theta$, show that

$$\left(\frac{\partial z}{\partial x} \right)^2 + \left(\frac{\partial z}{\partial y} \right)^2 = \left(\frac{\partial z}{\partial r} \right)^2 + \frac{1}{r^2} \left(\frac{\partial z}{\partial \theta} \right)^2.$$

(2) If z is a function of x & y & u & v are other two variables, such that $u = lx + my$, $v = ly - mx$. Show that

$$\frac{\partial^2 z}{\partial x^2} + \frac{\partial^2 z}{\partial y^2} = (l^2 + m^2) \left(\frac{\partial^2 z}{\partial u^2} + \frac{\partial^2 z}{\partial v^2} \right).$$

Taylor's series:

$$f(x, y) = f(a, b) + \frac{1}{1!} [h f_x(a, b) + k f_y(a, b)]$$

$$+ \frac{1}{2!} [h^2 f_{xx}(a, b) + 2hk f_{xy}(a, b) + k^2 f_{yy}(a, b)]$$

$$+ \frac{1}{3!} [h^3 f_{xxx}(a, b) + 3h^2 k f_{xxy}(a, b) + 3hk^2 f_{xyy}(a, b) + k^3 f_{yyy}(a, b)]$$

$$+ \dots$$

Q23) Expand $x^2y^2 + 2x^2y + 3xy^2$ in powers of $(x+2)$ & $(y-1)$ using Taylor's series upto third degree terms.

Sol:

Function	$x = -2, y = 1$ $(-2, 1)$
$f(x, y) = x^2y^2 + 2x^2y + 3xy^2$	$f = (-2)^2(1)^2 + 2(-2)^2(1) + 3(-2)(1)^2 = 4 + 8 - 6 = 6$
$f_x = 2xy^2 + 4xy + 3y^2$	$f_x = 2(-2)(1)^2 + 4(-2)(1) + 3(1)^2 = -4 - 8 + 3 = -9$
$f_y = 2x^2y + 2x^2 + 6xy$	$f_y = 2(-2)^2(1) + 2(-2)^2 + 6(-2)(1) = 8 + 8 - 12 = 4$
$f_{xx} = 2y^2 + 4y$	$f_{xx} = 2(1)^2 + 4(1) = 2 + 4 = 6$
$f_{xy} = 4xy + 4x + 6y$	$f_{xy} = 4(-2)(1) + 4(-2) + 6(1) = -8 - 8 + 6 = -10$
$f_{yy} = 2x^2 + 6x$	$f_{yy} = 2(-2)^2 + 6(-2) = 8 - 12 = -4$
$f_{xxx} = 0$	$f_{xxx} = 0$
$f_{xxy} = 4y + 4$	$f_{xxy} = 4(1) + 4 = 8$
$f_{xyy} = 4x + 6$	$f_{xyy} = 4(-2) + 6 = -8 + 6 = -2$
$f_{yyy} = 0$	$f_{yyy} = 0$

By Taylor's theorem,

$$f(x, y) = f(a, b) + \frac{1}{1!} [h f_x(a, b) + k f_y(a, b)]$$

$$+ \frac{1}{2!} [h^2 f_{xx}(a, b) + 2hk f_{xy}(a, b) + k^2 f_{yy}(a, b)]$$

$$+ \frac{1}{3!} [h^3 f_{xxx}(a, b) + 3h^2 k f_{xxy}(a, b) + 3hk^2 f_{xyy}(a, b) + k^3 f_{yyy}(a, b)]$$

$$+ \dots$$

Here $a = -2, b = 1$; $h = x - a = x - (-2) = x + 2$, $k = y - b = y - 1$

$$\begin{aligned}
 f(x,y) &= 6 + \frac{1}{1!} [(x+2)(-9) + (y-1)(4)] \\
 &+ \frac{1}{2!} [(x+2)^2(6) + 2(x+2)(y-1)(-10) + (y-1)^2(-4)] \\
 &+ \frac{1}{3!} [(x+2)^3(0) + 3(x+2)^2(y-1)(8) + 3(x+2)(y-1)^2(-2) + (y-1)^3(0)] \\
 &+ \dots \\
 &= 6 - 9(x+2) + 4(y-1) + \frac{1}{2} [6(x+2)^2 - 20(x+2)(y-1) - 4(y-1)^2] \\
 &+ \frac{1}{6} [24(x+2)^2(y-1) - 6(x+2)(y-1)^2] + \dots
 \end{aligned}$$

- 11.10
- Q1 Obtain the Taylor's series expansion of $x^3 + y^3 + xy^2$ in terms of powers of $(x-1)$ & $(y-2)$ up to third degree terms.
 - Q2 Find Taylor's series expansion of function of $f(x) = \sqrt{1+x+y^2}$ in powers of $(x-1)$ & y up to second degree terms.
 - Q3 Obtain the Taylor's series expansion of $e^x \sin y$ in terms of powers of x & y up to third degree terms.

Sol:

Function	$(0,0) [x=0, y=0]$
$f(x,y) = e^x \sin y$	$f = e^0 \sin 0 = (1)(0) = 0$
$f_x = e^x \sin y$	$f_x = e^0 \sin 0 = (1)(0) = 0$
$f_y = e^x \cos y$	$f_y = e^0 \cos 0 = (1)(1) = 1$
$f_{xx} = e^x \sin y$	$f_{xx} = e^0 \sin 0 = (1)(0) = 0$
$f_{xy} = e^x \cos y$	$f_{xy} = e^0 \cos 0 = (1)(1) = 1$
$f_{yy} = -e^x \sin y$	$f_{yy} = -e^0 \sin 0 = -(1)(0) = 0$
$f_{xxx} = e^x \sin y$	$f_{xxx} = e^0 \sin 0 = (1)(0) = 0$
$f_{xxy} = e^x \cos y$	$f_{xxy} = e^0 \cos 0 = (1)(1) = 1$
$f_{xyy} = -e^x \sin y$	$f_{xyy} = -e^0 \sin 0 = -(1)(0) = 0$
$f_{yyy} = -e^x \cos y$	$f_{yyy} = -e^0 \cos 0 = -(1)(1) = -1$

Here $a=0, b=0, h=x-a=x-0=x, k=y-b=y-0=y$

By Taylor's theorem,

$$f(x,y) = f(a,b) + \frac{1}{1!} [h f_x(a,b) + k f_y(a,b)]$$

$$+ \frac{1}{2!} [h^2 f_{xx}(a,b) + 2hk f_{xy}(a,b) + k^2 f_{yy}(a,b)]$$

$$+ \frac{1}{3!} [h^3 f_{xxx}(a,b) + 3h^2k f_{xxy}(a,b) + 3hk^2 f_{xyy}(a,b) + k^3 f_{yyy}(a,b)]$$

$$+ \dots$$

$$f(x,y) = 0 + \frac{1}{1!} [x(0) + y(1)] + \frac{1}{2!} [x^2(0) + 2xy(1) + y^2(0)]$$

$$+ \frac{1}{3!} [x^3(0) + 3x^2y(1) + 3xy^2(0) + y^3(-1)] + \dots$$

$$= y + \frac{1}{2} (2xy) + \frac{1}{6} (3x^2y - y^3) + \dots$$

$$= y + xy + \frac{1}{6} (3x^2y - y^3) + \dots$$

25) Expand the function $\sin xy$ in powers of $x-1$ & $y-\frac{\pi}{2}$ upto second degree terms, using Taylor's series.

Sol:

Function	$x=1, y=\frac{\pi}{2}$
$f(x,y) = \sin xy$	$f = \sin(1)(\frac{\pi}{2}) = \sin \frac{\pi}{2} = 1$
$f_x = \cos xy \cdot y$	$f_x = \cos(1)(\frac{\pi}{2}) \cdot \frac{\pi}{2} = \cos \frac{\pi}{2} \cdot \frac{\pi}{2} = 0 \cdot \frac{\pi}{2} = 0$
$f_y = \cos xy \cdot x$	$f_y = \cos(1)(\frac{\pi}{2}) \cdot 1 = \cos \frac{\pi}{2} = 0$
$f_{xx} = y(-\sin xy) \cdot y$ $= -y^2 \sin xy$	$f_{xx} = -(\frac{\pi}{2})^2 \sin(1)(\frac{\pi}{2}) = -\frac{\pi^2}{4} \sin \frac{\pi}{2} = -\frac{\pi^2}{4}$
$f_{xy} = \cos xy \cdot 1 + y(-\sin xy) \cdot x$ $= \cos xy - xy \sin xy$	$f_{xy} = \cos(1)(\frac{\pi}{2}) - (1)(\frac{\pi}{2}) \sin(1)(\frac{\pi}{2})$ $= \cos \frac{\pi}{2} - \frac{\pi}{2} \sin \frac{\pi}{2} = 0 - \frac{\pi}{2}(1) = -\frac{\pi}{2}$
$f_{yy} = x(-\sin xy) \cdot x$ $= -x^2 \sin xy$	$f_{yy} = -(1)^2 \sin(1)(\frac{\pi}{2}) = -\sin(\frac{\pi}{2}) = -1$

By Taylor's theorem, $f(x,y) = f(a,b) + \frac{1}{1!} [h f_x(a,b) + k f_y(a,b)]$

$$h = x - a = x - 1$$

$$k = y - b = y - \frac{\pi}{2}$$

$$+ \frac{1}{2!} [h^2 f_{xx}(a,b) + 2hk f_{xy}(a,b) + k^2 f_{yy}(a,b)] + \dots$$

$$f(x,y) = 1 + \frac{1}{1!} [(x-1)(0) + (y-\pi/2)(0)]$$

$$+ \frac{1}{2!} [(x-1)^2 \left(\frac{-\pi^2}{4}\right) + 2(x-1)(y-\pi/2)(-\pi/2) + (y-\pi/2)^2(-1)] + \dots$$

$$= 1 + \frac{1}{2} \left[-\frac{\pi^2}{4}(x-1)^2 - \pi(x-1)(y-\pi/2) - (y-\pi/2)^2 \right] + \dots$$

(26) Expand $e^x \log(1+y)$ in powers of x & y upto the third degree terms, using Taylor's series. $e^0 = 1, \log 1 = 0$

Sol:

Function	$x=0, y=0$
$f(x,y) = e^x \log(1+y)$	$f = e^0 \log(1+0) = e^0 \log 1 = (1)(0) = 0$
$f_x = e^x \log(1+y)$	$f_x = e^0 \log(1+0) = e^0 \log 1 = (1)(0) = 0$
$f_y = e^x \cdot \frac{1}{1+y} \cdot 1 = e^x (1+y)^{-1}$	$f_y = e^0 (1+0)^{-1} = e^0 (1)^{-1} = (1)(1) = 1$
$f_{xx} = e^x \log(1+y)$	$f_{xx} = e^0 \log(1+0) = e^0 \log 1 = (1)(0) = 0$
$f_{xy} = e^x (1+y)^{-1}$	$f_{xy} = e^0 (1+0)^{-1} = (1)(1) = 1$
$f_{yy} = e^x (-1)(1+y)^{-2} = -e^x (1+y)^{-2}$	$f_{yy} = -e^0 (1+0)^{-2} = -(1)(1)^{-2} = -1$
$f_{xxx} = e^x \log(1+y)$	$f_{xxx} = e^0 \log(1+0) = (1)(0) = 0$
$f_{xxy} = e^x (1+y)^{-1}$	$f_{xxy} = e^0 (1+0)^{-1} = (1)(1) = 1$
$f_{xyy} = -e^x (1+y)^{-2}$	$f_{xyy} = -e^0 (1+0)^{-2} = -(1)(1) = -1$
$f_{yyy} = -e^x (-2)(1+y)^{-3} = 2e^x (1+y)^{-3}$	$f_{yyy} = 2e^0 (1+0)^{-3} = 2(1)(1) = 2$

By Taylor's Theorem,

$$f(x,y) = f(a,b) + \frac{1}{1!} [h f_x(a,b) + k f_y(a,b)]$$

$$+ \frac{1}{2!} [h^2 f_{xx}(a,b) + 2hk f_{xy}(a,b) + k^2 f_{yy}(a,b)]$$

$$+ \frac{1}{3!} [h^3 f_{xxx}(a,b) + 3h^2 k f_{xxy}(a,b) + 3h k^2 f_{xyy}(a,b) + k^3 f_{yyy}(a,b)]$$

$$+ \dots$$

Here $a=0, b=0$

$$h = x - a = x - 0 = x, \quad k = y - b = y - 0 = y$$

$$f(x, y) = 0 + \frac{1}{1!} [x(0) + y(1)] + \frac{1}{2!} [x^2(0) + 2xy(1) + y^2(-1)]$$

$$+ \frac{1}{3!} [x^3(0) + 3x^2y(1) + 3xy^2(-1) + y^3(2)] + \dots$$

$$= y + \frac{1}{2} (2xy - y^2) + \frac{1}{6} (3x^2y - 3xy^2 + 2y^3) + \dots$$

- H.W
- ① Expand $e^x \cos y$ about $(0, \pi/2)$ upto the third term using Taylor's series.
 - ② Obtain terms upto the third degree in the Taylor's series expansion of $e^x \sin y$ around the point $(1, \pi/2)$.
 - ③ Expand $f(x, y) = e^{xy}$ in Taylor series at $(1, 1)$ upto second degree.

Maxima & minima for functions of two variables:

Definitions:

Extremum value:

$f(a, b)$ is said to be an extremum value of $f(x, y)$ if it is either a maximum or a minimum.

Notations: $\frac{\partial f}{\partial x} = f_x$, $\frac{\partial f}{\partial y} = f_y$, $\frac{\partial^2 f}{\partial x^2} = f_{xx}$, $\frac{\partial^2 f}{\partial x \partial y} = f_{xy}$, $\frac{\partial^2 f}{\partial y^2} = f_{yy}$

Sufficient conditions:

If $f_x(a, b) = 0$, $f_y(a, b) = 0$ & $f_{xx}(a, b) = A$, $f_{xy}(a, b) = B$, $f_{yy}(a, b) = C$,

then

- (i) $f(a, b)$ is maximum value if $AC - B^2 > 0$ & $A < 0$ (or $B < 0$)
- (ii) $f(a, b)$ is minimum value if $AC - B^2 > 0$ & $A > 0$ (or $B > 0$)
- (iii) $f(a, b)$ is not an extremum (saddle) if $AC - B^2 < 0$ &
- (iv) If $AC - B^2 = 0$, then the test is inconclusive.

Stationary value:

A function $f(x, y)$ is said to be stationary at (a, b) or $f(a, b)$ is said to be a stationary value of $f(x, y)$ if $f_x(a, b) = 0$ & $f_y(a, b) = 0$.

Note: Every extremum value is a stationary value but a stationary value need not be an extremum value.

(27) Examine $f(x, y) = x^3 + 3xy^2 - 15x^2 - 15y^2 + 72x$ for extreme values.

Sol: Given $f(x, y) = x^3 + 3xy^2 - 15x^2 - 15y^2 + 72x$

$$f_x = 3x^2 + 3y^2 - 30x + 72$$

$$f_y = 6xy - 30y$$

$$\begin{array}{r|l} x & + \\ \hline 24 & -10 \\ -6 & -4 \\ x-6 & x-4 \end{array}$$

Stationary points:

$$f_x = 0$$

$$3x^2 + 3y^2 - 30x + 72 = 0 \text{ --- (1)}$$

$$y = 0 \text{ in (1)}$$

$$3x^2 - 30x + 72 = 0 \Rightarrow x^2 - 10x + 24 = 0$$

$$\Rightarrow (x-6)(x-4) = 0$$

$$\Rightarrow x = 4, 6$$

$$f_y = 0$$

$$6xy - 30y = 0 \Rightarrow 6y(x-5) = 0$$

$$\Rightarrow y = 0, x = 5$$

\therefore The points are $(4, 0)$ & $(6, 0)$

$$x = 5 \text{ in (1)}$$

$$75 + 3y^2 - 150 + 72 = 0 \Rightarrow 3y^2 - 3 = 0 \Rightarrow 3y^2 = 3 \Rightarrow y^2 = 1 \Rightarrow y = \pm\sqrt{1} = \pm 1$$

\therefore The points are $(5, 1)$ & $(5, -1)$.

Hence the stationary points are $(4, 0), (6, 0), (5, 1)$ & $(5, -1)$.

$$A = f_{xx} = 6x - 30 \quad ; \quad B = f_{xy} = 6y \quad ; \quad C = f_{yy} = 6x - 30$$

	$(4, 0)$	$(6, 0)$	$(5, 1)$	$(5, -1)$
$A = 6x - 30$	$-6 < 0$	$6 > 0$	0	0
$B = 6y$	0	0	6	-6
$C = 6x - 30$	-6	6	0	0
$Ac - B^2$	$36 > 0$	$36 > 0$	$-36 < 0$	$-36 < 0$
Conclusion	Maximum value	Minimum value	Saddle point	Saddle point

$$f(4, 0) = (4)^3 + 3(4)(0)^2 - 15(4)^2 - 15(0)^2 + 72(4) = 112$$

$$f(6, 0) = (6)^3 + 3(6)(0)^2 - 15(6)^2 - 15(0)^2 + 72(6) = 108$$

Hence the maximum value is 112 & the minimum value is 108.

Q28 Find the maxima & minima of $f(x,y) = x^4 + y^4 - 2x^2 + 4xy - 2y^2$.

Sol: Given $f(x,y) = x^4 + y^4 - 2x^2 + 4xy - 2y^2$.

$$f_x = 4x^3 - 4x + 4y$$

$$A = f_{xx} = 12x^2 - 4$$

$$C = f_{yy} = 12y^2 - 4$$

$$f_y = 4y^3 + 4x - 4y$$

$$B = f_{xy} = 4$$

Stationary points:

$$f_x = 0$$

$$\Rightarrow 4x^3 - 4x + 4y = 0$$

$$\Rightarrow x^3 - x + y = 0 \text{ --- (1)}$$

$$f_y = 0$$

$$\Rightarrow 4y^3 + 4x - 4y = 0$$

$$\Rightarrow y^3 + x - y = 0 \text{ --- (2)}$$

$$\text{(1) + (2)} \Rightarrow x^3 - x + y + y^3 + x - y = 0 \Rightarrow x^3 + y^3 = 0 \Rightarrow x^3 = -y^3 \Rightarrow x = -y$$

$$\Rightarrow \boxed{y = -x} \text{ --- (3)}$$

Substituting (3) in (1),

$$x^3 - x - x = 0 \Rightarrow x^3 - 2x = 0 \Rightarrow x(x^2 - 2) = 0$$

$$\Rightarrow x = 0, x^2 - 2 = 0$$

$$\Rightarrow x = 0, x^2 = 2$$

$$\Rightarrow x = 0, x = \pm\sqrt{2} \text{ --- (4)}$$

Substituting (4) in (3),

$$x = 0 \Rightarrow y = 0; \quad x = \sqrt{2} \Rightarrow y = -\sqrt{2}; \quad x = -\sqrt{2} \Rightarrow y = \sqrt{2}$$

Hence the stationary points are $(0,0), (\sqrt{2}, -\sqrt{2})$ & $(-\sqrt{2}, \sqrt{2})$.

	$(0,0)$	$(\sqrt{2}, -\sqrt{2})$	$(-\sqrt{2}, \sqrt{2})$
$A = 12x^2 - 4$	-4	$20 > 0$	$20 > 0$
$B = 4$	4	4	4
$C = 12y^2 - 4$	-4	20	20
$AC - B^2$	0	$384 > 0$	$384 > 0$
Conclusion	Inconclusive	Minimum value	Minimum value.

$$f(\sqrt{2}, -\sqrt{2}) = (\sqrt{2})^4 + (-\sqrt{2})^4 - 2(\sqrt{2})^2 + 4(\sqrt{2})(-\sqrt{2}) - 2(-\sqrt{2})^2 = 4 + 4 - 4 - 8 - 4 = -8$$

$$f(-\sqrt{2}, \sqrt{2}) = (-\sqrt{2})^4 + (\sqrt{2})^4 - 2(-\sqrt{2})^2 + 4(-\sqrt{2})(\sqrt{2}) - 2(\sqrt{2})^2 = 4 + 4 - 4 - 8 - 4 = -8$$

Hence the minimum value is -8.

29) Find the extreme values of $f(x, y) = x^3 y^2 (1 - x - y)$.

Sol: Given $f(x, y) = x^3 y^2 (1 - x - y) = x^3 y^2 - x^4 y^2 - x^3 y^3$

$$f_x = 3x^2 y^2 - 4x^3 y^2 - 3x^2 y^3$$

$$f_y = 2x^3 y - 2x^4 y - 3x^3 y^2$$

$$A = f_{xx} = 6xy^2 - 12x^2 y^2 - 6xy^3$$

$$B = f_{xy} = 6x^2 y - 8x^3 y - 9x^2 y^2$$

$$C = f_{yy} = 2x^3 - 2x^4 - 6x^3 y$$

Stationary points:

$$f_x = 0$$

$$\Rightarrow 3x^2 y^2 - 4x^3 y^2 - 3x^2 y^3 = 0$$

$$\Rightarrow x^2 y^2 (3 - 4x - 3y) = 0$$

$$\Rightarrow x = 0, y = 0, 3 - 4x - 3y = 0$$

$$\Rightarrow x = 0, y = 0, 4x + 3y = 3 \text{ --- (1)}$$

$$f_y = 0$$

$$\Rightarrow 2x^3 y - 2x^4 y - 3x^3 y^2 = 0$$

$$\Rightarrow x^3 y (2 - 2x - 3y) = 0$$

$$\Rightarrow x = 0, y = 0, 2 - 2x - 3y = 0$$

$$\Rightarrow x = 0, y = 0, 2x + 3y = 2 \text{ --- (2)}$$

$$\text{(1) - (2)} \Rightarrow 4x + 3y - 2x - 3y = 3 - 2 \Rightarrow 2x = 1 \Rightarrow \boxed{x = \frac{1}{2}}$$

Substituting $x = \frac{1}{2}$ in (2),

$$2\left(\frac{1}{2}\right) + 3y = 2 \Rightarrow 1 + 3y = 2 \Rightarrow 3y = 2 - 1 \Rightarrow 3y = 1 \Rightarrow \boxed{y = \frac{1}{3}}$$

Hence the stationary points are $(0, 0)$ & $(\frac{1}{2}, \frac{1}{3})$.

	$(0, 0)$	$(\frac{1}{2}, \frac{1}{3})$
$A = 6xy^2 - 12x^2 y^2 - 6xy^3$	0	$-\frac{1}{9} < 0$
$B = 6x^2 y - 8x^3 y - 9x^2 y^2$	0	$-\frac{1}{12}$
$C = 2x^3 - 2x^4 - 6x^3 y$	0	$-\frac{1}{8}$
$AC - B^2$	0	$\frac{1}{144} > 0$
Conclusion	Inconclusive	Maximum value

$$f\left(\frac{1}{2}, \frac{1}{3}\right) = \left(\frac{1}{2}\right)^3 \left(\frac{1}{3}\right)^2 \left(1 - \frac{1}{2} - \frac{1}{3}\right) = \frac{1}{8} \times \frac{1}{9} \times \left(\frac{1}{6}\right) = \frac{1}{432}$$

Hence the maximum value is $\frac{1}{432}$.

30) Discuss the maxima & minima of the function $f(x,y) = x^3 + y^3 - 3axy$.

Sol. Given $f(x,y) = x^3 + y^3 - 3axy$.

$f_x = 3x^2 - 3ay$ $A = f_{xx} = 6x$ $C = f_{yy} = 6y$

$f_y = 3y^2 - 3ax$ $B = f_{xy} = -3a$

Stationary points:

$f_x = 0$
 $\Rightarrow 3x^2 - 3ay = 0$
 $\Rightarrow x^2 - ay = 0 \Rightarrow x^2 = ay \text{ --- (1)}$

$f_y = 0$
 $\Rightarrow 3y^2 - 3ax = 0 \Rightarrow y^2 - ax = 0$
 $\Rightarrow y^2 = ax \text{ --- (2)}$

(1) $\Rightarrow y = \frac{x^2}{a} \text{ --- (3)}$

Substituting (3) in (2), $\left(\frac{x^2}{a}\right)^2 = ax \Rightarrow \frac{x^4}{a^2} = ax \Rightarrow \frac{x^4}{x} = a^3 \Rightarrow x^3 = a^3$
 $\Rightarrow x = a \text{ --- (4)}$

Substituting (4) in (3), $y = \frac{a^2}{a} = a \Rightarrow y = a$

Hence the stationary point is (a, a) .

	(a, a)
$A = 6x$	$6a$
$B = -3a$	$-3a$
$C = 6y$	$6a$
$Ac - B^2$	$36a^2 - 9a^2 = 27a^2 > 0$
Conclusion	

If $a > 0$, then $A > 0 \Rightarrow$
 Minimum value at (a, a) .
 If $a < 0$, then $A < 0 \Rightarrow$
 Maximum value at (a, a) .

$f(a, a) = a^3 + a^3 - 3a(a)(a) = 2a^3 - 3a^3 = -a^3$

Hence the maximum or minimum value at (a, a) is $-a^3$.

10) 1) Find the maximum or minimum values of $f(x,y) = 3x^2 - y^2 + x^3$.


2) Find the maximum or minimum values of $f(x,y) = x^2 + y^2 + 6x + 12$.

3) Examine $x^3y^2(12-x-y)$ for extreme values.

4) Find the maxima & minima of $xy(a-x-y)$.

31 If $z = f(x, y)$ where $x = r \cos \theta$ & $y = r \sin \theta$, show that

$$\left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2 = \left(\frac{\partial z}{\partial r}\right)^2 + \frac{1}{r^2} \left(\frac{\partial z}{\partial \theta}\right)^2$$



Sol: Given $z = f(x, y)$, $x = r \cos \theta$, $y = r \sin \theta$

$$\frac{\partial x}{\partial r} = \cos \theta = \frac{x}{r} \quad \left| \quad \frac{\partial y}{\partial r} = \sin \theta = \frac{y}{r}$$

$$\frac{\partial x}{\partial \theta} = -r \sin \theta = -y \quad \left| \quad \frac{\partial y}{\partial \theta} = r \cos \theta = x$$

$$\frac{\partial z}{\partial r} = \frac{\partial z}{\partial x} \cdot \frac{\partial x}{\partial r} + \frac{\partial z}{\partial y} \cdot \frac{\partial y}{\partial r} = \frac{x}{r} \frac{\partial z}{\partial x} + \frac{y}{r} \frac{\partial z}{\partial y}$$

$$\frac{\partial z}{\partial \theta} = \frac{\partial z}{\partial x} \cdot \frac{\partial x}{\partial \theta} + \frac{\partial z}{\partial y} \cdot \frac{\partial y}{\partial \theta} = -y \frac{\partial z}{\partial x} + x \frac{\partial z}{\partial y}$$

$$\begin{aligned} \left(\frac{\partial z}{\partial r}\right)^2 &= \left(\frac{x}{r} \frac{\partial z}{\partial x} + \frac{y}{r} \frac{\partial z}{\partial y}\right)^2 = \frac{1}{r^2} \left(x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y}\right)^2 \\ &= \frac{1}{r^2} \left(x^2 \left(\frac{\partial z}{\partial x}\right)^2 + y^2 \left(\frac{\partial z}{\partial y}\right)^2 + 2xy \frac{\partial z}{\partial x} \frac{\partial z}{\partial y}\right) \end{aligned}$$

$$\begin{aligned} \left(\frac{\partial z}{\partial \theta}\right)^2 &= \left(-y \frac{\partial z}{\partial x} + x \frac{\partial z}{\partial y}\right)^2 = \left(x \frac{\partial z}{\partial y} - y \frac{\partial z}{\partial x}\right)^2 \\ &= x^2 \left(\frac{\partial z}{\partial y}\right)^2 + y^2 \left(\frac{\partial z}{\partial x}\right)^2 - 2xy \frac{\partial z}{\partial x} \frac{\partial z}{\partial y} \end{aligned}$$

$$\begin{aligned} \therefore \left(\frac{\partial z}{\partial r}\right)^2 + \frac{1}{r^2} \left(\frac{\partial z}{\partial \theta}\right)^2 &= \frac{1}{r^2} \left(x^2 \left(\frac{\partial z}{\partial x}\right)^2 + y^2 \left(\frac{\partial z}{\partial y}\right)^2 + 2xy \frac{\partial z}{\partial x} \frac{\partial z}{\partial y}\right) \\ &\quad + \frac{1}{r^2} \left(x^2 \left(\frac{\partial z}{\partial y}\right)^2 + y^2 \left(\frac{\partial z}{\partial x}\right)^2 - 2xy \frac{\partial z}{\partial x} \frac{\partial z}{\partial y}\right) \\ &= \frac{1}{r^2} \left(x^2 \left(\frac{\partial z}{\partial x}\right)^2 + y^2 \left(\frac{\partial z}{\partial y}\right)^2 + x^2 \left(\frac{\partial z}{\partial y}\right)^2 + y^2 \left(\frac{\partial z}{\partial x}\right)^2\right) \\ &= \frac{1}{r^2} \left[\left(\frac{\partial z}{\partial x}\right)^2 (x^2 + y^2) + \left(\frac{\partial z}{\partial y}\right)^2 (x^2 + y^2)\right] \\ &= \frac{1}{r^2} \left[(x^2 + y^2) \left(\left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2\right)\right] \\ &= \frac{1}{r^2} \times r^2 \times \left(\left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2\right) \\ &= \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2 \end{aligned}$$

Hence $\left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2 = \left(\frac{\partial z}{\partial r}\right)^2 + \frac{1}{r^2} \left(\frac{\partial z}{\partial \theta}\right)^2$

Lagrange's method of undetermined multipliers:

(32) A thin closed rectangular box is to have one edge equal to twice the other & constant volume 72 m³. Find the least surface area of the box.

Sol: Let x, y, 2y be the length, breadth & height of the box respectively.

Surface area = 2xy + 2(y)(2y) + 2(x)(2y) = 2xy + 4y² + 4xy = 6xy + 4y²

Volume = (x)(y)(2y) = 2xy² = 72 ⇒ xy² = $\frac{72}{2}$ = 36 ⇒ xy² = 36 — (*)

F = (6xy + 4y²) + λ(xy² - 36) = 6xy + 4y² + λxy² - 36λ

F_x = 6y + λy² ; F_y = 6x + 8y + 2λxy

F_x = 0
⇒ 6y + λy² = 0 ⇒ 6y = -λy²
⇒ 6 = -λy ⇒ $\frac{6}{y} = -\lambda$ — ①

F_y = 0
⇒ 6x + 8y + 2λxy = 0 ⇒ 6x + 8y = -2λxy
⇒ 3x + 4y = -λxy ⇒ $\frac{3x + 4y}{xy} = -\lambda$
⇒ $\frac{3}{y} + \frac{4}{x} = -\lambda$ — ②

From ① & ②, $\frac{6}{y} = \frac{3}{y} + \frac{4}{x} \Rightarrow \frac{6}{y} - \frac{3}{y} = \frac{4}{x} \Rightarrow \frac{3}{y} = \frac{4}{x}$
⇒ 3x = 4y ⇒ $y = \frac{3}{4}x$ — ③

Substituting ③ in (*),

$x(\frac{3}{4}x)^2 = 36 \Rightarrow x \frac{9}{16}x^2 = 36 \Rightarrow \frac{9}{16}x^3 = 36 \Rightarrow x^3 = \frac{36 \times 16}{9} = 64 = 4^3$
⇒ $x = 4$

∴ y = $\frac{3}{4}(4) = 3 \Rightarrow y = 3$

∴ Least surface area = 6xy + 4y² = 6(4)(3) + 4(3)² = 108.

(33) Find the dimensions of the rectangular box without a top of maximum capacity, whose surface area is 108 sq.cm.

Sol: Let x, y, z be the length, breadth & height of the box.

Surface area = xy + 2yz + 2zx = 108 — ①

Volume = xyz

$F = xyz + \lambda(xy + 2yz + 2zx - 108) = xyz + \lambda xy + 2\lambda yz + 2\lambda zx - 108\lambda$

$F_x = yz + \lambda y + 2\lambda z$; $F_y = xz + \lambda x + 2\lambda z$; $F_z = xy + 2\lambda y + 2\lambda x$

$F_x = 0$

$\Rightarrow yz + \lambda(y + 2z) = 0$

$\Rightarrow yz = -\lambda(y + 2z)$

$\Rightarrow \frac{yz}{y + 2z} = -\lambda$

$\Rightarrow \frac{y + 2z}{yz} = \frac{-1}{\lambda}$

$\Rightarrow \frac{1}{z} + \frac{2}{y} = \frac{-1}{\lambda}$ — (2)

From (2) & (3),

$\frac{1}{z} + \frac{2}{y} = \frac{1}{z} + \frac{2}{x}$

$\Rightarrow \frac{2}{y} = \frac{2}{x} \Rightarrow 2x = 2y$

$\Rightarrow \boxed{x = y}$ — (5)

$F_y = 0$

$\Rightarrow xz + \lambda(x + 2z) = 0$

$\Rightarrow xz = -\lambda(x + 2z)$

$\Rightarrow \frac{xz}{x + 2z} = -\lambda$

$\Rightarrow \frac{x + 2z}{xz} = \frac{-1}{\lambda}$

$\Rightarrow \frac{1}{z} + \frac{2}{x} = \frac{-1}{\lambda}$ — (3)

From (3) & (4),

$\frac{1}{z} + \frac{2}{x} = \frac{2}{x} + \frac{2}{y}$

$\Rightarrow \frac{1}{z} = \frac{2}{y}$

$\Rightarrow \boxed{y = 2z}$ — (6)

$F_z = 0$

$\Rightarrow xy + \lambda(2y + 2x) = 0$

$\Rightarrow xy = -\lambda(2y + 2x)$

$\Rightarrow \frac{xy}{2y + 2x} = -\lambda$

$\Rightarrow \frac{2y + 2x}{xy} = \frac{-1}{\lambda}$

$\Rightarrow \frac{2}{x} + \frac{2}{y} = \frac{-1}{\lambda}$ — (4)

From (5) & (6), $x = y = 2z$

$\therefore \textcircled{1} \Rightarrow 2y + 2yz + 2zx = 108 \Rightarrow (2z)(2z) + 2(2z)z + 2z(2z) = 108$

$\Rightarrow 4z^2 + 4z^2 + 4z^2 = 108 \Rightarrow 12z^2 = 108 \Rightarrow z^2 = 9 \Rightarrow \boxed{z = 3}$

$\therefore x = 6, y = 6, z = 3$

Maximum volume = $xyz = (6)(6)(3) = 108$.

34) Find the shortest & the longest distances from the point $(1, 2, -1)$ to the sphere $x^2 + y^2 + z^2 = 24$.

Sol: Given $x^2 + y^2 + z^2 = 24$ & $(1, 2, -1)$

$d = \sqrt{(x-1)^2 + (y-2)^2 + (z+1)^2}$

$\Rightarrow d^2 = (x-1)^2 + (y-2)^2 + (z+1)^2$

$F = (x-1)^2 + (y-2)^2 + (z+1)^2 + \lambda(x^2 + y^2 + z^2 - 24)$

$F_x = 2(x-1) + 2x\lambda$; $F_y = 2(y-2) + 2y\lambda$; $F_z = 2(z+1) + 2z\lambda$

Here $x_1 = 1, y_1 = 2, z_1 = -1$
 $d = \sqrt{(x-x_1)^2 + (y-y_1)^2 + (z-z_1)^2}$

$$F_x = 0$$

$$2(x-1) + 2x\lambda = 0$$

$$x-1 + x\lambda = 0$$

$$x + x\lambda = 1$$

$$x(1+\lambda) = 1$$

$$x = \frac{1}{1+\lambda} \text{ --- (1)}$$

$$F_y = 0$$

$$2(y-2) + 2y\lambda = 0$$

$$y-2 + y\lambda = 0$$

$$y + y\lambda = 2$$

$$y(1+\lambda) = 2$$

$$\frac{y}{2} = \frac{1}{1+\lambda} \text{ --- (2)}$$

$$F_z = 0$$

$$2(z+1) + 2z\lambda = 0$$

$$z+1 + z\lambda = 0$$

$$z + z\lambda = -1$$

$$z(1+\lambda) = -1$$

$$-z = \frac{1}{1+\lambda} \text{ --- (3)}$$

From (1), (2) & (3),

$$x = \frac{y}{2} = -z \Rightarrow x = -z, \frac{y}{2} = -z \Rightarrow x = -z, y = -2z$$

$$\therefore x^2 + y^2 + z^2 = 24 \Rightarrow (-z)^2 + (-2z)^2 + z^2 = 24$$

$$\Rightarrow z^2 + 4z^2 + z^2 = 24 \Rightarrow 6z^2 = 24 \Rightarrow z^2 = 4 \Rightarrow z = \pm\sqrt{4} = \pm 2$$

$$z = 2 \Rightarrow x = -2, y = -2(2) = -4$$

$$z = -2 \Rightarrow x = 2, y = -2(-2) = 4$$

Hence the stationary points are $(-2, -4, 2)$ & $(2, 4, -2)$.

$$d = \sqrt{(-2-1)^2 + (-4-2)^2 + (2+1)^2} = \sqrt{9 + 36 + 9} = \sqrt{54} = 3\sqrt{6}$$

$$d = \sqrt{(2-1)^2 + (4-2)^2 + (-2+1)^2} = \sqrt{1 + 4 + 1} = \sqrt{6}$$

Hence the shortest & longest distances are $\sqrt{6}$ & $3\sqrt{6}$ respectively.

10 (35) Find the minimum distance from the point $(1, 2, 0)$ to the cone

$$z^2 = x^2 + y^2$$

Sol: Given $z^2 = x^2 + y^2$ & $(1, 2, 0)$

$$d = \sqrt{(x-1)^2 + (y-2)^2 + (z-0)^2}$$

$$d^2 = (x-1)^2 + (y-2)^2 + z^2$$

$$F = (x-1)^2 + (y-2)^2 + z^2 + \lambda(z^2 - x^2 - y^2)$$

$$F_x = 2(x-1) - 2x\lambda$$

$$; F_y = 2(y-2) - 2y\lambda$$

$$; F_z = 2z + 2z\lambda$$

$$F_x = 0$$

$$F_y = 0$$

$$F_z = 0$$

$$2(x-1) - 2x\lambda = 0$$

$$2(y-2) - 2y\lambda = 0$$

$$2z + 2z\lambda = 0$$

$$x-1 - x\lambda = 0$$

$$y-2 - y\lambda = 0$$

$$z + z\lambda = 0$$

$$x-1 = x\lambda$$

$$y-2 = y\lambda$$

$$z = -z\lambda$$

$$\frac{x-1}{x} = \lambda$$

$$\frac{y-2}{y} = \lambda$$

$$\frac{z}{-z} = \lambda$$

$$1 - \frac{1}{x} = \lambda \text{ --- (1)}$$

$$1 - \frac{2}{y} = \lambda \text{ --- (2)}$$

$$\boxed{\lambda = -1} \text{ --- (3)}$$

Substituting (3) in (1) & (2),

$$1 - \frac{1}{x} = -1 \Rightarrow 1 + 1 = \frac{1}{x} \Rightarrow 2 = \frac{1}{x} \Rightarrow \boxed{x = \frac{1}{2}}$$

$$1 - \frac{2}{y} = -1 \Rightarrow 1 + 1 = \frac{2}{y} \Rightarrow 2 = \frac{2}{y} \Rightarrow y = \frac{2}{2} \Rightarrow \boxed{y = 1}$$

$$\therefore z^2 = x^2 + y^2 \Rightarrow z^2 = \left(\frac{1}{2}\right)^2 + 1^2 = \frac{1}{4} + 1 = \frac{5}{4} \Rightarrow z = \pm \sqrt{\frac{5}{4}} = \pm \frac{\sqrt{5}}{2}$$

Hence the stationary points are $\left(\frac{1}{2}, 1, \frac{\sqrt{5}}{2}\right)$ & $\left(\frac{1}{2}, 1, -\frac{\sqrt{5}}{2}\right)$.

$$d = \sqrt{\left(\frac{1}{2}-1\right)^2 + (1-2)^2 + \left(\frac{\sqrt{5}}{2}\right)^2} = \sqrt{\frac{1}{4} + 1 + \frac{5}{4}} = \sqrt{\frac{3}{2} + 1} = \sqrt{\frac{5}{2}}$$

$$d = \sqrt{\left(\frac{1}{2}-1\right)^2 + (1-2)^2 + \left(-\frac{\sqrt{5}}{2}\right)^2} = \sqrt{\frac{5}{2}}$$

Hence the minimum distance is $\sqrt{\frac{5}{2}}$.

(36) Find the maximum volume of the largest rectangular parallelepiped that can be inscribed in an ellipsoid $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$.

Sol: Given $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ — (1)

Volume of parallelepiped = $(2x)(2y)(2z) = 8xyz$

$$F = 8xyz + \lambda \left(\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} - 1 \right)$$

$$F = 8xyz + \lambda \frac{x^2}{a^2} + \lambda \frac{y^2}{b^2} + \lambda \frac{z^2}{c^2} - \lambda$$

$$F_x = 8yz + \frac{2x\lambda}{a^2} \quad ; \quad F_y = 8xz + \frac{2\lambda y}{b^2} \quad ; \quad F_z = 8xy + \frac{2\lambda z}{c^2}$$

$$F_x = 0 \\ 8yz + \frac{2x\lambda}{a^2} = 0$$

$$8yz = -\frac{2x\lambda}{a^2}$$

$$4yz = -\frac{x\lambda}{a^2}$$

$$4xyz = -\frac{x^2\lambda}{a^2}$$

$$\frac{4xyz}{-\lambda} = \frac{x^2}{a^2} \quad \text{--- (2)}$$

$$F_y = 0 \\ 8xz + \frac{2\lambda y}{b^2} = 0$$

$$8xz = -\frac{2\lambda y}{b^2}$$

$$4xz = -\frac{\lambda y}{b^2}$$

$$4xyz = -\frac{\lambda y^2}{b^2}$$

$$\frac{4xyz}{-\lambda} = \frac{y^2}{b^2} \quad \text{--- (3)}$$

$$F_z = 0 \\ 8xy + \frac{2\lambda z}{c^2} = 0$$

$$8xy = -\frac{2\lambda z}{c^2}$$

$$4xy = -\frac{\lambda z}{c^2}$$

$$4xyz = -\frac{\lambda z^2}{c^2}$$

$$\frac{4xyz}{-\lambda} = \frac{z^2}{c^2} \quad \text{--- (4)}$$

From (2), (3) & (4), $\frac{x^2}{a^2} = \frac{y^2}{b^2} = \frac{z^2}{c^2}$ — (5)

Substituting (5) in (1),

$$\frac{x^2}{a^2} + \frac{x^2}{a^2} + \frac{x^2}{a^2} = 1 \Rightarrow \frac{3x^2}{a^2} = 1 \Rightarrow x^2 = \frac{a^2}{3} \Rightarrow \boxed{x = \frac{a}{\sqrt{3}}}$$

Similarly, $\boxed{y = \frac{b}{\sqrt{3}}}$ & $\boxed{z = \frac{c}{\sqrt{3}}}$

Hence the stationary point is $(\frac{a}{\sqrt{3}}, \frac{b}{\sqrt{3}}, \frac{c}{\sqrt{3}})$.

$$\text{Maximum volume} = 8xyz = 8\left(\frac{a}{\sqrt{3}}\right)\left(\frac{b}{\sqrt{3}}\right)\left(\frac{c}{\sqrt{3}}\right) = \frac{8abc}{3\sqrt{3}}$$

(37) Find the maximum value of $x^m y^n z^p$, when $x+y+z=a$.

Sol: Given $x+y+z=a$ — (*)

$$F = x^m y^n z^p + \lambda(x+y+z-a) = x^m y^n z^p + \lambda x + \lambda y + \lambda z - \lambda a$$

$$F_x = mx^{m-1} y^n z^p + \lambda \quad ; \quad F_y = nx^m y^{n-1} z^p + \lambda \quad ; \quad F_z = px^m y^n z^{p-1} + \lambda$$

$$\begin{aligned} F_x &= 0 \\ mx^{m-1} y^n z^p + \lambda &= 0 \\ mx^{m-1} y^n z^p &= -\lambda \\ \frac{mx^m y^n z^p}{x} &= -\lambda \quad \text{--- (1)} \end{aligned}$$

$$\begin{aligned} F_y &= 0 \\ nx^m y^{n-1} z^p + \lambda &= 0 \\ nx^m y^{n-1} z^p &= -\lambda \\ \frac{nx^m y^n z^p}{y} &= -\lambda \quad \text{--- (2)} \end{aligned}$$

$$\begin{aligned} F_z &= 0 \\ px^m y^n z^{p-1} + \lambda &= 0 \\ px^m y^n z^{p-1} &= -\lambda \\ \frac{px^m y^n z^p}{z} &= -\lambda \quad \text{--- (3)} \end{aligned}$$

From (1), (2) & (3),

$$\frac{mx^m y^n z^p}{x} = \frac{nx^m y^n z^p}{y} = \frac{px^m y^n z^p}{z}$$

Dividing by $x^m y^n z^p$, we get

$$\frac{m}{x} = \frac{n}{y} = \frac{p}{z}$$

$$\frac{m}{x} = \frac{p}{z}$$

$$\Rightarrow \boxed{x = \frac{mz}{p}} \quad \text{--- (4)}$$

$$\frac{n}{y} = \frac{p}{z}$$

$$\Rightarrow \boxed{y = \frac{nz}{p}} \quad \text{--- (5)}$$

∴ (*) becomes,

$$\frac{mz}{p} + \frac{nz}{p} + z = a \Rightarrow z\left(\frac{m}{p} + \frac{n}{p} + 1\right) = a$$

$$\Rightarrow z\left(\frac{m+n+p}{p}\right) = a \Rightarrow z = \frac{ap}{m+n+p} \quad \text{--- (6)}$$

Substituting (6) in (4) & (5),

$$x = \frac{map}{p(m+n+p)} = \frac{ma}{m+n+p}$$

$$y = \frac{nap}{p(m+n+p)} = \frac{an}{m+n+p}$$

Hence the stationary point is $(\frac{am}{m+n+p}, \frac{an}{m+n+p}, \frac{ap}{m+n+p})$.

$$\begin{aligned} \text{Maximum value of } x^m y^n z^p &= \left(\frac{am}{m+n+p}\right)^m \left(\frac{an}{m+n+p}\right)^n \left(\frac{ap}{m+n+p}\right)^p \\ &= \frac{a^m m^m}{(m+n+p)^m} \cdot \frac{a^n n^n}{(m+n+p)^n} \cdot \frac{a^p p^p}{(m+n+p)^p} \\ &= \frac{a^{m+n+p} m^m n^n p^p}{(m+n+p)^{m+n+p}} \end{aligned}$$

A.w
① Find the minimum values of $x^2 y z^3$ subject to the condition

$$2x + y + 3z = a.$$

② Find the maximum value of $400xy^2z^2$ subject to the condition

$$x^2 + y^2 + z^2 = 1.$$

③ Find the dimensions of the rectangular box without top of maximum capacity with surface area 432 square metres.

④ A rectangular box open at the top, is to have a volume of 32 cc. Find the dimensions of the box, that requires the least material for its construction.

INTEGRAL CALCULUS

1

(A0) Fundamental theorem of calculus:

Suppose f is continuous on $[a, b]$.

(i) If $g(x) = \int_a^x f(t) dt$, then $g'(x) = f(x)$.

(ii) $\int_a^b f(x) dx = F(b) - F(a)$, where F is any anti-derivative of f , that is $F' = f$.

(A0) ① Find the derivative of $G(x) = \int_x^1 \cos \sqrt{t} dt$.

Sol: Given $G(x) = \int_x^1 \cos \sqrt{t} dt = -\int_1^x \cos \sqrt{t} dt$

Here $f(t) = \cos \sqrt{t}$ is continuous.

$\therefore G'(x) = -\cos \sqrt{x}$

(A0) ② Evaluate $\int_0^3 (x^3 - 6x) dx$ by using Riemann sum with n sub intervals.

Sol: Take n sub intervals, we have $\Delta x = \frac{b-a}{n} = \frac{3-0}{n} = \frac{3}{n}$

$x_0 = 0, x_1 = \frac{3}{n}, x_2 = \frac{6}{n}, x_3 = \frac{9}{n}, \dots, x_i = \frac{3i}{n}$. Here $f(x) = x^3 - 6x$

$\therefore \int_0^3 (x^3 - 6x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(\frac{3i}{n}\right) \left(\frac{3}{n}\right)$

$= \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left(\left(\frac{3i}{n}\right)^3 - 6\left(\frac{3i}{n}\right) \right)$

$= \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left(\frac{27i^3}{n^3} - \frac{18i}{n} \right)$

$= \lim_{n \rightarrow \infty} \frac{81}{n^4} \sum_{i=1}^n i^3 - \lim_{n \rightarrow \infty} \frac{54}{n^2} \sum_{i=1}^n i$

$= \lim_{n \rightarrow \infty} \frac{81}{n^4} \left[\frac{n(n+1)}{2} \right]^2 - \lim_{n \rightarrow \infty} \frac{54}{n^2} \left[\frac{n(n+1)}{2} \right]$

$= \lim_{n \rightarrow \infty} \frac{81}{n^4} \left[\frac{n^2(1+1/n)}{2} \right]^2 - \lim_{n \rightarrow \infty} \frac{54}{n^2} \left[\frac{n^2(1+1/n)}{2} \right]$

$= \lim_{n \rightarrow \infty} \frac{81}{n^4} \times n^4 \frac{(1+1/n)^2}{4} - \lim_{n \rightarrow \infty} \frac{54}{n^2} \times n^2 \frac{(1+1/n)}{2}$

$= \lim_{n \rightarrow \infty} \frac{81}{4} (1+1/n)^2 - \lim_{n \rightarrow \infty} 27(1+1/n) = \frac{81}{4} - 27 = -\frac{27}{4}$

Note:

$$\textcircled{1} \sum_{i=1}^n i = \frac{n(n+1)}{2}$$

$$\textcircled{2} \sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\textcircled{3} \sum_{i=1}^n i^3 = \left(\frac{n(n+1)}{2}\right)^2$$

(H.W) $\textcircled{1}$ Evaluate $\int_0^3 (x^2 - 2x) dx$ by using Riemann sum with n sub intervals.

(Au) $\textcircled{3}$ What is wrong with the equation $\int_{-1}^2 \frac{4}{x^3} dx = \left[\frac{-2}{x^2} \right]_{-1}^2 = \frac{3}{2}$?

Sol: Here $f(x) = \frac{4}{x^3}$ is not continuous in the interval $[-1, 2]$.

Since $f(x) = \frac{4}{x^3}$ is discontinuous at $x=0$.

$\therefore \int_{-1}^2 \frac{4}{x^3} dx$ doesn't exist.

Formulae:

$$\textcircled{1} \int x^n dx = \frac{x^{n+1}}{n+1} + c, \quad n \neq -1$$

$$\textcircled{2} \int \frac{1}{x} dx = \log x + c$$

$$\textcircled{3} \int e^x dx = e^x + c$$

$$\textcircled{4} \int e^{2x} = \frac{e^{2x}}{2} + c$$

$$\textcircled{5} \int dx = x + c$$

$$\textcircled{6} \int a dx = ax + c, \text{ where } a \text{ is a constant.}$$

$\textcircled{4}$ Evaluate the following:

$$(i) \int \left(\frac{6}{x^2} + \sqrt{x} + x^{3/2} + \frac{5}{x} + 1 \right) dx$$

$$\text{Sol: } \int \left(\frac{6}{x^2} + \sqrt{x} + x^{3/2} + \frac{5}{x} + 1 \right) dx = \int \left(6x^{-2} + x^{1/2} + x^{3/2} + \frac{5}{x} + 1 \right) dx$$

$$= 6x^{\frac{-2+1}{-2+1}} + \frac{x^{\frac{1/2+1}{1/2+1}}}{\frac{1/2+1}{1/2+1}} + \frac{x^{\frac{3/2+1}{3/2+1}}}{\frac{3/2+1}{3/2+1}} + 5 \log x + x + c$$

$$= \frac{6x^{-1}}{-1} + \frac{x^{3/2}}{3/2} + \frac{x^{5/2}}{5/2} + 5 \log x + x + c$$

$$= -\frac{6}{x} + \frac{2}{3}x^{3/2} + \frac{2}{5}x^{5/2} + 5 \log x + x + c$$

$$(ii) \int \frac{x^2 + 3x - 5}{\sqrt{x}} dx$$

$$\text{Sol: } \int \frac{x^2 + 3x - 5}{\sqrt{x}} dx = \int x^{-1/2} (x^2 + 3x - 5) dx$$

$$\begin{aligned}
&= \int (x^{-1/2} x^2 + 3x^{-1/2} x - 5x^{-1/2}) dx \\
&= \int (x^{-1/2+2} + 3x^{-1/2+1} - 5x^{-1/2}) dx = \int (x^{3/2} + 3x^{1/2} - 5x^{-1/2}) dx \\
&= \frac{x^{3/2+1}}{3/2+1} + 3 \frac{x^{1/2+1}}{1/2+1} - 5 \frac{x^{-1/2+1}}{-1/2+1} + c \\
&= \frac{x^{5/2}}{5/2} + 3 \frac{x^{3/2}}{3/2} - 5 \frac{x^{1/2}}{1/2} + c = \frac{2}{5} x^{5/2} + 2x^{3/2} - 10\sqrt{x} + c
\end{aligned}$$

(iii) $\int (e^{2x} + 3x - 7) dx$

Sol: $\int (e^{2x} + 3x - 7) dx = \frac{e^{2x}}{2} + 3 \frac{x^2}{2} - 7x + c$

(iv) $\int (e^{\log x} + 2) dx$

Sol: $\int (e^{\log x} + 2) dx = \int (x + 2) dx = \frac{x^2}{2} + 2x + c$

(v) $\int x^2(1-x)^2 dx$

Sol: $\int x^2(1-x)^2 dx = \int x^2(1+x^2-2x) dx$
 $= \int (x^2 + x^4 - 2x^3) dx$
 $= \frac{x^3}{3} + \frac{x^5}{5} - \frac{2x^4}{4} + c$
 $= \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^4}{2} + c$

H.W
 (1) Evaluate the following:

(i) $\int (x^4 - \frac{1}{2}x^3 + \frac{1}{4}x - 2) dx$ (ii) $\int \frac{x^3 - 2\sqrt{x}}{x} dx$

(iii) $\int (x^{2/5} - x^{-3/5})^2 dx$ (iv) $\int (e^x + x^2 + 8) dx$

Q5 If f is continuous & $\int_0^4 f(x) dx = 10$, find $\int_0^2 f(2x) dx$.

Sol: Take $2x = t$
 $2dx = dt \Rightarrow dx = \frac{dt}{2}$ When $x=0 \Rightarrow t=0$
 $x=2 \Rightarrow t=2(2)=4$

$\therefore \int_0^2 f(2x) dx = \int_0^4 f(t) \frac{dt}{2} = \frac{1}{2} \int_0^4 f(t) dt = \frac{1}{2} (10) = 5$

$(\because \int_0^4 f(x) dx = \int_0^4 f(t) dt = 10)$

Formulae:

- ① $\int \sin x dx = -\cos x + c$
- ② $\int \cos x dx = \sin x + c$
- ③ $\int \sec^2 x dx = \tan x + c$
- ④ $\int \operatorname{cosec}^2 x dx = -\cot x + c$
- ⑤ $\int \sec x \tan x dx = \sec x + c$
- ⑥ $\int \operatorname{cosec} x \cot x dx = -\operatorname{cosec} x + c$
- ⑦ $\int \cosh x dx = \sinh x + c$
- ⑧ $\int \sinh x dx = \cosh x + c$
- ⑨ $\int \frac{1}{1+x^2} dx = \tan^{-1} x + c$
- ⑩ $\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + c$
- ⑪ $\int \frac{1}{\sqrt{x^2-1}} dx = \log(x + \sqrt{x^2-1}) + c$
- ⑫ $\int \frac{1}{\sqrt{x^2+1}} dx = \log(x + \sqrt{x^2+1}) + c$
- ⑬ $\int \frac{1}{x\sqrt{x^2-1}} dx = \sec^{-1} x + c$
- ⑭ $\int \sin 2x dx = \frac{-\cos 2x}{2} + c$

Ans (b) Evaluate $\int \frac{\tan x}{\sec x + \cos x} dx$.

Sol:
$$\int \frac{\tan x}{\sec x + \cos x} dx = \int \frac{\frac{\sin x}{\cos x}}{\frac{1}{\cos x} + \cos x} dx$$

$$= \int \frac{\frac{\sin x}{\cos x}}{\frac{1 + \cos^2 x}{\cos x}} dx = \int \frac{\sin x}{\cos x} \times \frac{\cos x}{1 + \cos^2 x} dx$$

$$= \int \frac{\sin x}{1 + \cos^2 x} dx$$

$$= \int \frac{-dt}{1+t^2} = -\int \frac{dt}{1+t^2}$$

$$= -\tan^{-1} t + c = -\tan^{-1}(\cos x) + c$$

Put $\cos x = t$
 $-\sin x dx = dt$
 $\sin x dx = -dt$

⑦ Evaluate the following:

(i) $\int \frac{1}{1+\sin x} dx$

Sol:
$$\int \frac{1}{1+\sin x} dx = \int \frac{1}{1+\sin x} \times \frac{1-\sin x}{1-\sin x} dx$$

$$= \int \frac{1-\sin x}{1-\sin^2 x} dx = \int \frac{1-\sin x}{\cos^2 x} dx$$

$$= \int \left(\frac{1}{\cos^2 x} - \frac{\sin x}{\cos^2 x} \right) dx = \int (\sec^2 x - \tan x \sec x) dx$$

$$= \tan x - \sec x + c$$

(ii) $\int \frac{\cos^2 x}{1 - \sin x} dx$

Sol: $\int \frac{\cos^2 x}{1 - \sin x} dx = \int \frac{1 - \sin^2 x}{1 - \sin x} dx$

$$= \int \frac{(1 + \sin x)(1 - \sin x)}{1 - \sin x} dx = \int (1 + \sin x) dx$$

$$= x - \cos x + C$$

(iii) $\int (\tan x - 2 \cot x)^2 dx$

Sol: $\int (\tan x - 2 \cot x)^2 dx = \int (\tan^2 x + 4 \cot^2 x - 4 \tan x \cot x) dx$

$$= \int \left(\sec^2 x - 1 + 4(\operatorname{cosec}^2 x - 1) - 4 \tan x \frac{1}{\tan x} \right) dx$$

$$= \int (\sec^2 x - 1 + 4 \operatorname{cosec}^2 x - 4 - 4) dx$$

$$= \int (\sec^2 x + 4 \operatorname{cosec}^2 x - 9) dx$$

$$= \tan x + 4(-\cot x) - 9x + C$$

$$= \tan x - 4 \cot x - 9x + C$$

(H.w) ① Evaluate the following:

(i) $\int \frac{\sin^2 x}{1 + \cos x} dx$ (ii) $\int \frac{1}{1 - \cos x} dx$ (iii) $\int \left(\frac{3}{\sqrt{1-x^2}} + e^x + 8 \right) dx$

⑧ Evaluate the following:

(i) $\int_1^4 (x^2 + 2x - 5) dx$

Sol: $\int_1^4 (x^2 + 2x - 5) dx = \left[\frac{x^3}{3} + \frac{2x^2}{2} - 5x \right]_1^4 = \left[\frac{x^3}{3} + x^2 - 5x \right]_1^4$

$$= \left[\frac{64}{3} + 16 - 20 - \left(\frac{1}{3} + 1 - 5 \right) \right]$$

$$= \frac{64}{3} + 16 - 20 - \frac{1}{3} - 1 + 5 = 21$$

(ii) $\int_{-1}^1 (2 - |x|) dx$

Sol: $\int_{-1}^1 (2 - |x|) dx$

Here $f(x) = 2 - |x|$ is an even function.

$$\int_{-a}^a f(x) dx = \begin{cases} 2 \int_0^a f(x) dx & \text{if } f(x) \text{ is even} \\ 0 & \text{if } f(x) \text{ is odd} \end{cases}$$

$$f(x) = 2 - |x|$$

$$f(-x) = 2 - |-x| = 2 - |x| = f(x)$$

$$\int_{-1}^1 (2-1 \times 1) dx = 2 \int_0^1 (2-x) dx = 2 \left[2x - \frac{x^2}{2} \right]_0^1$$

$$= 2 \left[2 - \frac{1}{2} \right] = 2 \times \frac{3}{2} = 3$$

(A0) (9) Evaluate $\int_0^{\pi/2} \frac{1}{1+\tan x} dx$. (or) $\int_0^{\pi/2} \frac{\cos x}{\sin x + \cos x} dx$

Sol: Let $I = \int_0^{\pi/2} \frac{\cos x}{\sin x + \cos x} dx$ — (1)

$$= \int_0^{\pi/2} \frac{\cos(\frac{\pi}{2}-x)}{\sin(\frac{\pi}{2}-x) + \cos(\frac{\pi}{2}-x)} dx$$

($\because \int_0^a f(x) dx = \int_0^a f(a-x) dx$)

$$\therefore I = \int_0^{\pi/2} \frac{\sin x}{\cos x + \sin x} dx$$
 — (2)

(1) + (2) $\Rightarrow 2I = \int_0^{\pi/2} \frac{\cos x}{\sin x + \cos x} dx + \int_0^{\pi/2} \frac{\sin x}{\cos x + \sin x} dx$

$$= \int_0^{\pi/2} \frac{\sin x + \cos x}{\sin x + \cos x} dx = \int_0^{\pi/2} dx = (x)_0^{\pi/2} = \frac{\pi}{2} - 0 = \frac{\pi}{2}$$

$$\therefore I = \frac{\pi}{2 \times 2} = \frac{\pi}{4}$$

$$\therefore \int_0^{\pi/2} \frac{1}{1+\tan x} dx = \frac{\pi}{4}$$

(H.W) (1) Evaluate: (i) $\int_0^1 (4+3x^2) dx$ (ii) $\int_2^1 \left(1 + \frac{z}{2}\right) dz$

(2) Evaluate: (i) $\int_0^{\pi/2} \frac{\sin x}{\sin x + \cos x} dx$ (ii) $\int_0^{\pi/2} \frac{\sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$

(10) Evaluate $\int_0^{\pi/2} \log(\tan x) dx$.

Sol: Let $I = \int_0^{\pi/2} \log(\tan x) dx$ — (1)

$$I = \int_0^{\pi/2} \log(\tan(\frac{\pi}{2}-x)) dx = \int_0^{\pi/2} \log(\cot x) dx$$
 — (2)

(1) + (2) $\Rightarrow 2I = \int_0^{\pi/2} \log(\tan x) dx + \int_0^{\pi/2} \log(\cot x) dx$

$$= \int_0^{\pi/2} (\log(\tan x) + \log(\cot x)) dx = \int_0^{\pi/2} \log(\tan x \cdot \cot x) dx$$

$$= \int_0^{\pi/2} \log 1 \, dx = \int_0^{\pi/2} 0 \, dx = 0$$

$$\therefore I = 0 \Rightarrow \int_0^{\pi/2} \log(\tan x) \, dx = 0$$

Substitution rule:

⑪ Evaluate: $\int (x+1)\sqrt{2x+x^2} \, dx$

Sol: Put $u = 2x+x^2$
 $du = (2+2x) \, dx = 2(1+x) \, dx \Rightarrow (x+1) \, dx = \frac{du}{2}$

$$\begin{aligned} \therefore \int (x+1)\sqrt{2x+x^2} \, dx &= \int \sqrt{u} \frac{du}{2} = \frac{1}{2} \int \sqrt{u} \, du = \frac{1}{2} \int u^{1/2} \, du \\ &= \frac{1}{2} \left[\frac{u^{1/2+1}}{1/2+1} \right] + c = \frac{1}{2} \left[\frac{u^{3/2}}{3/2} \right] + c \\ &= \frac{1}{2} \times \frac{2}{3} u^{3/2} + c = \frac{1}{3} (2x+x^2)^{3/2} + c \end{aligned}$$

⑫ Evaluate: $\int \frac{x^2}{\sqrt{x+5}} \, dx$

$u^2 = x+5 \Rightarrow u = \sqrt{x+5} = (x+5)^{1/2}$

Sol: Put $u^2 = x+5 \Rightarrow x = u^2 - 5$
 $2u \, du = dx$

$$\begin{aligned} \therefore \int \frac{x^2}{\sqrt{x+5}} \, dx &= \int \frac{(u^2-5)^2}{\sqrt{u^2}} 2u \, du = \int \frac{(u^2-5)^2}{u} 2u \, du = 2 \int (u^2-5)^2 \, du \\ &= 2 \int (u^4 + 25 - 10u^2) \, du = 2 \left(\frac{u^5}{5} + 25u - 10 \frac{u^3}{3} \right) + c \\ &= 2 \left(\frac{(x+5)^{5/2}}{5} + 25\sqrt{x+5} - \frac{10(x+5)^{3/2}}{3} \right) + c \end{aligned}$$

⑬ Evaluate: $\int_1^e \frac{\log x}{x} \, dx$

Sol: Put $u = \log x$
 $du = \frac{1}{x} \, dx$

when $x=1 \Rightarrow u = \log 1 = 0$
 $x=e \Rightarrow u = \log e = 1$

$$\therefore \int_1^e \frac{\log x}{x} \, dx = \int_0^1 u \, du = \left(\frac{u^2}{2} \right)_0^1 = \frac{1}{2}$$

⑭ Evaluate: $\int \frac{\sec^2(\log x)}{x} \, dx$

Sol: Put $u = \log x \Rightarrow du = \frac{1}{x} \, dx$

$$\int \frac{\sec^2(\log x)}{x} dx = \int \sec^2 u du = \tan u + c = \tan(\log x) + c$$

15) Evaluate: $\int_1^2 \frac{e^{1/x}}{x^2} dx$

Sol: Put $u = e^{1/x}$

$$du = e^{1/x} \cdot \left(-\frac{1}{x^2}\right) dx \Rightarrow \frac{dx}{x^2} = \frac{-du}{e^{1/x}} = \frac{-du}{u}$$

When $x=1 \Rightarrow u=e$

$x=2 \Rightarrow u = e^{1/2} = \sqrt{e}$

$$\therefore \int_1^2 \frac{e^{1/x}}{x^2} dx = \int_e^{\sqrt{e}} u \left(\frac{-du}{u}\right) = - \int_e^{\sqrt{e}} du = -(u)_e^{\sqrt{e}} = -(\sqrt{e} - e) = e - \sqrt{e}$$

16) Evaluate: $\int e^{\tan^{-1} x} \left(\frac{1+x+x^2}{1+x^2}\right) dx$.

Sol: Put $u = \tan^{-1} x \Rightarrow x = \tan u$

$$du = \frac{1}{1+x^2} dx$$

$$1+x+x^2 = 1 + \tan u + \tan^2 u = \tan u + \sec^2 u$$

$$\therefore \int e^{\tan^{-1} x} \left(\frac{1+x+x^2}{1+x^2}\right) dx = \int e^u (\tan u + \sec^2 u) du$$

Put $t = e^u \tan u$

$$dt = (e^u \sec^2 u + \tan u e^u) du = e^u (\tan u + \sec^2 u) du$$

$$\therefore \int e^{\tan^{-1} x} \left(\frac{1+x+x^2}{1+x^2}\right) dx = \int dt = t + c$$

$$= e^u \tan u + c = e^{\tan^{-1} x} \tan(\tan^{-1} x) + c$$

$$= x e^{\tan^{-1} x} + c$$

H.w 1) Evaluate the following:

(i) $\int \cos^3 \theta \sin \theta d\theta$

(ii) $\int \sec^2 \theta \tan^2 \theta d\theta$

(iii) $\int \frac{e^x}{e^x + 1} dx$

(iv) $\int_0^1 \frac{e^x + 1}{e^x + x} dx$

(v) $\int \frac{(\log x)^2}{x} dx$

(vi) $\int \frac{(1+\sqrt{x})^{1/3}}{\sqrt{x}} dx$

Integration by parts:

①7 Evaluate: $\int x \cos 5x dx$

$\int u dv = uv - \int v du$

Sol: Let $u = x$, $dv = \cos 5x dx$
 $du = dx$, $\int dv = \int \cos 5x dx$

$v = \frac{\sin 5x}{5}$

$\therefore \int x \cos 5x dx = x \frac{\sin 5x}{5} - \int \frac{\sin 5x}{5} dx$
 $= \frac{x}{5} \sin 5x - \frac{1}{5} \left(\frac{-\cos 5x}{5} \right) + C$
 $= \frac{x}{5} \sin 5x + \frac{1}{25} \cos 5x + C$

①8 Evaluate: $\int x^5 e^x dx$

Bernoulli's formula:
 $\int u dv = uv - u'v_1 + u''v_2 - \dots$

Sol: $u = x^5$ $dv = e^x dx$
 $u' = 5x^4$ $v = e^x$
 $u'' = 20x^3$ $v_1 = e^x$
 $u''' = 60x^2$ $v_2 = e^x$
 $u^{IV} = 120x$ $v_3 = e^x$
 $u^V = 120$ $v_4 = e^x$
 $v_5 = e^x$

$\therefore \int x^5 e^x dx = x^5 e^x - 5x^4 e^x + 20x^3 e^x - 60x^2 e^x + 120x e^x - 120 e^x + C$

(Au) ①9 Using integration by parts, evaluate $\int \frac{(\ln x)^2}{x^2} dx$.

$\ln x = \log x$

Sol: $u = (\log x)^2$ $dv = \frac{dx}{x^2} = x^{-2} dx$
 $du = 2 \log x \cdot \frac{1}{x} dx$ $v = \frac{x^{-2+1}}{-2+1} = \frac{x^{-1}}{-1} = -\frac{1}{x}$

$\int u dv = uv - \int v du$

$\therefore \int \frac{(\ln x)^2}{x^2} dx = \int \frac{(\log x)^2}{x^2} dx$
 $= (\log x)^2 \cdot \frac{-1}{x} - \int \frac{-1}{x} \cdot 2 \log x \cdot \frac{1}{x} dx$
 $= -\frac{1}{x} (\log x)^2 + 2 \int \frac{\log x}{x^2} dx$

$u = \log x$ $dv = \frac{dx}{x^2} = x^{-2} dx$
 $du = \frac{1}{x} dx$ $v = -\frac{1}{x}$

$$\int \frac{(\ln x)^2}{x^2} dx = -\frac{1}{x} (\log x)^2 + 2 \left[\log x \cdot \frac{-1}{x} - \int \frac{-1}{x} \cdot \frac{1}{x} dx \right]$$

$$= -\frac{1}{x} (\log x)^2 - \frac{2}{x} \log x + 2 \int \frac{dx}{x^2}$$

$$= -\frac{1}{x} (\log x)^2 - \frac{2}{x} \log x + 2 \int x^{-2} dx$$

$$= -\frac{1}{x} (\log x)^2 - \frac{2}{x} \log x + 2 \left(\frac{x^{-2+1}}{-2+1} \right) + c$$

$$= -\frac{1}{x} (\log x)^2 - \frac{2}{x} \log x + 2 \left(\frac{x^{-1}}{-1} \right) + c$$

$$= -\frac{1}{x} (\log x)^2 - \frac{2}{x} \log x - \frac{2}{x} + c$$

20 Evaluate $\int e^{ax} \cos bx dx$ using integration by parts.

Sol:

$u = e^{ax}$ $du = e^{ax} \cdot a dx$	$dv = \cos bx dx$ $v = \frac{\sin bx}{b}$
--	--

$\int u dv = uv - \int v du$

$$\text{Let } \int = \int e^{ax} \cos bx dx = e^{ax} \frac{\sin bx}{b} - \int \frac{\sin bx}{b} e^{ax} a dx$$

$$= \frac{1}{b} e^{ax} \sin bx - \frac{a}{b} \int e^{ax} \sin bx dx$$

$u = e^{ax}$ $du = e^{ax} a dx$	$dv = \sin bx dx$ $v = -\frac{\cos bx}{b}$
------------------------------------	---

$$\therefore \int = \frac{1}{b} e^{ax} \sin bx - \frac{a}{b} \left[-e^{ax} \frac{\cos bx}{b} - \int -\frac{\cos bx}{b} e^{ax} a dx \right]$$

$$\int = \frac{1}{b} e^{ax} \sin bx + \frac{a}{b^2} e^{ax} \cos bx - \frac{a^2}{b^2} \int e^{ax} \cos bx dx$$

$$\int = \frac{1}{b} e^{ax} \sin bx + \frac{a}{b^2} e^{ax} \cos bx - \frac{a^2}{b^2} \int$$

$$\int + \frac{a^2}{b^2} \int = \frac{1}{b} e^{ax} \sin bx + \frac{a}{b^2} e^{ax} \cos bx$$

$$\int \left(1 + \frac{a^2}{b^2} \right) = \frac{1}{b} e^{ax} \sin bx + \frac{a}{b^2} e^{ax} \cos bx$$

$$\int = \frac{b^2}{a^2 + b^2} \left(\frac{1}{b} e^{ax} \sin bx + \frac{a}{b^2} e^{ax} \cos bx \right)$$

$$\therefore \int = \frac{1}{a^2 + b^2} \left(b e^{ax} \sin bx + a e^{ax} \cos bx \right) + c$$

(A.V.) (21) Evaluate $\int e^x \sin x dx$ by using integration by parts.

Sol:

$u = e^x$	$dv = \sin x dx$
$du = e^x dx$	$v = -\cos x$

$$\int u dv = uv - \int v du$$

$$\begin{aligned} \text{Let } I &= \int e^x \sin x dx = e^x(-\cos x) - \int -\cos x e^x dx \\ &= -e^x \cos x + \int e^x \cos x dx \end{aligned}$$

$u = e^x$	$dv = \cos x dx$
$du = e^x dx$	$v = \sin x$

$$\begin{aligned} \therefore I &= -e^x \cos x + \left[e^x \sin x - \int \sin x e^x dx \right] \\ &= -e^x \cos x + e^x \sin x - \int e^x \sin x dx \end{aligned}$$

$$\therefore I = -e^x \cos x + e^x \sin x - I$$

$$2I = -e^x \cos x + e^x \sin x \Rightarrow I = \frac{1}{2} e^x (\sin x - \cos x) + c$$

(H.w) (1) Evaluate $\int e^{ax} \sin bx dx$ using integration by parts.

(2) Evaluate $\int e^x \cos x dx$ using integration by parts.

Reduction formula:

(22) Establish a reduction formula for $I_n = \int \sin^n x dx$. Hence find $\int_0^{\pi/2} \sin^n x dx$.

Sol: Given $I_n = \int \sin^n x dx$ — (1)

$$= \int \sin^{n-1} x \sin x dx$$

$u = \sin^{n-1} x$	$dv = \sin x dx$
$du = (n-1) \sin^{n-2} x \cos x dx$	$v = -\cos x$

$$\int u dv = uv - \int v du$$

$$\therefore I_n = \sin^{n-1} x (-\cos x) - \int (-\cos x)(n-1) \sin^{n-2} x \cos x dx$$

$$= -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x \cos^2 x dx$$

$$= -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x (1 - \sin^2 x) dx \quad (\because \sin^2 x + \cos^2 x = 1)$$

$$= -\sin^{n-1} x \cos x + (n-1) \int (\sin^{n-2} x - \sin^n x) dx$$

$$= -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x dx - (n-1) \int \sin^n x dx$$

$$\therefore I_n = -\sin^{n-1}x \cos x + (n-1)I_{n-2} - (n-1)I_n \quad (\because \text{by } \textcircled{1})$$

$$I_n + (n-1)I_n = -\sin^{n-1}x \cos x + (n-1)I_{n-2}$$

$$I_n(1+n-1) = -\sin^{n-1}x \cos x + (n-1)I_{n-2}$$

$$nI_n = -\sin^{n-1}x \cos x + (n-1)I_{n-2}$$

$$\therefore I_n = -\frac{1}{n} \sin^{n-1}x \cos x + \frac{n-1}{n} I_{n-2}$$

$$\int \sin^n x dx = -\frac{1}{n} \sin^{n-1}x \cos x + \frac{n-1}{n} \int \sin^{n-2}x dx$$

$$I_0 = \int \sin^0 x dx = \int dx = x + C$$

$$I_1 = \int \sin x dx = -\cos x + C$$

Now consider, $I_n = \int_0^{\pi/2} \sin^n x dx$

$$I_n = \left(-\frac{1}{n} \sin^{n-1}x \cos x \right)_0^{\pi/2} + \frac{n-1}{n} \int_0^{\pi/2} \sin^{n-2}x dx$$

$$= (0+0) + \frac{n-1}{n} \int_0^{\pi/2} \sin^{n-2}x dx = \frac{n-1}{n} I_{n-2}$$

$$\therefore I_n = \frac{n-1}{n} I_{n-2} \quad \text{--- } \textcircled{2}$$

$$I_{n-2} = \frac{n-2-1}{n-2} I_{n-2-2} = \frac{n-3}{n-2} I_{n-4}$$

$$I_{n-4} = \frac{n-4-1}{n-4} I_{n-4-2} = \frac{n-5}{n-4} I_{n-6}$$

⋮

$$\therefore I_n = \begin{cases} \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdot \frac{n-5}{n-4} \cdots \frac{1}{2} I_0 & \text{if } n \text{ is even} \\ \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdot \frac{n-5}{n-4} \cdots \frac{2}{3} I_1 & \text{if } n \text{ is odd} \end{cases}$$

$$I_0 = \int_0^{\pi/2} \sin^0 x dx = \int_0^{\pi/2} dx = (x)_0^{\pi/2} = \frac{\pi}{2}$$

$$I_1 = \int_0^{\pi/2} \sin x dx = (-\cos x)_0^{\pi/2} = -\cos \frac{\pi}{2} + \cos 0 = -0 + 1 = 1$$

$$\therefore I_n = \begin{cases} \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdot \frac{n-5}{n-4} \cdots \frac{1}{2} \cdot \frac{\pi}{2} & \text{if } n \text{ is even} \\ \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdot \frac{n-5}{n-4} \cdots \frac{2}{3} \cdot 1 & \text{if } n \text{ is odd} \end{cases}$$

$$\textcircled{2} \Rightarrow I_2 = \frac{2-1}{2} I_0$$

$$I_2 = \frac{1}{2} I_0$$

$$\textcircled{2} \Rightarrow I_3 = \frac{3-1}{3} I_{3-2}$$

$$I_3 = \frac{2}{3} I_1$$

(23) Establish a reduction formula for $I_n = \int \cos^n x dx$. Hence find $\int_0^{\pi/2} \cos^n x dx$.

Sol: Given $I_n = \int \cos^n x dx$ — (1)
 $= \int \cos^{n-1} x \cos x dx$

$\int u dv = uv - \int v du$

$u = \cos^{n-1} x$ $du = (n-1) \cos^{n-2} x (-\sin x) dx$	$dv = \cos x dx$ $v = \sin x$
--	----------------------------------

$\therefore I_n = \cos^{n-1} x \sin x - \int \sin x (n-1) \cos^{n-2} x (-\sin x) dx$
 $= \cos^{n-1} x \sin x + (n-1) \int \cos^{n-2} x \sin^2 x dx$
 $= \cos^{n-1} x \sin x + (n-1) \int \cos^{n-2} x (1 - \cos^2 x) dx$
 $= \cos^{n-1} x \sin x + (n-1) \int \cos^{n-2} x dx - (n-1) \int \cos^n x dx$

$I_n = \cos^{n-1} x \sin x + (n-1) I_{n-2} - (n-1) I_n$

$I_n + (n-1) I_n = \cos^{n-1} x \sin x + (n-1) I_{n-2}$

$I_n (1+n-1) = \cos^{n-1} x \sin x + (n-1) I_{n-2}$

$\therefore n I_n = \cos^{n-1} x \sin x + (n-1) I_{n-2}$

$I_n = \frac{1}{n} \cos^{n-1} x \sin x + \frac{n-1}{n} I_{n-2}$

$\therefore \int \cos^n x dx = \frac{1}{n} \cos^{n-1} x \sin x + \frac{n-1}{n} \int \cos^{n-2} x dx$

$I_0 = \int \cos^0 x dx = \int dx = x + c$

$I_1 = \int \cos x dx = \sin x + c$

Now consider, $I_n = \int_0^{\pi/2} \cos^n x dx$

$I_n = \left(\frac{1}{n} \cos^{n-1} x \sin x \right)_0^{\pi/2} + \frac{n-1}{n} \int_0^{\pi/2} \cos^{n-2} x dx$

$= 0 + \frac{n-1}{n} I_{n-2}$

$\therefore I_n = \frac{n-1}{n} I_{n-2}$

$I_{n-2} = \frac{n-2-1}{n-2} I_{n-2-2} = \frac{n-3}{n-2} I_{n-4}$

$I_{n-4} = \frac{n-4-1}{n-4} I_{n-4-2} = \frac{n-5}{n-4} I_{n-6}$

⋮

$\therefore I_n = \left\{ \begin{array}{l} \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdot \frac{n-5}{n-4} \dots \frac{1}{2} \cdot I_0 \\ \text{if } n \text{ is even} \\ \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdot \frac{n-5}{n-4} \dots \frac{2}{3} I_1 \\ \text{if } n \text{ is odd} \end{array} \right.$

$$I_0 = \int_0^{\pi/2} \cos^0 x dx = \int_0^{\pi/2} dx = (x)_0^{\pi/2} = \frac{\pi}{2}$$

$$I_1 = \int_0^{\pi/2} \cos x dx = (\sin x)_0^{\pi/2} = \sin \pi/2 - \sin 0 = 1 - 0 = 1$$

$$\therefore I_n = \begin{cases} \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdot \frac{n-5}{n-4} \dots \frac{1}{2} \cdot \frac{\pi}{2} & \text{if } n \text{ is even} \\ \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdot \frac{n-5}{n-4} \dots \frac{2}{3} \cdot 1 & \text{if } n \text{ is odd} \end{cases}$$

- (24) Find the value of (i) $\int_0^{\pi/2} \sin^3 x dx$ (ii) $\int_0^{\pi/2} \sin^4 x dx$ (iii) $\int_0^{\pi/2} \sin^7 x dx$
 (iv) $\int_0^{\pi/2} \sin^8 x dx$ (v) $\int_0^{\pi/2} \sin^{2n} x dx$.

Sol: We know that $\int \sin^n x dx = -\frac{1}{n} \sin^{n-1} x \cos x + \frac{n-1}{n} \int \sin^{n-2} x dx$

(i) $\int \sin^3 x dx = -\frac{1}{3} \sin^2 x \cos x + \frac{3-1}{3} \int \sin x dx$ (Here $n=3$)
 $= -\frac{1}{3} \sin^2 x \cos x + \frac{2}{3} (-\cos x) + c$

$\therefore \int \sin^3 x dx = -\frac{1}{3} \sin^2 x \cos x - \frac{2}{3} \cos x + c$

(ii) $\int \sin^4 x dx = -\frac{1}{4} \sin^3 x \cos x + \frac{3}{4} \int \sin^2 x dx$ (Here $n=4$)

$= -\frac{1}{4} \sin^3 x \cos x + \frac{3}{4} \left[-\frac{1}{2} \sin x \cos x + \frac{1}{2} \int \sin^0 x dx \right]$ (Here $n=2$)

$= -\frac{1}{4} \sin^3 x \cos x - \frac{3}{8} \sin x \cos x + \frac{3}{8} x + c$

$= -\frac{1}{4} \sin^3 x \cos x - \frac{3}{8} \frac{\sin 2x}{2} + \frac{3}{8} x + c$ ($\because 2 \sin x \cos x = \sin 2x$)

$= -\frac{1}{4} \sin^3 x \cos x - \frac{3}{16} \sin 2x + \frac{3}{8} x + c$

(iii) $\int_0^{\pi/2} \sin^7 x dx$ (Here $n=7 \Rightarrow$ odd)

We know that $\int_0^{\pi/2} \sin^n x dx = \begin{cases} \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdot \frac{n-5}{n-4} \dots \frac{1}{2} \cdot \frac{\pi}{2} & \text{if } n \text{ is even} \\ \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdot \frac{n-5}{n-4} \dots \frac{2}{3} \cdot 1 & \text{if } n \text{ is odd} \end{cases}$

$\therefore \int_0^{\pi/2} \sin^7 x dx = \frac{7-1}{7} \cdot \frac{7-3}{7-2} \cdot \frac{7-5}{7-4} \cdot 1 = \frac{6}{7} \times \frac{4}{5} \times \frac{2}{3} \times 1 = \frac{16}{35}$

(iv) $\int_0^{\pi/2} \sin^8 x dx = \frac{7}{8} \times \frac{5}{6} \times \frac{3}{4} \times \frac{1}{2} \times \frac{\pi}{2} = \frac{35}{256} \pi$

(Here $n=8 \rightarrow$ even)

$$(v) \int_0^{\pi/2} \sin^{2n} x dx = \frac{2n-1}{2n} \cdot \frac{2n-3}{2n-2} \cdot \frac{2n-5}{2n-4} \dots \frac{1}{2} \cdot \frac{\pi}{2} \quad (\text{Here } 2n \text{ is even})$$

(10) Evaluate $\int_0^{\pi/2} \cos^5 x dx$.

Sol: We know that $\int_0^{\pi/2} \cos^n x dx = \begin{cases} \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdot \frac{n-5}{n-4} \dots \frac{1}{2} \cdot \frac{\pi}{2} & \text{if } n \text{ is even} \\ \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdot \frac{n-5}{n-4} \dots \frac{2}{3} \cdot 1 & \text{if } n \text{ is odd} \end{cases}$

Here $n=5$. $\therefore n$ is odd.

$$\therefore \int_0^{\pi/2} \cos^5 x dx = \frac{4}{5} \times \frac{2}{3} \times 1 = \frac{8}{15}$$

(H.w) (1) Find the value of (i) $\int_0^{\pi/2} \sin^2 x dx$ (ii) $\int_0^{\pi/2} \sin^5 x dx$ (iii) $\int_0^{\pi/2} \cos^3 x dx$

(iv) $\int_0^{\pi/2} \cos^{10} x dx$

(26) Find the reduction formula for $\int \sec^n x dx$, $n \geq 2$ is an integer.

Sol: Let $I_n = \int \sec^n x dx = \int \sec^{n-2} x \sec^2 x dx$

$u = \sec^{n-2} x$	$dv = \sec^2 x dx$
$du = (n-2) \sec^{n-3} x (\sec x \tan x) dx$	$v = \tan x$

$\int u dv = uv - \int v du$

$$\begin{aligned} \therefore I_n &= \sec^{n-2} x \tan x - \int \tan x (n-2) \sec^{n-3} x \sec x \tan x dx \\ &= \sec^{n-2} x \tan x - (n-2) \int \sec^{n-2} x \tan^2 x dx \\ &= \sec^{n-2} x \tan x - (n-2) \int \sec^{n-2} x (\sec^2 x - 1) dx \\ &= \sec^{n-2} x \tan x - (n-2) \int (\sec^n x - \sec^{n-2} x) dx \\ &= \sec^{n-2} x \tan x - (n-2) \int \sec^n x dx + (n-2) \int \sec^{n-2} x dx \end{aligned}$$

$$\begin{aligned} I_n &= \sec^{n-2} x \tan x - (n-2) I_n + (n-2) I_{n-2} \\ \Rightarrow I_n + (n-2) I_n &= \sec^{n-2} x \tan x + (n-2) I_{n-2} \\ I_n (1+n-2) &= \sec^{n-2} x \tan x + (n-2) I_{n-2} \end{aligned}$$

$I_n = \frac{1}{n-1} \sec^{n-2} x \tan x + \frac{n-2}{n-1} I_{n-2}$

$$I_0 = \int \sec^0 x dx = \int dx = x + c$$

$$I_1 = \int \sec x dx = \log(\sec x + \tan x) + c$$

(27) Find the reduction formula for $\int \operatorname{cosec}^n x dx$, $n \geq 2$ is an integer.

Sol: Let $I_n = \int \operatorname{cosec}^n x dx = \int \operatorname{cosec}^{n-2} x \operatorname{cosec}^2 x dx$

$u = \operatorname{cosec}^{n-2} x$	$dv = \operatorname{cosec}^2 x dx$
$du = (n-2) \operatorname{cosec}^{n-3} x (-\operatorname{cosec} x \cot x) dx$	$v = -\cot x$

$$\begin{aligned} \therefore I_n &= \operatorname{cosec}^{n-2} x (-\cot x) - \int (-\cot x) (n-2) \operatorname{cosec}^{n-3} x (-\operatorname{cosec} x \cot x) dx \\ &= -\operatorname{cosec}^{n-2} x \cot x + (n-2) \int \operatorname{cosec}^{n-2} x \cot^2 x dx \\ &= -\operatorname{cosec}^{n-2} x \cot x + (n-2) \int \operatorname{cosec}^{n-2} x (\operatorname{cosec}^2 x - 1) dx \\ &= -\operatorname{cosec}^{n-2} x \cot x + (n-2) \int \operatorname{cosec}^n x dx + (n-2) \int \operatorname{cosec}^{n-2} x dx \end{aligned}$$

$$I_n = -\operatorname{cosec}^{n-2} x \cot x + (n-2) I_n + (n-2) I_{n-2}$$

$$\therefore I_n + (n-2) I_n = -\operatorname{cosec}^{n-2} x \cot x + (n-2) I_{n-2}$$

$$I_n (1+n-2) = -\operatorname{cosec}^{n-2} x \cot x + (n-2) I_{n-2}$$

$$\therefore I_n = \frac{-1}{n-1} \operatorname{cosec}^{n-2} x \cot x + \frac{n-2}{n-1} I_{n-2}$$

$$I_0 = \int \operatorname{cosec}^0 x dx = \int dx = x + c$$

$$I_1 = \int \operatorname{cosec} x dx = \log(\operatorname{cosec} x - \cot x) + c$$

(28) Find the reduction formula for $\int \tan^n x dx$.

Sol: Let $I_n = \int \tan^n x dx = \int \tan^{n-2} x \tan^2 x dx$

$$\begin{aligned} &= \int \tan^{n-2} x (\sec^2 x - 1) dx \\ &= \int \tan^{n-2} x \sec^2 x dx - \int \tan^{n-2} x dx \\ &= \int u^{n-2} du - \int \tan^{n-2} x dx \\ &= \frac{u^{n-2+1}}{n-2+1} - I_{n-2} = \frac{u^{n-1}}{n-1} - I_{n-2} \end{aligned}$$

Put $u = \tan x$ $du = \sec^2 x dx$
--

$$\therefore I_n = \frac{\tan^{n-1} x}{n-1} - I_{n-2}$$

$$I_0 = \int \tan^0 x dx = \int dx = x + c$$

$$I_1 = \int \tan x dx = \log(\sec x) + c$$

(29) Find the reduction formula for $\int \cot^n x dx$, $n \neq 1$.

Sol: Let $I_n = \int \cot^n x dx$

$$= \int \cot^{n-2} x \cot^2 x dx = \int \cot^{n-2} x (\operatorname{cosec}^2 x - 1) dx$$

$$= \int \cot^{n-2} x \operatorname{cosec}^2 x dx - \int \cot^{n-2} x dx$$

$$= \int u^{n-2} (-du) - I_{n-2}$$

$$= - \left[\frac{u^{n-2+1}}{n-2+1} \right] - I_{n-2}$$

$$= - \frac{u^{n-1}}{n-1} - I_{n-2} = - \frac{\cot^{n-1} x}{n-1} - I_{n-2}$$

$$\therefore I_n = - \frac{\cot^{n-1} x}{n-1} - I_{n-2}$$

$$I_0 = \int \cot^0 x dx = \int dx = x + c$$

$$I_1 = \int \cot x dx = \int \frac{\cos x}{\sin x} dx = \log(\sin x) + c$$

Put $u = \cot x$
 $du = -\operatorname{cosec}^2 x dx$
 $-du = \operatorname{cosec}^2 x dx$

(30) Evaluate: $\int \sin^6 x \cos^3 x dx$.

Sol: Let $I = \int \sin^6 x \cos^3 x dx = \int \sin^6 x \cos^2 x \cos x dx = \int \sin^6 x (1 - \sin^2 x) \cos x dx$

Put $u = \sin x$
 $du = \cos x dx$

$$\therefore I = \int u^6 (1 - u^2) du = \int (u^6 - u^8) du = \left(\frac{u^7}{7} - \frac{u^9}{9} \right) + c$$

$$= \frac{\sin^7 x}{7} - \frac{\sin^9 x}{9} + c$$

(31) Evaluate: $\int \sin^5 x \cos^2 x dx$.

Sol: Let $I = \int \sin^5 x \cos^2 x dx = \int \sin^4 x \cos^2 x \sin x dx$
 $= \int (1 - \cos^2 x)^2 \cos^2 x \sin x dx$

Put $u = \cos x$
 $du = -\sin x dx \Rightarrow -du = \sin x dx$

$$\therefore I = \int (1 - u^2)^2 u^2 (-du) = - \int (1 + u^4 - 2u^2) u^2 du = - \int (u^2 + u^6 - 2u^4) du$$

$$= - \left(\frac{u^3}{3} + \frac{u^7}{7} - \frac{2u^5}{5} \right) + c = - \frac{\cos^3 x}{3} - \frac{\cos^7 x}{7} + \frac{2\cos^5 x}{5} + c$$

32 Evaluate: $\int \cos^2 x \sin 2x dx$.

Sol: Let $I = \int \cos^2 x \sin 2x dx = \int \cos^2 x 2 \sin x \cos x dx$
 $= 2 \int \cos^3 x \sin x dx$ ($\because \sin 2x = 2 \sin x \cos x$)

Put $u = \cos x$

$du = -\sin x dx \Rightarrow \sin x dx = -du$

$\therefore I = 2 \int u^3 (-du) = -2 \int u^3 du = -2 \left(\frac{u^4}{4} \right) + c = -\frac{1}{2} \cos^4 x + c$

33 Evaluate: $\int_0^\pi \sin^2 x \cos^4 x dx$

Sol: Let $I = \int_0^\pi \sin^2 x \cos^4 x dx = \int_0^\pi \left(\frac{1 - \cos 2x}{2} \right) \left(\frac{1 + \cos 2x}{2} \right)^2 dx$
 $= \frac{1}{8} \int_0^\pi (1 - \cos 2x)(1 + \cos^2 2x + 2 \cos 2x) dx$
 $= \frac{1}{8} \int_0^\pi (1 + \cos^2 2x + 2 \cos 2x - \cos 2x - \cos^3 2x - 2 \cos^2 2x) dx$
 $= \frac{1}{8} \int_0^\pi (1 - \cos^2 2x + \cos 2x - \cos^3 2x) dx$ — (1)

$\int \cos^2 2x dx = \int \left(\frac{1 + \cos 4x}{2} \right) dx = \frac{1}{2} \left(x + \frac{\sin 4x}{4} \right)$

$\int \cos^3 2x dx = \int \cos^2 2x \cos 2x dx = \int (1 - \sin^2 2x) \cos 2x dx$

Put $u = \sin 2x$

$du = 2 \cos 2x dx \Rightarrow \cos 2x dx = \frac{du}{2}$

$\therefore \int \cos^3 2x dx = \int (1 - u^2) \frac{du}{2} = \frac{1}{2} \left(u - \frac{u^3}{3} \right) = \frac{1}{2} \left(\sin 2x - \frac{\sin^3 2x}{3} \right)$


$\therefore I = \frac{1}{8} \left[x - \frac{1}{2} x - \frac{1}{8} \sin 4x + \frac{\sin 2x}{2} - \frac{1}{2} \sin 2x + \frac{1}{6} \sin^3 2x \right]_0^\pi$

$= \frac{1}{8} \left[\frac{1}{2} x - \frac{1}{8} \sin 4x + \frac{1}{6} \sin^3 2x \right]_0^\pi$

$= \frac{1}{8} \left[\frac{\pi}{2} \right] = \frac{\pi}{16}$


$\sin n\pi = 0$
$\sin 0 = 0$

Trigonometric substitution:




$$\tan \theta = \frac{x}{a}$$

$$x = a \tan \theta$$



$$\sin \theta = \frac{x}{a}$$

$$x = a \sin \theta$$



$$\cos \theta = \frac{a}{x}$$

$$x = \frac{a}{\cos \theta} = a \sec \theta$$

$$x = a \sec \theta$$

34 Evaluate $\int \frac{x^2}{\sqrt{9-x^2}} dx$.

Sol: Put $x = a \sin \theta$. Here $a = 3$

$$x = 3 \sin \theta$$

$$dx = 3 \cos \theta d\theta$$

$$\therefore \int \frac{x^2}{\sqrt{9-x^2}} dx = \int \frac{(3 \sin \theta)^2}{\sqrt{9-(3 \sin \theta)^2}} 3 \cos \theta d\theta$$

$$= \int \frac{9 \sin^2 \theta}{\sqrt{9-9 \sin^2 \theta}} 3 \cos \theta d\theta = 9 \int \frac{\sin^2 \theta}{3 \sqrt{1-\sin^2 \theta}} 3 \cos \theta d\theta$$

$$= 9 \int \frac{\sin^2 \theta}{\cos \theta} \cos \theta d\theta = 9 \int \sin^2 \theta d\theta = 9 \int \left(\frac{1-\cos 2\theta}{2} \right) d\theta$$


$$= \frac{9}{2} \left[\theta - \frac{\sin 2\theta}{2} \right] + c = \frac{9}{2} \left[\theta - \frac{2 \sin \theta \cos \theta}{2} \right] + c$$

$$= \frac{9}{2} \left[\theta - \sin \theta \cos \theta \right] + c$$

$$= \frac{9}{2} \left[\sin^{-1} \left(\frac{x}{3} \right) - \frac{x}{3} \frac{\sqrt{9-x^2}}{3} \right] + c$$


$$= \frac{9}{2} \sin^{-1} \left(\frac{x}{3} \right) - \frac{1}{2} x \sqrt{9-x^2} + c$$

$$\begin{aligned} x &= a \sin \theta \\ \theta &= \sin^{-1} \left(\frac{x}{a} \right) \\ \theta &= \sin^{-1} \left(\frac{x}{3} \right) \end{aligned}$$



35) Evaluate $\int \sqrt{a^2 - x^2} dx$ by using substitution rule.

Sol: Put $x = a \sin \theta \Rightarrow \theta = \sin^{-1}(x/a)$
 $dx = a \cos \theta d\theta$



$$\begin{aligned} \therefore \int \sqrt{a^2 - x^2} dx &= \int \sqrt{a^2 - (a \sin \theta)^2} a \cos \theta d\theta \\ &= \int \sqrt{a^2 - a^2 \sin^2 \theta} a \cos \theta d\theta \\ &= a^2 \int \sqrt{1 - \sin^2 \theta} \cos \theta d\theta = a^2 \int \cos^2 \theta d\theta \\ &= a^2 \int \left(\frac{1 + \cos 2\theta}{2} \right) d\theta \\ &= \frac{a^2}{2} \left[\theta + \frac{\sin 2\theta}{2} \right] + c \\ &= \frac{a^2}{2} \theta + \frac{a^2}{4} (2 \sin \theta \cos \theta) + c \\ &= \frac{a^2}{2} \theta + \frac{a^2}{2} \sin \theta \cos \theta + c \\ &= \frac{a^2}{2} \sin^{-1} \left(\frac{x}{a} \right) + \frac{a^2}{2} \frac{x}{a} \frac{\sqrt{a^2 - x^2}}{a} + c \\ &= \frac{a^2}{2} \sin^{-1} \left(\frac{x}{a} \right) + \frac{x}{2} \sqrt{a^2 - x^2} + c \end{aligned}$$

36) Evaluate $\int \frac{x}{\sqrt{3 - 2x - x^2}} dx$.

Sol: Consider, $3 - 2x - x^2 = -(x^2 + 2x) + 3$
 $= -(x^2 + 2x + 1 - 1) + 3$
 $= -[(x+1)^2 - 1] + 3$
 $= -(x+1)^2 + 1 + 3 = 4 - (x+1)^2$

Put $u = x+1 \Rightarrow x = u-1$
 $du = dx$


$$\therefore \int \frac{x}{\sqrt{3 - 2x - x^2}} dx = \int \frac{u-1}{\sqrt{4 - u^2}} du$$

Put $u = a \sin \theta$, Here $a = 2$

$$u = 2 \sin \theta \Rightarrow \theta = \sin^{-1} \frac{u}{2}$$

$$du = 2 \cos \theta d\theta$$

$$\begin{aligned} \therefore \int \frac{x}{\sqrt{3-2x-x^2}} dx &= \int \frac{u-1}{\sqrt{4-u^2}} du \\ &= \int \frac{2\sin\theta - 1}{\sqrt{4-4\sin^2\theta}} 2\cos\theta d\theta \\ &= \int \frac{2\sin\theta - 1}{2\cos\theta} 2\cos\theta d\theta = \int (2\sin\theta - 1) d\theta \\ &= 2(-\cos\theta) - \theta + C \\ &= -2 \frac{\sqrt{4-u^2}}{2} - \sin^{-1} \frac{u}{2} + C \\ &= -\sqrt{4-(x+1)^2} - \sin^{-1} \left(\frac{x+1}{2} \right) + C \\ &= -\sqrt{4-x^2-1-2x} - \sin^{-1} \left(\frac{x+1}{2} \right) + C \\ &= -\sqrt{3-x^2-2x} - \sin^{-1} \left(\frac{x+1}{2} \right) + C \end{aligned}$$



(A0) (37) Evaluate $\int_{\sqrt{2}/3}^{2/3} \frac{dx}{x^5 \sqrt{9x^2-1}}$

Sol: $\int_{\sqrt{2}/3}^{2/3} \frac{dx}{x^5 \sqrt{9(x^2-1/9)}} = \frac{1}{3} \int_{\sqrt{2}/3}^{2/3} \frac{dx}{x^5 \sqrt{x^2 - (1/3)^2}} = I \text{ (Say)}$

Put $x = a \sec\theta$

Here $a = 1/3$. $x = 1/3 \sec\theta$

$dx = 1/3 \sec\theta \tan\theta d\theta$

When $x = 2/3 \Rightarrow \frac{2}{3} = \frac{1}{3} \sec\theta$
 $\Rightarrow \frac{2}{3} \times 3 = \frac{1}{\cos\theta} \Rightarrow \cos\theta = \frac{1}{2}$
 $\Rightarrow \theta = \frac{\pi}{3}$

$\sqrt{x^2 - (1/3)^2} = \sqrt{1/9 \sec^2\theta - 1/9} = \frac{1}{3} \sqrt{\sec^2\theta - 1} = \frac{1}{3} \sqrt{\tan^2\theta} = \frac{1}{3} \tan\theta$

$\therefore I = \frac{1}{3} \int_{\pi/4}^{\pi/3} \frac{1/3 \sec\theta \tan\theta}{(1/3)^5 \sec^5\theta \cdot 1/3 \tan\theta} d\theta$

$= \frac{1}{9} \times 3^6 \int_{\pi/4}^{\pi/3} \frac{1}{\sec^4\theta} d\theta$

$= 81 \int_{\pi/4}^{\pi/3} \cos^4\theta d\theta = 81 \int_{\pi/4}^{\pi/3} \left(\frac{1+\cos 2\theta}{2} \right)^2 d\theta$

When $x = \sqrt{2}/3 \Rightarrow \frac{\sqrt{2}}{3} = \frac{1}{3} \sec\theta$
 $\Rightarrow \frac{\sqrt{2}}{3} \times 3 = \frac{1}{\cos\theta} \Rightarrow \cos\theta = \frac{1}{\sqrt{2}}$
 $\Rightarrow \theta = \frac{\pi}{4}$

$$\begin{aligned}
&= \frac{81}{4} \int_{\pi/4}^{\pi/3} (1 + \cos 2\theta)^2 d\theta \\
&= \frac{81}{4} \int_{\pi/4}^{\pi/3} (1 + \cos^2 2\theta + 2\cos 2\theta) d\theta \\
&= \frac{81}{4} \int_{\pi/4}^{\pi/3} \left(1 + \frac{1 + \cos 4\theta}{2} + 2\cos 2\theta\right) d\theta \\
&= \frac{81}{4} \int_{\pi/4}^{\pi/3} \left(1 + \frac{1}{2} + \frac{\cos 4\theta}{2} + 2\cos 2\theta\right) d\theta \\
&= \frac{81}{4} \int_{\pi/4}^{\pi/3} \left(\frac{3}{2} + \frac{\cos 4\theta}{2} + 2\cos 2\theta\right) d\theta \\
&= \frac{81}{4} \left[\frac{3}{2}\theta + \frac{\sin 4\theta}{8} + \frac{2\sin 2\theta}{2} \right]_{\pi/4}^{\pi/3} \\
&= \frac{81}{4} \left[\frac{3}{2}\left(\frac{\pi}{3}\right) + \frac{\sin 4\pi/3}{8} + \sin \frac{2\pi}{3} - \frac{3}{2}\left(\frac{\pi}{4}\right) - \frac{\sin \pi}{8} - \frac{\sin \pi}{2} \right] \\
&= \frac{81}{4} \left[\frac{\pi}{2} - \frac{\sin \pi/3}{8} + \cos \frac{\pi}{6} - \frac{3\pi}{8} - 1 \right] \\
&= \frac{81}{4} \left[\frac{\pi}{2} - \frac{\sqrt{3}}{2 \times 8} + \frac{\sqrt{3}}{2} - \frac{3\pi}{8} - 1 \right] \\
&= \frac{81}{4} \left[\frac{4\pi}{8} - \frac{3\pi}{8} - \frac{\sqrt{3}}{16} + \frac{\sqrt{3}}{2} - 1 \right] = \frac{81}{4} \left[\frac{\pi}{8} - \frac{\sqrt{3}}{16} + \frac{8\sqrt{3}}{16} - 1 \right] \\
&= \frac{81}{4} \left[\frac{\pi}{8} + \frac{7\sqrt{3}}{16} - 1 \right] = \frac{81}{32} \left[\pi + \frac{7\sqrt{3}}{2} - 8 \right]
\end{aligned}$$


38) Evaluate $\int \frac{1}{x^2 \sqrt{x^2+4}} dx$.

Sol: Here $a=2$

Put $x = a \tan \theta \Rightarrow x = 2 \tan \theta$
 $dx = 2 \sec^2 \theta d\theta$

$$\therefore \int \frac{1}{x^2 \sqrt{x^2+4}} dx = \int \frac{1}{4 \tan^2 \theta \sqrt{4 \tan^2 \theta + 4}} \cdot 2 \sec^2 \theta d\theta$$

$$\begin{aligned}
 &= \frac{1}{2} \int \frac{1}{\tan^2 \theta \cdot 2\sqrt{1+\tan^2 \theta}} \sec^2 \theta d\theta \\
 &= \frac{1}{4} \int \frac{1}{\tan^2 \theta \sec \theta} \sec^2 \theta d\theta = \frac{1}{4} \int \frac{\sec \theta}{\tan^2 \theta} d\theta \\
 &= \frac{1}{4} \int \frac{1/\cos \theta}{\frac{\sin^2 \theta}{\cos^2 \theta}} d\theta = \frac{1}{4} \int \frac{1}{\cos \theta} \times \frac{\cos^2 \theta}{\sin^2 \theta} d\theta \\
 &= \frac{1}{4} \int \frac{\cos \theta}{\sin^2 \theta} d\theta = \frac{1}{4} \int \operatorname{cosec} \theta \cot \theta d\theta \\
 &= -\frac{1}{4} \operatorname{cosec} \theta + C \\
 &= -\frac{1}{4} \frac{\sqrt{4+x^2}}{x} + C
 \end{aligned}$$



Integration of rational functions by partial fraction:


(A0) 39 Evaluate $\int_0^{\pi/2} \frac{\sin x \cos x}{\cos^2 x + 3 \cos x + 2} dx$.

Sol: Put $u = \cos x$
 $du = -\sin x dx \Rightarrow \sin x dx = -du$

When $x = \pi/2 \Rightarrow u = \cos \pi/2 = 0$

$x = 0 \Rightarrow u = \cos 0 = 1$

$$\begin{aligned}
 \therefore \int_0^{\pi/2} \frac{\sin x \cos x}{\cos^2 x + 3 \cos x + 2} dx &= \int_1^0 \frac{u(-du)}{u^2 + 3u + 2} \\
 &= - \int_1^0 \frac{u du}{(u+1)(u+2)} = \int_0^1 \frac{u du}{(u+1)(u+2)}
 \end{aligned}$$



Consider, $\frac{u}{(u+1)(u+2)} = \frac{A}{u+1} + \frac{B}{u+2} = \frac{A(u+2) + B(u+1)}{(u+1)(u+2)}$

$\therefore u = A(u+2) + B(u+1)$

Put $u = -2$

Put $u = -1$

$-2 = -B \Rightarrow \boxed{B = 2}$

$\boxed{-1 = A}$

$\therefore \frac{u}{(u+1)(u+2)} = \frac{-1}{u+1} + \frac{2}{u+2}$

$\therefore \int = \int_0^1 \frac{u du}{(u+1)(u+2)} = \int_0^1 \left(\frac{-1}{u+1} + \frac{2}{u+2} \right) du$

$$\begin{aligned}
&= \left(-\log(u+1) + 2\log(u+2) \right)'_0 \\
&= -\log 2 + 2\log 3 + \log 1 - 2\log 2 \\
&= -3\log 2 + 2\log 3 = \log(2)^{-3} + \log(3)^2 \\
&= \log \frac{1}{2^3} + \log 9 = \log \frac{1}{8} + \log 9 \\
&= \log \left(\frac{1}{8} \times 9 \right) = \log \frac{9}{8}
\end{aligned}$$

④ Evaluate $\int \frac{x^2+1}{(x-3)(x-2)^2} dx$.

Sol: Consider, $\frac{x^2+1}{(x-3)(x-2)^2} = \frac{A}{x-3} + \frac{B}{x-2} + \frac{C}{(x-2)^2}$

$$x^2+1 = A(x-2)^2 + B(x-3)(x-2) + C(x-3)$$

Put $x=2$

$$5 = C(-1)$$

$$-C = 5 \Rightarrow \boxed{C = -5}$$

Put $x=3$

$$\boxed{10 = A}$$

Put $x=0$

$$1 = 4A + 6B - 3C$$

$$1 = 40 + 6B + 15 \Rightarrow 6B = -54$$

$$\boxed{B = -9}$$

$$\therefore \frac{x^2+1}{(x-3)(x-2)^2} = \frac{10}{x-3} - \frac{9}{x-2} - \frac{5}{(x-2)^2}$$

$$\therefore \int \frac{x^2+1}{(x-3)(x-2)^2} dx = 10 \int \frac{dx}{x-3} - 9 \int \frac{dx}{x-2} - 5 \int \frac{dx}{(x-2)^2}$$

$$= 10 \log(x-3) - 9 \log(x-2) - 5 \int (x-2)^{-2} dx$$

$$= 10 \log(x-3) - 9 \log(x-2) - 5 \left[\frac{(x-2)^{-2+1}}{-2+1} \right] + C$$

$$= 10 \log(x-3) - 9 \log(x-2) + 5 \frac{1}{x-2} + C$$

④ Evaluate $\int \frac{2x^2-x+4}{x^3+4x} dx$.

Sol: Consider, $\frac{2x^2-x+4}{x^3+4x} = \frac{2x^2-x+4}{x(x^2+4)} = \frac{A}{x} + \frac{Bx+C}{x^2+4}$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + c)x$$

Put $x=0$

$$4 = 4A \Rightarrow \boxed{A=1}$$

Put $x=1$

$$2 - 1 + 4 = 5A + B + c$$

$$5 = 5 + B + c$$

$$\Rightarrow B + c = 0 \text{ --- (1)}$$

Put $x=-1$

$$2 + 1 + 4 = 5A - (B(-1) + c)$$

$$7 = 5 + B - c$$

$$B - c = 2 \text{ --- (2)}$$

$$\text{(1) + (2)} \Rightarrow 2B = 2 \Rightarrow \boxed{B=1}$$

Subst. $B=1$ in (1), $1 + c = 0 \Rightarrow \boxed{c=-1}$

$$\begin{aligned} \therefore \int \frac{2x^2 - x + 4}{x^2 + 4x} dx &= \int \frac{1}{x} dx + \int \frac{x-1}{x^2+4} dx \\ &= \log x + \int \frac{x}{x^2+4} dx - \int \frac{dx}{x^2+4} \\ &= \log x + \int \frac{du/2}{u} - \frac{1}{2} \tan^{-1} \frac{x}{2} \\ &= \log x + \frac{1}{2} \log u - \frac{1}{2} \tan^{-1} \frac{x}{2} + c \\ &= \log x + \frac{1}{2} \log(x^2+4) - \frac{1}{2} \tan^{-1} \frac{x}{2} + c \end{aligned}$$

Put $x^2+4 = u$
 $2x dx = du$
 $x dx = \frac{du}{2}$

(42) Evaluate $\int \frac{x^2}{x+2} dx$.

Sol:

$x+2$	$x-2$
x^2	x^2
$x^2 + 2x$	$x^2 + 2x$
$(-)(-)$	$(-)(-)$
$-2x$	$-2x$
$-2x - 4$	$-2x - 4$
$(+)(+)$	$(+)(+)$
4	4

$$\frac{x^2}{x+2} = x - 2 + \frac{4}{x+2}$$

$$\therefore \int \frac{x^2}{x+2} dx = \int \left(x - 2 + \frac{4}{x+2} \right) dx = \frac{x^2}{2} - 2x + 4 \log(x+2) + c$$

Working rule: $\int \frac{px+q}{\sqrt{ax^2+bx+c}} dx$

Put $px+q = A \frac{d}{dx}(ax^2+bx+c) + B$

(A0) (43) Evaluate $\int \frac{2x+5}{\sqrt{x^2-2x+10}} dx$.

Sol: Put $2x+5 = A \frac{d}{dx}(x^2-2x+10) + B$

$$2x+5 = A(2x-2) + B \Rightarrow 2x+5 = 2Ax - 2A + B$$

Equating like coefficients on both sides, we get

$$2 = 2A \Rightarrow \boxed{A=1}$$

$$5 = -2A + B \Rightarrow 5 = -2 + B \Rightarrow \boxed{B=7}$$

$$\therefore 2x+5 = (2x-2) + 7$$

$$\int \frac{2x+5}{\sqrt{x^2-2x+10}} dx = \int \frac{2x-2}{\sqrt{x^2-2x+10}} dx + \int \frac{7}{\sqrt{x^2-2x+10}} dx$$

Put $u = x^2 - 2x + 10$
 $du = (2x-2) dx$

$$= \int \frac{du}{\sqrt{u}} + 7 \int \frac{dx}{\sqrt{x^2-2x+1-1+10}}$$

$$= \int u^{-1/2} du + 7 \int \frac{dx}{\sqrt{(x-1)^2 + 9}}$$

Put $t = x-1$
 $dt = dx$

$$= \frac{u^{-1/2+1}}{-1/2+1} + 7 \int \frac{dt}{\sqrt{t^2+3^2}}$$

$$= \frac{u^{1/2}}{1/2} + 7 \sinh^{-1} \frac{t}{3} + C$$

$$= 2\sqrt{x^2-2x+10} + 7 \sinh^{-1} \left(\frac{x-1}{3} \right) + C$$

$$\int \frac{dx}{\sqrt{a^2-x^2}} = \sin^{-1} \frac{x}{a} + C$$

$$\int \frac{dx}{\sqrt{x^2-a^2}} = \cosh^{-1} \frac{x}{a} + C$$

$$\int \frac{dx}{\sqrt{a^2+x^2}} = \sinh^{-1} \frac{x}{a} + C$$

(A0) (44) Evaluate $\int \frac{x}{\sqrt{x^2+x+1}} dx$

Sol: Put $x = A \frac{d}{dx}(x^2+x+1) + B$

$$x = A(2x+1) + B \Rightarrow x = 2Ax + A + B$$

Equating like coefficients on both sides, we get

$$1 = 2A \Rightarrow \boxed{A=1/2}$$

$$0 = A + B \Rightarrow 0 = 1/2 + B \Rightarrow \boxed{B=-1/2}$$

$$\therefore x = 1/2(2x+1) - 1/2$$

$$\begin{aligned} \therefore \int \frac{x}{\sqrt{x^2+x+1}} dx &= \frac{1}{2} \int \frac{2x+1}{\sqrt{x^2+x+1}} dx - \frac{1}{2} \int \frac{dx}{\sqrt{x^2+x+1}} \\ &= \frac{1}{2} \int \frac{du}{\sqrt{u}} - \frac{1}{2} \int \frac{dx}{\sqrt{x^2+2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1}} \\ &= \frac{1}{2} \int u^{-1/2} du - \frac{1}{2} \int \frac{dx}{\sqrt{(x+1/2)^2 + 3/4}} \\ &= \frac{1}{2} \frac{u^{-1/2+1}}{-1/2+1} - \frac{1}{2} \int \frac{dt}{\sqrt{t^2 + (\frac{\sqrt{3}}{2})^2}} \\ &= \frac{1}{2} \frac{u^{1/2}}{1/2} - \frac{1}{2} \sinh^{-1} \frac{t}{\sqrt{3/2}} + c \\ &= \sqrt{x^2+x+1} - \frac{1}{2} \sinh^{-1} \left(\frac{2}{\sqrt{3}} (x+1/2) \right) + c \end{aligned}$$

Put $u = x^2 + x + 1$
 $du = (2x+1) dx$

Put $t = x + 1/2$
 $dt = dx$

(A0) 45 Evaluate $\int_3^{\infty} \frac{dx}{(x-2)^{3/2}}$ & determine whether it is convergent or divergent.

Sol: $\int_3^{\infty} \frac{dx}{(x-2)^{3/2}} = \lim_{t \rightarrow \infty} \int_3^t \frac{dx}{(x-2)^{3/2}} = \lim_{t \rightarrow \infty} \int_3^t (x-2)^{-3/2} dx$

$$= \lim_{t \rightarrow \infty} \left[\frac{(x-2)^{-3/2+1}}{-3/2+1} \right]_3^t = \lim_{t \rightarrow \infty} \left[\frac{(x-2)^{-1/2}}{-1/2} \right]_3^t$$

$$= \lim_{t \rightarrow \infty} \left[-2 \frac{1}{\sqrt{x-2}} \right]_3^t = \lim_{t \rightarrow \infty} \left[\frac{-2}{\sqrt{t-2}} + \frac{2}{\sqrt{1}} \right]$$

$$= \frac{-2}{\infty} + 2 = 0 + 2 = 2$$

$\therefore \int_3^{\infty} \frac{dx}{(x-2)^{3/2}}$ is convergent.

(A0) 46 Evaluate $\int_4^{\infty} \frac{1}{\sqrt{x}} dx$ & determine whether it is convergent or divergent.

Sol: $\int_4^{\infty} \frac{1}{\sqrt{x}} dx = \lim_{t \rightarrow \infty} \int_4^t x^{-1/2} dx = \lim_{t \rightarrow \infty} \left[\frac{x^{-1/2+1}}{-1/2+1} \right]_4^t$

$$= \lim_{t \rightarrow \infty} \left[\frac{x^{1/2}}{1/2} \right]_4^t = \lim_{t \rightarrow \infty} \left[2\sqrt{x} \right]_4^t = \lim_{t \rightarrow \infty} \left[2\sqrt{t} - 2\sqrt{4} \right]$$

$$= \lim_{t \rightarrow \infty} \left[2\sqrt{t} - 4 \right] = \infty - 4 = \infty$$

$\therefore \int_4^{\infty} \frac{1}{\sqrt{x}} dx$ is divergent.

(AU) (47) Determine whether the given integral $\int_0^{\infty} e^x dx$ is convergent or divergent.

Sol: $\int_0^{\infty} e^x dx = \lim_{t \rightarrow \infty} \int_0^t e^x dx = \lim_{t \rightarrow \infty} (e^x)_0^t = \lim_{t \rightarrow \infty} (e^t - e^0)$
 $= \lim_{t \rightarrow \infty} (e^t - 1) = e^{\infty} - 1 = \infty - 1 = \infty$

$\therefore \int_0^{\infty} e^x dx$ is divergent.

(AU) (48) For what values of p is $\int_1^{\infty} \frac{1}{x^p} dx$ convergent?

Sol: If $p \neq 1$,

$$\lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \int_1^t x^{-p} dx = \lim_{t \rightarrow \infty} \left[\frac{x^{-p+1}}{-p+1} \right]_1^t$$

$$= \lim_{t \rightarrow \infty} \left[\frac{t^{-p+1}}{-p+1} - \frac{1^{-p+1}}{-p+1} \right]$$

$$= \lim_{t \rightarrow \infty} \left[\frac{t^{-p+1}}{-p+1} - \frac{1}{-p+1} \right] = \lim_{t \rightarrow \infty} \left[\frac{1}{1-p} (t^{-p+1} - 1) \right]$$

$$= \lim_{t \rightarrow \infty} \left[\frac{1}{1-p} (t^{-(p-1)} - 1) \right] = \lim_{t \rightarrow \infty} \left[\frac{1}{1-p} \left(\frac{1}{t^{p-1}} - 1 \right) \right]$$

$$= \frac{1}{1-p} \left[\frac{1}{\infty} - 1 \right] = \frac{1}{1-p} [0 - 1] = \frac{-1}{1-p} = \frac{1}{p-1} \rightarrow \text{Rough work}$$

$$= \lim_{t \rightarrow \infty} \left[\frac{1}{p-1} \left(1 - \frac{1}{t^{p-1}} \right) \right]$$

$$= \begin{cases} \frac{1}{p-1}, & p > 1, \text{ converges} \\ \infty, & p \leq 1, \text{ diverges} \end{cases}$$

MULTIPLE INTEGRALS

(AV) ① Evaluate $\int_1^a \int_2^b \frac{dx dy}{xy}$.

Sol: $\int_1^a \int_2^b \frac{dx dy}{xy} = \int_1^a \int_2^b \frac{dx}{x} \frac{dy}{y} = \int_1^a (\log x)_2^b \frac{dy}{y}$

$= \int_1^a (\log b - \log 2) \frac{dy}{y}$

$= (\log b - \log 2) (\log y)_1^a = (\log b - \log 2) (\log a - \log 1)$

$= (\log b - \log 2) \log a = \log\left(\frac{b}{2}\right) \log a$

(AV) ② Find the value of $\int_0^\infty \int_0^y \left(\frac{e^{-y}}{y}\right) dx dy$.

Sol: $\int_0^\infty \int_0^y \left(\frac{e^{-y}}{y}\right) dx dy = \int_0^\infty \left(\frac{e^{-y}}{y}\right) (x)_0^y dy$

$= \int_0^\infty \left(\frac{e^{-y}}{y}\right) \times y dy = \int_0^\infty e^{-y} dy$

$= \left(\frac{e^{-y}}{-1}\right)_0^\infty = -(e^{-y})_0^\infty = -(e^{-\infty} - e^{-0})$

$= -(0 - 1) = 1$

(AV) ③ Evaluate $\int_1^{\ln 8} \int_0^{\ln y} e^{x+y} dx dy$.

Sol: $\int_1^{\ln 8} \int_0^{\ln y} e^{x+y} dx dy = \int_1^{\ln 8} \int_0^{\ln y} e^x e^y dx dy = \int_1^{\ln 8} (e^x)_0^{\ln y} e^y dy$

$= \int_1^{\ln 8} (e^{\ln y} - e^0) e^y dy = \int_1^{\ln 8} (y - 1) e^y dy$

$= \int_1^{\ln 8} (ye^y - e^y) dy$

$= (ye^y)_1^{\ln 8} - (e^y)_1^{\ln 8}$

$= \ln 8 e^{\ln 8} - e - (e^{\ln 8})_1^{\ln 8} - (e^{\ln 8} - e)$

$u = y$	$dv = e^y dy$
$du = dy$	$v = e^y$
$\int u dv = uv - \int v du$	

$$= \ln 8 \cdot 8 - e - (e^{\ln 8} - e) - (8 - e)$$

$$= 8 \ln 8 - e - 8 + e - 8 + e = 8 \ln 8 + e - 16$$

(10) (4) Evaluate $\int_1^2 \int_0^{x^2} x \, dx \, dy$

Sol: $\int_1^2 \int_0^{x^2} x \, dx \, dy = \int_{x=1}^2 \int_{y=0}^{x^2} x \, dy \, dx$ (correct form)

$$= \int_1^2 x (y)_0^{x^2} \, dx = \int_1^2 x (x^2 - 0) \, dx = \int_1^2 x^3 \, dx$$

$$= \left(\frac{x^4}{4} \right)_1^2 = \frac{2^4}{4} - \frac{1}{4} = \frac{16}{4} - \frac{1}{4} = \frac{15}{4}$$

(10) (5) Evaluate $\int_0^{2a} \int_0^x \int_y^x xyz \, dz \, dy \, dx$.

Sol: $\int_0^{2a} \int_0^x \int_y^x xyz \, dz \, dy \, dx = \int_0^{2a} \int_0^x xy \left(\frac{z^2}{2} \right)_y^x \, dy \, dx$

$$= \int_0^{2a} \int_0^x \frac{xy}{2} (x^2 - y^2) \, dy \, dx$$

$$= \frac{1}{2} \int_0^{2a} \int_0^x x (yx^2 - y^3) \, dy \, dx$$

$$= \frac{1}{2} \int_0^{2a} x \left(\frac{y^2 x^2}{2} - \frac{y^4}{4} \right)_0^x \, dx$$

$$= \frac{1}{2} \int_0^{2a} x \left(\frac{x^4}{2} - \frac{x^4}{4} \right) \, dx$$

$$= \frac{1}{2} \int_0^{2a} x \left(\frac{2x^4 - x^4}{4} \right) \, dx = \frac{1}{2} \int_0^{2a} x \left(\frac{x^4}{4} \right) \, dx$$

$$= \frac{1}{8} \int_0^{2a} x^5 \, dx = \frac{1}{8} \left(\frac{x^6}{6} \right)_0^{2a} = \frac{1}{48} (x^6)_0^{2a}$$

$$= \frac{1}{48} ((2a)^6 - 0) = \frac{64a^6}{48} = \frac{4}{3} a^6$$


(6) Find the limits of integration $\iint_R f(x,y) dx dy$ where R is the triangle bounded by $x=0, y=0, x+y=2$.

Sol: Given $x=0, y=0, x+y=2$
 $x+y=2 \Rightarrow y=2-x$

$x:$	-2	-1	0	1	2
$y:$	4	3	2	1	0

From the graph, we get
 $x=0, x=2-y$ & $y=0, y=2$

$$\therefore \iint_R f(x,y) dx dy = \int_0^2 \int_0^{2-y} f(x,y) dx dy$$




(7) Find the limits of integration in the double integral $\iint_R f(x,y) dx dy$ where R is in the first quadrant & bounded $x=1, y=0, y^2=4x$.

Sol: Given $x=1, y=0, y^2=4x$
 $y^2=4x \Rightarrow y = \pm \sqrt{4x} = \pm 2\sqrt{x}$

$x:$	0	1	4
$y:$	0	± 2	± 4

From the graph, we get
 $x = \frac{y^2}{4}, x=1$ & $y=0, y=2$

$$\therefore \iint_R f(x,y) dx dy = \int_0^2 \int_{\frac{y^2}{4}}^1 f(x,y) dx dy$$



Change the order of integration:

(8) Change the order of integration in $\int_0^1 \int_{y^2}^y f(x,y) dx dy$.

Sol: Given $y=0, y=1, x=y^2$ & $x=y$.

$x=y$

$x:$	-2	-1	0	1	2
$y:$	-2	-1	0	1	2


$$\underline{x=y^2} \rightarrow y=\pm\sqrt{x}$$

x	0	1	4
y	0	± 1	± 2

From the graph, we get

$$y=\sqrt{x}, y=x \text{ \& } x=0, x=1.$$

$$\therefore \int_0^1 \int_{y^2}^y f(x,y) dx dy = \int_0^1 \int_{\sqrt{x}}^x f(x,y) dy dx$$



10) Change the order of integration in $\int_{x=0}^{\infty} \int_{y=x}^{\infty} \frac{e^{-y}}{y} dy dx$ & then evaluate it.

Sol: Given $x=0, x=\infty, y=x$ & $y=\infty$.

From the graph, we get


$$x=0, x=y \text{ \& } y=0, y=\infty.$$

$$\therefore \int_0^{\infty} \int_x^{\infty} \frac{e^{-y}}{y} dy dx = \int_0^{\infty} \int_0^y \frac{e^{-y}}{y} dx dy$$

$$= \int_0^{\infty} \left(\frac{e^{-y}}{y}\right) (x)_0^y dy$$

$$= \int_0^{\infty} \left(\frac{e^{-y}}{y}\right) (y) dy = \int_0^{\infty} e^{-y} dy = \left(\frac{e^{-y}}{-1}\right)_0^{\infty}$$

$$= -(e^{-y})_0^{\infty} = -(e^{-\infty} - e^{-0}) = -(0 - 1) = 1$$




10) Change the order of integration for the given integral $\int_{x=0}^a \int_0^{2\sqrt{ax}} x^2 dy dx$ & evaluate it.

Sol: Given $x=0, x=a, y=0$ & $y=2\sqrt{ax}$

$$\underline{y=2\sqrt{ax}} \Rightarrow y^2=4ax \Rightarrow x=\frac{y^2}{4a}$$

x	0	a	a/4
y	0	2a	a



From the graph, we get

$$x = \frac{y^2}{4a}, \quad x=a \quad \& \quad y=0, \quad y=2a$$

$$\begin{aligned} \therefore \int_0^a \int_0^{2\sqrt{ax}} x^2 dy dx &= \int_0^{2a} \int_{\frac{y^2}{4a}}^a x^2 dx dy \\ &= \int_0^{2a} \left(\frac{x^3}{3} \right)_{\frac{y^2}{4a}}^a dy = \frac{1}{3} \int_0^{2a} \left(a^3 - \left(\frac{y^2}{4a} \right)^3 \right) dy \\ &= \frac{1}{3} \int_0^{2a} \left(a^3 - \frac{y^6}{64a^3} \right) dy = \frac{1}{3} \left[a^3 y - \frac{y^7}{7 \times 64 a^3} \right]_0^{2a} \\ &= \frac{1}{3} \left[a^3(2a) - \frac{(2a)^7}{7 \times 64 a^3} \right] \\ &= \frac{1}{3} \left[2a^4 - \frac{2 \times 64 a^7}{7 \times 64 a^3} \right] = \frac{1}{3} \left[2a^4 - \frac{2a^4}{7} \right] \\ &= \frac{a^4}{3} \left(2 - \frac{2}{7} \right) = \frac{a^4}{3} \left(\frac{14-2}{7} \right) = \frac{a^4}{3} \left(\frac{12}{7} \right) = \frac{4}{7} a^4 \end{aligned}$$

Q11) Change the order of integration for the given integral $\int_{x=0}^a \int_{y=x/a}^{\sqrt{x/a}} (x^2+y^2) dy dx$ & evaluate it.


Sol: Given $x=0, x=a, y=\frac{x}{a}$ & $y=\sqrt{\frac{x}{a}}$

$y = \frac{x}{a}$

x	0	a	2a	-a	-2a
y	0	1	2	-1	-2

$y = \sqrt{\frac{x}{a}}$

x	0	a	4a
y	0	1	2



From the graph, we have

$$y=0, y=1, x=ay^2 \quad \& \quad x=ay$$

$$\therefore \int_0^1 \int_{ay^2}^{ay} (x^2+y^2) dx dy$$

$$\begin{aligned}
 &= \int_0^1 \left(\frac{x^3}{3} + xy^2 \right)_{ay^2}^{ay} dy \\
 &= \int_0^1 \left(\frac{a^3 y^3}{3} + ay^2 - \frac{a^3 y^6}{3} - ay^2 y^2 \right) dy \\
 &= \int_0^1 \left(\frac{a^3 y^3}{3} + ay^3 - \frac{a^3 y^6}{3} - ay^4 \right) dy \\
 &= \left(\frac{a^3 y^4}{12} - \frac{ay^4}{4} - \frac{a^3 y^7}{21} - \frac{ay^5}{5} \right)_0^1 = \frac{a^3}{12} - \frac{a}{4} - \frac{a^3}{21} - \frac{a}{5} \\
 &= a^3 \left(\frac{1}{12} - \frac{1}{21} \right) - a \left(\frac{1}{4} + \frac{1}{5} \right) = \frac{a^3}{28} - \frac{9a}{20} \\
 &= \frac{a}{4} \left(\frac{a^2}{7} - \frac{9}{5} \right)
 \end{aligned}$$

(12) Change the order of integration & hence evaluate $\int_0^1 \int_{x^2}^{2-x} xy \, dy \, dx$.

Sol:


Given $x=0, x=1, y=x^2$ & $y=2-x$.

$y=x^2$

x	-2	-1	0	1	2
y	4	1	0	1	4

$y=2-x$

x	-2	-1	0	1	2
y	4	3	2	1	0



From the graph, we get


I_1 : $x=0, x=\sqrt{y}, y=0$ & $y=1$

I_2 : $x=0, x=2-y, y=1$ & $y=2$

$$\begin{aligned}
 \int_0^1 \int_{x^2}^{2-x} xy \, dy \, dx &= \int_0^1 \int_0^{\sqrt{y}} xy \, dx \, dy + \int_1^2 \int_0^{2-y} xy \, dx \, dy \\
 &= \int_0^1 \left(\frac{x^2}{2} \right)_0^{\sqrt{y}} y \, dy + \int_1^2 \left(\frac{x^2}{2} \right)_0^{2-y} y \, dy
 \end{aligned}$$

$$\begin{aligned}
&= \int_0^1 \frac{y}{2} y dy + \int_1^2 \frac{(2-y)^2}{2} y dy \\
&= \frac{1}{2} \int_0^1 y^2 dy + \frac{1}{2} \int_1^2 (4 + y^2 - 4y) y dy \\
&= \frac{1}{2} \left(\frac{y^3}{3} \right)_0^1 + \frac{1}{2} \int_1^2 (4y + y^3 - 4y^2) dy \\
&= \frac{1}{2} \left(\frac{1}{3} \right) + \frac{1}{2} \left[\frac{4y^2}{2} + \frac{y^4}{4} - \frac{4y^3}{3} \right]_1^2 \\
&= \frac{1}{6} + \frac{1}{2} \left[\frac{16}{2} + \frac{16}{4} - \frac{32}{3} - \left(\frac{4}{2} + \frac{1}{4} - \frac{4}{3} \right) \right] \\
&= \frac{1}{6} + \frac{1}{2} \left[8 + 4 - \frac{32}{3} - 2 - \frac{1}{4} + \frac{4}{3} \right] = \frac{3}{8}
\end{aligned}$$

(13) Evaluate $\iint xy dx dy$ over the region in the positive quadrant bounded by $\frac{x}{a} + \frac{y}{b} = 1$.



Sol: Given $x=0, y=0, \frac{x}{a} + \frac{y}{b} = 1$.

From the graph, we get

$$\begin{aligned}
x=0, \quad \frac{x}{a} + \frac{y}{b} = 1 &\Rightarrow \frac{x}{a} = 1 - \frac{y}{b} \Rightarrow x = a \left(1 - \frac{y}{b} \right) \\
&\Rightarrow x=0, \quad x = a \left(1 - \frac{y}{b} \right)
\end{aligned}$$

$$\begin{aligned}
\therefore \int_0^b \int_0^{a(1-\frac{y}{b})} xy dx dy &= \int_0^b \left(\frac{x^2}{2} \right)_0^{a(1-\frac{y}{b})} y dy \\
&= \frac{1}{2} \int_0^b a^2 \left(1 - \frac{y}{b} \right)^2 y dy = \frac{a^2}{2} \int_0^b \left(1 + \frac{y^2}{b^2} - \frac{2y}{b} \right) y dy \\
&= \frac{a^2}{2} \int_0^b \left[y + \frac{y^3}{b^2} - \frac{2y^2}{b} \right] dy \\
&= \frac{a^2}{2} \left[\frac{y^2}{2} + \frac{y^4}{4b^2} - \frac{2y^3}{3b} \right]_0^b = \frac{a^2}{2} \left[\frac{b^2}{2} + \frac{b^4}{4b^2} - \frac{2b^3}{3b} \right] \\
&= \frac{a^2}{2} \left[\frac{b^2}{2} + \frac{b^2}{4} - \frac{2b^2}{3} \right] = \frac{a^2 b^2}{2} \left(\frac{1}{2} + \frac{1}{4} - \frac{2}{3} \right) = \frac{a^2 b^2}{24}
\end{aligned}$$

14) Using double integral, find the area bounded by $y=x$ & $y=x^2$.


Sol: Given $y=x$ & $y=x^2$.

$y=x$

x	-2	-1	0	1	2
y	-2	-1	0	1	2

$y=x^2$

x	-2	-1	0	1	2
y	4	1	0	1	4



From the graph, we get

$y=x, y=x^2, x=0$ & $x=1$

$$\int_0^1 \int_x^{x^2} dy dx = \int_0^1 (y)_x^{x^2} dx$$

$$= \int_0^1 (x^2 - x) dx = \left(\frac{x^3}{3} - \frac{x^2}{2} \right)_0^1 = \frac{1}{3} - \frac{1}{2} = -\frac{1}{6}$$

Hence the required area is $\frac{1}{6}$.

15) Evaluate $\iint xy(x+y) dx dy$ over the area between $y=x^2$ & $y=x$.

Sol: By using Problem no. 14, we have $y=x^2, y=x, x=0$ & $x=1$.

$$\int_0^1 \int_{x^2}^x xy(x+y) dy dx = \int_0^1 \int_{x^2}^x x(xy+y^2) dy dx$$

$$= \int_0^1 x \left(x \frac{y^2}{2} + \frac{y^3}{3} \right)_{x^2}^x dx$$

$$= \int_0^1 x \left(\frac{x^3}{2} + \frac{x^3}{3} - \frac{x^5}{2} - \frac{x^6}{3} \right) dx$$

$$= \int_0^1 x \left(\frac{5x^3}{6} - \frac{x^5}{2} - \frac{x^6}{3} \right) dx$$

$$= \int_0^1 \left(\frac{5x^4}{6} - \frac{x^6}{2} - \frac{x^7}{3} \right) dx$$

$$= \left(\frac{5x^5}{30} - \frac{x^7}{14} - \frac{x^8}{24} \right)_0^1 = \frac{5}{30} - \frac{1}{14} - \frac{1}{24}$$

$$= \frac{3}{56}$$